



Full wwPDB X-ray Structure Validation Report i

Oct 15, 2023 – 01:56 PM EDT

PDB ID : 2DEX
Title : Crystal structure of human peptidylarginine deiminase 4 in complex with histone H3 N-terminal peptide including Arg17
Authors : Arita, K.; Shimizu, T.; Hashimoto, H.; Hidaka, Y.; Yamada, M.; Sato, M.
Deposited on : 2006-02-18
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 5190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein-arginine deiminase type IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	X	627	4937	3153	828	922	34	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	-7	GLY	-	cloning artifact	UNP Q9UM07
X	-6	PRO	-	cloning artifact	UNP Q9UM07
X	-5	LEU	-	cloning artifact	UNP Q9UM07
X	-4	GLY	-	cloning artifact	UNP Q9UM07
X	-3	SER	-	cloning artifact	UNP Q9UM07
X	-2	PRO	-	cloning artifact	UNP Q9UM07
X	-1	GLU	-	cloning artifact	UNP Q9UM07
X	0	PHE	-	cloning artifact	UNP Q9UM07
X	645	ALA	CYS	engineered mutation	UNP Q9UM07

- Molecule 2 is a protein called 10-mer peptide from histone H3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	A	5	41	25	10	6	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	X	5	Total Ca 5 5	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	X	1	Total O S 5 4 1	0	0
4	X	1	Total O S 5 4 1	0	0
4	X	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	X	190	Total O 190 190	0	0
5	A	2	Total O 2 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein-arginine deiminase type IV



- Molecule 2: 10-mer peptide from histone H3



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	146.07Å 60.10Å 115.70Å 90.00° 124.32° 90.00°	Depositor
Resolution (Å)	33.13 – 2.10 33.13 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.6 (33.13-2.10) 94.6 (33.13-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.46 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R , R_{free}	0.203 , 0.247 0.210 , 0.250	Depositor DCC
R_{free} test set	4628 reflections (10.05%)	wwPDB-VP
Wilson B-factor (Å ²)	54.4	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 67.5	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5190	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	X	0.57	1/5056 (0.0%)	0.74	7/6861 (0.1%)
2	A	0.58	0/41	0.77	0/53
All	All	0.57	1/5097 (0.0%)	0.74	7/6914 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	311	CYS	CB-SG	-5.93	1.72	1.81

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	650	ARG	NE-CZ-NH1	6.23	123.42	120.30
1	X	650	ARG	NE-CZ-NH2	-6.13	117.23	120.30
1	X	203	VAL	CB-CA-C	-5.94	100.11	111.40
1	X	372	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	X	427	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	X	367	VAL	CB-CA-C	-5.14	101.64	111.40
1	X	201	LEU	CA-CB-CG	5.13	127.10	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	313	ILE	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	4937	0	4900	72	0
2	A	41	0	48	2	0
3	X	5	0	0	0	0
4	X	15	0	0	0	0
5	A	2	0	0	1	0
5	X	190	0	0	7	0
All	All	5190	0	4948	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:454:GLN:O	5:X:909:HOH:O	1.79	1.00
1:X:425:LEU:HD12	1:X:456:VAL:HG13	1.58	0.86
1:X:315:GLU:HG3	1:X:316:ASN:HD22	1.43	0.82
1:X:574:LYS:NZ	1:X:580:GLU:OE1	2.11	0.81
1:X:313:ILE:HB	1:X:338:PRO:HA	1.62	0.79
1:X:572:LYS:HE3	1:X:574:LYS:HZ2	1.47	0.78
1:X:26:GLN:OE1	1:X:77:THR:HB	1.84	0.76
1:X:316:ASN:O	5:X:1097:HOH:O	2.07	0.73
2:A:1:ALA:N	5:A:188:HOH:O	2.23	0.72
1:X:195:THR:HG22	1:X:196:ASN:ND2	2.05	0.72
1:X:572:LYS:HE3	1:X:574:LYS:NZ	2.07	0.70
1:X:236:HIS:ND1	5:X:968:HOH:O	2.26	0.69
1:X:310:ALA:O	1:X:336:ILE:HA	1.95	0.66
1:X:203:VAL:CG2	1:X:229:LEU:HD13	2.27	0.64
1:X:312:SER:HB2	1:X:320:LEU:HD12	1.82	0.62
1:X:662:VAL:O	1:X:662:VAL:HG23	2.00	0.61
1:X:52:ILE:HD12	1:X:76:VAL:HG22	1.81	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:315:GLU:HG3	1:X:316:ASN:ND2	2.16	0.57
1:X:586:MET:HA	1:X:599:PRO:HG2	1.85	0.57
1:X:203:VAL:HG22	1:X:229:LEU:HD13	1.86	0.57
1:X:312:SER:CB	1:X:320:LEU:HD12	2.35	0.57
1:X:594:LYS:NZ	5:X:1009:HOH:O	2.20	0.56
1:X:311:CYS:CB	1:X:337:CYS:HB3	2.36	0.56
1:X:45:SER:OG	1:X:90:GLN:NE2	2.38	0.56
1:X:334:LEU:C	1:X:334:LEU:HD13	2.27	0.55
1:X:27:LEU:HB3	1:X:29:ILE:CD1	2.36	0.55
1:X:123:ARG:O	1:X:124:THR:OG1	2.24	0.54
1:X:307:GLU:OE2	1:X:335:THR:CG2	2.57	0.53
1:X:311:CYS:HB3	1:X:337:CYS:HB3	1.91	0.53
1:X:307:GLU:OE2	1:X:335:THR:HG21	2.09	0.51
1:X:153:ASN:HB3	1:X:166:CYS:HB3	1.93	0.51
1:X:514:LEU:HD13	1:X:515:PHE:CE2	2.45	0.51
1:X:441:ARG:HH11	1:X:441:ARG:HG3	1.76	0.51
1:X:441:ARG:NH1	5:X:921:HOH:O	2.44	0.51
1:X:425:LEU:HD12	1:X:456:VAL:CG1	2.35	0.51
1:X:108:LEU:HG	1:X:110:LEU:HD13	1.93	0.51
1:X:158:ASN:HD22	1:X:159:LEU:N	2.10	0.50
1:X:158:ASN:HD22	1:X:158:ASN:C	2.15	0.50
1:X:123:ARG:HD3	1:X:659:TRP:CD1	2.47	0.49
1:X:166:CYS:HB2	1:X:254:LEU:HD22	1.94	0.49
1:X:186:SER:OG	1:X:246:ASN:ND2	2.46	0.49
1:X:434:CYS:HB2	5:X:961:HOH:O	2.14	0.47
1:X:633:PHE:CE1	1:X:638:ILE:HD11	2.50	0.46
1:X:596:LEU:HD11	1:X:625:LEU:HG	1.98	0.46
1:X:311:CYS:HB2	1:X:342:ASN:OD1	2.15	0.46
1:X:126:LYS:N	5:X:1036:HOH:O	2.47	0.46
1:X:633:PHE:O	1:X:638:ILE:HG12	2.15	0.45
1:X:379:PHE:HB3	1:X:380:PRO:HD3	1.98	0.45
1:X:623:LEU:HB2	1:X:625:LEU:HD22	1.98	0.45
1:X:74:VAL:HG13	1:X:74:VAL:O	2.17	0.44
1:X:313:ILE:CB	1:X:338:PRO:HA	2.40	0.44
1:X:320:LEU:HD23	1:X:320:LEU:O	2.17	0.44
1:X:553:LEU:O	1:X:557:LEU:HB2	2.18	0.44
1:X:427:ARG:HG2	1:X:460:VAL:HG23	2.00	0.44
1:X:71:ASP:O	1:X:74:VAL:HG12	2.18	0.43
1:X:80:MET:CE	1:X:90:GLN:HG3	2.48	0.43
1:X:42:ILE:CD1	1:X:76:VAL:HG11	2.48	0.43
1:X:535:LEU:HD22	1:X:539:ASN:ND2	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:311:CYS:SG	1:X:311:CYS:O	2.77	0.43
1:X:594:LYS:HA	1:X:625:LEU:HD12	2.00	0.43
1:X:296:TRP:CZ3	1:X:298:MET:HG3	2.53	0.43
1:X:198:THR:HG23	1:X:272:LEU:HD12	2.00	0.42
1:X:374:ARG:O	1:X:377:LYS:HB2	2.19	0.42
1:X:427:ARG:HG3	1:X:458:ALA:O	2.19	0.42
1:X:313:ILE:O	1:X:342:ASN:ND2	2.53	0.42
1:X:203:VAL:HG23	1:X:229:LEU:HD13	1.99	0.42
1:X:334:LEU:HD11	1:X:336:ILE:CD1	2.50	0.41
1:X:114:GLU:O	1:X:187:THR:HA	2.20	0.41
1:X:80:MET:HE3	1:X:90:GLN:HG3	2.02	0.41
1:X:232:LYS:HD2	1:X:232:LYS:O	2.20	0.41
1:X:365:PRO:HD2	1:X:388:ASP:O	2.21	0.41
1:X:212:ARG:HD3	1:X:227:VAL:CG1	2.51	0.41
1:X:469:VAL:HG11	2:A:3:ARG:HD3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	617/671 (92%)	591 (96%)	20 (3%)	6 (1%)	15 11
2	A	3/10 (30%)	3 (100%)	0	0	100 100
All	All	620/681 (91%)	594 (96%)	20 (3%)	6 (1%)	15 11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	484	ARG
1	X	124	THR
1	X	315	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	458	ALA
1	X	485	LYS
1	X	519	LYS

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	557/593 (94%)	508 (91%)	49 (9%)	10 6
2	A	4/8 (50%)	4 (100%)	0	100 100
All	All	561/601 (93%)	512 (91%)	49 (9%)	10 7

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	8	ARG
1	X	10	THR
1	X	67	THR
1	X	70	LEU
1	X	77	THR
1	X	87	THR
1	X	110	LEU
1	X	116	SER
1	X	117	LEU
1	X	124	THR
1	X	150	LEU
1	X	158	ASN
1	X	169	ASP
1	X	175	GLU
1	X	185	LEU
1	X	186	SER
1	X	201	LEU
1	X	203	VAL
1	X	232	LYS
1	X	244	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	276	ASN
1	X	314	PHE
1	X	328	MET
1	X	331	LYS
1	X	335	THR
1	X	364	LEU
1	X	376	LEU
1	X	382	LYS
1	X	389	PHE
1	X	392	VAL
1	X	404	LEU
1	X	416	VAL
1	X	427	ARG
1	X	429	LEU
1	X	489	LEU
1	X	500	LEU
1	X	514	LEU
1	X	530	LEU
1	X	535	LEU
1	X	544	ARG
1	X	557	LEU
1	X	570	LEU
1	X	592	LEU
1	X	594	LYS
1	X	606	ILE
1	X	609	ARG
1	X	614	GLU
1	X	625	LEU
1	X	635	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	90	GLN
1	X	158	ASN
1	X	178	GLN
1	X	196	ASN
1	X	246	ASN
1	X	303	GLN
1	X	316	ASN
1	X	346	GLN
1	X	448	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	505	GLN
1	X	538	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	X	907	-	4,4,4	0.15	0	6,6,6	0.16	0
4	SO4	X	906	-	4,4,4	0.13	0	6,6,6	0.30	0
4	SO4	X	905	-	4,4,4	0.12	0	6,6,6	0.40	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	627/671 (93%)	0.72	99 (15%) 2 2	51, 63, 80, 95	0
2	A	5/10 (50%)	2.51	3 (60%) 0 0	67, 71, 75, 77	0
All	All	632/681 (92%)	0.73	102 (16%) 1 2	51, 63, 80, 95	0

All (102) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	312	SER	8.1
1	X	314	PHE	6.5
1	X	587	VAL	5.3
2	A	1	ALA	5.0
1	X	476	LEU	4.9
1	X	74	VAL	4.7
1	X	124	THR	4.6
1	X	520	LYS	4.4
1	X	38	THR	4.3
1	X	410	LEU	4.3
1	X	98	GLY	4.1
1	X	546	ILE	4.0
1	X	217	THR	3.9
1	X	484	ARG	3.8
1	X	328	MET	3.7
1	X	137	ARG	3.7
1	X	47	GLY	3.6
1	X	462	LEU	3.6
1	X	159	LEU	3.6
1	X	491	LEU	3.6
2	A	2	PRO	3.5
1	X	367	VAL	3.5
1	X	475	PHE	3.4
1	X	136	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	354	ILE	3.4
1	X	522	LYS	3.3
1	X	649	VAL	3.3
1	X	4	GLY	3.2
1	X	313	ILE	3.2
1	X	586	MET	3.1
1	X	366	VAL	3.1
1	X	311	CYS	3.1
1	X	567	ILE	3.1
1	X	588	ASN	3.1
1	X	233	TRP	3.0
1	X	126	LYS	3.0
1	X	519	LYS	3.0
1	X	477	SER	3.0
1	X	99	PRO	2.9
1	X	125	GLY	2.9
1	X	585	ASN	2.9
1	X	591	VAL	2.8
1	X	321	LYS	2.8
1	X	315	GLU	2.8
1	X	343	MET	2.8
1	X	52	ILE	2.7
1	X	589	MET	2.7
1	X	161	SER	2.7
1	X	644	HIS	2.7
1	X	68	TRP	2.7
1	X	398	THR	2.7
1	X	490	LEU	2.7
1	X	49	VAL	2.6
1	X	104	VAL	2.6
1	X	400	GLY	2.6
1	X	412	VAL	2.6
1	X	472	VAL	2.6
1	X	81	LYS	2.6
1	X	232	LYS	2.5
1	X	13	GLN	2.5
2	A	5	GLN	2.5
1	X	70	LEU	2.5
1	X	357	ILE	2.5
1	X	51	ASP	2.4
1	X	71	ASP	2.4
1	X	583	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	645	ALA	2.4
1	X	331	LYS	2.4
1	X	467	LEU	2.4
1	X	100	LYS	2.4
1	X	296	TRP	2.4
1	X	473	ASP	2.4
1	X	298	MET	2.3
1	X	598	ILE	2.3
1	X	489	LEU	2.3
1	X	50	VAL	2.3
1	X	79	THR	2.3
1	X	152	VAL	2.3
1	X	162	SER	2.3
1	X	278	GLU	2.2
1	X	344	ASP	2.2
1	X	297	ILE	2.2
1	X	408	GLY	2.2
1	X	146	GLN	2.2
1	X	492	ALA	2.2
1	X	420	GLY	2.1
1	X	599	PRO	2.1
1	X	643	VAL	2.1
1	X	576	PHE	2.1
1	X	355	GLY	2.1
1	X	411	GLU	2.1
1	X	569	GLN	2.1
1	X	316	ASN	2.1
1	X	353	GLU	2.1
1	X	438	ASN	2.1
1	X	144	CYS	2.1
1	X	333	LYS	2.1
1	X	428	ILE	2.1
1	X	172	LEU	2.0
1	X	590	LEU	2.0
1	X	160	GLU	2.0
1	X	349	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	X	904	1/1	0.94	0.08	62,62,62,62	0
4	SO4	X	907	5/5	0.95	0.24	87,88,88,89	0
4	SO4	X	905	5/5	0.96	0.12	73,73,74,75	0
4	SO4	X	906	5/5	0.97	0.16	86,86,87,87	0
3	CA	X	902	1/1	0.97	0.07	65,65,65,65	0
3	CA	X	901	1/1	0.98	0.04	61,61,61,61	0
3	CA	X	903	1/1	0.98	0.06	62,62,62,62	0
3	CA	X	900	1/1	1.00	0.10	53,53,53,53	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.