



Full wwPDB X-ray Structure Validation Report i

Dec 4, 2023 – 07:53 am GMT

PDB ID : 2JIO

Title : A New Catalytic Mechanism of Periplasmic Nitrate Reductase from Desulfovibrio desulfuricans ATCC 27774 from Crystallographic and EPR Data and based on detailed analysis of the sixth ligand

Authors : Najmudin, S.; Gonzalez, P.J.; Trincao, J.; Coelho, C.; Mukhopadhyay, A.; Romao, C.C.; Moura, I.; Moura, J.J.G.; Brondino, C.D.; Romao, M.J.

Deposited on : 2007-06-28

Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references \(1\)](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for $>=3$, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	723	 87% 12% .

2 Entry composition [\(i\)](#)


There are 6 unique types of molecules in this entry. The entry contains 6580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PERIPLASMIC NITRATE REDUCTASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	720	Total	C 5666	N 3573	O 1019	S 1033	41	0

- Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe 8	S 4	0	0

- Molecule 3 is MOLYBDENUM ATOM (three-letter code: MO) (formula: Mo).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mo 1	0	0

- Molecule 4 is 2-AMINO-5,6-DIMERCAPTO-7-METHYL-3,7,8A,9-TETRAHYDRO-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-4-ONE GUANOSINE DINUCLEOTIDE

(three-letter code: MGD) (formula: C₂₀H₂₆N₁₀O₁₃P₂S₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
4	A	1	47	20	10	13	2	2	0	0
4	A	1	Total	C	N	O	P	S	0	0
			47	20	10	13	2	2		

- Molecule 5 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total X 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	810	Total O 810 810	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PERIPLASMIC NITRATE REDUCTASE



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.19 Å 106.19 Å 135.13 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	92.06 – 2.20 76.03 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (92.06-2.20) 99.4 (76.03-2.20)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.06 (at 2.20 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.150 , 0.209 0.164 , 0.216	Depositor DCC
R_{free} test set	2267 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.676	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6580	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MO, SF4, MGD, UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/5868	0.78	14/7959 (0.2%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	335	ARG	NE-CZ-NH2	-8.98	115.81	120.30
1	A	702[A]	ASP	CB-CA-C	-8.91	92.58	110.40
1	A	702[B]	ASP	CB-CA-C	-8.91	92.58	110.40
1	A	138	ARG	NE-CZ-NH2	-7.35	116.63	120.30
1	A	335	ARG	NE-CZ-NH1	6.90	123.75	120.30
1	A	580	ARG	NE-CZ-NH2	-6.71	116.94	120.30
1	A	67	LEU	CA-CB-CG	6.26	129.69	115.30
1	A	525	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	A	580	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	A	666	ARG	NE-CZ-NH1	-5.66	117.47	120.30
1	A	477	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	666	ARG	NE-CZ-NH2	5.16	122.88	120.30
1	A	76	LEU	CA-CB-CG	5.15	127.15	115.30
1	A	525	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5666	0	5568	86	0
2	A	8	0	0	0	0
3	A	1	0	0	0	0
4	A	94	0	44	12	0
5	A	1	0	0	0	0
6	A	810	0	0	18	0
All	All	6580	0	5612	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:540[A]:ARG:NH1	6:A:2615:HOH:O	1.84	1.07
1:A:168:THR:HG22	1:A:299:SER:HB3	1.56	0.86
1:A:373:LYS:HE3	6:A:2451:HOH:O	1.77	0.85
1:A:541[A]:GLU:OE1	1:A:544:ARG:NH1	2.09	0.84
1:A:714:LYS:HE3	4:A:812:MGD:C5'	2.09	0.82
1:A:368:ALA:H	1:A:374:HIS:HD2	1.29	0.80
1:A:714:LYS:HE3	4:A:812:MGD:H5'1	1.68	0.75
1:A:121:ASN:HD21	1:A:125:LYS:NZ	1.89	0.71
1:A:597:PRO:HD2	6:A:2687:HOH:O	1.89	0.71
1:A:368:ALA:H	1:A:374:HIS:CD2	2.10	0.70
1:A:714:LYS:HE3	4:A:812:MGD:H5'2	1.74	0.68
1:A:443:ILE:O	4:A:811:MGD:N2	2.26	0.66
1:A:121:ASN:HD22	1:A:133:VAL:HG11	1.59	0.66
1:A:168:THR:HG21	6:A:2414:HOH:O	1.95	0.65
1:A:617:ARG:HH22	4:A:811:MGD:H15	1.45	0.65
1:A:345:GLY:HA3	4:A:812:MGD:C12	2.29	0.63
1:A:237:ILE:HD11	1:A:329:ILE:HG12	1.82	0.61
1:A:580:ARG:HD3	6:A:2338:HOH:O	1.99	0.61
1:A:617:ARG:HH21	1:A:623:HIS:CE1	2.17	0.61
1:A:11:GLY:HA2	1:A:481:THR:HA	1.83	0.61
1:A:15:TYR:O	1:A:347:PRO:HD3	2.01	0.61
1:A:617:ARG:HH21	1:A:623:HIS:HE1	1.50	0.59
1:A:134:ASP:OD1	1:A:138:ARG:HD2	2.04	0.58
1:A:580:ARG:NH2	6:A:2667:HOH:O	2.38	0.56
1:A:666:ARG:NH1	6:A:2757:HOH:O	2.29	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:623:HIS:CE1	4:A:812:MGD:H15	2.23	0.56
1:A:381:LEU:HD22	1:A:526:MET:CE	2.37	0.55
1:A:101:PRO:HB2	1:A:130:THR:HB	1.89	0.54
1:A:424:PRO:HG2	1:A:702[A]:ASP:OD1	2.08	0.54
1:A:714:LYS:CE	4:A:812:MGD:H5'2	2.39	0.53
1:A:657:HIS:CD2	1:A:676:VAL:H	2.27	0.53
1:A:371:ASN:OD1	1:A:373:LYS:HE2	2.09	0.52
1:A:540[A]:ARG:HH11	1:A:540[A]:ARG:HG3	1.74	0.52
1:A:540[A]:ARG:HH11	1:A:540[A]:ARG:CG	2.23	0.51
1:A:657:HIS:HD2	1:A:676:VAL:H	1.58	0.51
1:A:545:LYS:NZ	6:A:2621:HOH:O	2.44	0.49
1:A:545:LYS:HD2	6:A:2623:HOH:O	2.12	0.49
1:A:617:ARG:HD2	4:A:812:MGD:H102	1.95	0.49
1:A:136:ASN:OD1	1:A:137:PRO:HD3	2.13	0.49
1:A:121:ASN:HD21	1:A:125:LYS:HZ1	1.59	0.49
1:A:649:ASP:O	1:A:653:THR:HG22	2.13	0.48
1:A:493:PRO:HB2	1:A:496:ASN:HD22	1.78	0.48
1:A:648:GLU:HG2	6:A:2745:HOH:O	2.14	0.48
1:A:657:HIS:HD2	1:A:676:VAL:N	2.12	0.47
1:A:647[B]:GLU:HG2	1:A:677:SER:HA	1.96	0.47
1:A:121:ASN:HD21	1:A:125:LYS:HZ2	1.63	0.47
1:A:430:HIS:HA	1:A:433[A]:MET:HE2	1.97	0.46
1:A:104:VAL:HG12	1:A:411:CYS:HB2	1.97	0.46
1:A:226:LEU:HD11	1:A:319:ALA:HB2	1.97	0.46
1:A:615:SER:HA	1:A:687:VAL:O	2.15	0.46
1:A:113:LEU:HD13	1:A:469:GLY:HA3	1.98	0.46
1:A:237:ILE:HD12	1:A:263:PHE:CE1	2.51	0.45
1:A:381:LEU:HD22	1:A:526:MET:HE2	1.97	0.45
1:A:237:ILE:HD12	1:A:263:PHE:HE1	1.82	0.45
1:A:363:LEU:HB2	1:A:367:ARG:HB2	1.98	0.45
1:A:510:GLN:HG2	6:A:2594:HOH:O	2.16	0.45
1:A:525:ARG:HD3	1:A:538:MET:O	2.15	0.45
1:A:362:LEU:HD22	6:A:2446:HOH:O	2.16	0.45
1:A:280:GLU:HG3	6:A:2393:HOH:O	2.16	0.44
1:A:134:ASP:OD1	1:A:138:ARG:CD	2.65	0.44
1:A:618:VAL:HG13	6:A:2730:HOH:O	2.18	0.44
1:A:375:ARG:HD2	6:A:2473:HOH:O	2.18	0.43
1:A:136:ASN:N	1:A:137:PRO:CD	2.82	0.43
1:A:367:ARG:HA	1:A:374:HIS:CD2	2.53	0.43
1:A:617:ARG:CD	4:A:812:MGD:H102	2.49	0.42
1:A:119:VAL:O	1:A:123:ILE:HG13	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:430:HIS:HD2	1:A:433[A]:MET:CE	2.32	0.42
1:A:524:TRP:CE2	1:A:535:PHE:HZ	2.37	0.42
1:A:138:ARG:HD3	1:A:396:HIS:HB2	2.00	0.42
1:A:533:TYR:O	1:A:534:ASP:C	2.58	0.42
1:A:503:ARG:NE	6:A:2583:HOH:O	2.52	0.42
1:A:207:ARG:CZ	6:A:2270:HOH:O	2.67	0.41
1:A:345:GLY:HA3	4:A:812:MGD:S12	2.60	0.41
1:A:425:ASN:ND2	1:A:702[B]:ASP:OD2	2.45	0.41
1:A:345:GLY:HA3	4:A:812:MGD:C13	2.51	0.41
1:A:368:ALA:N	1:A:374:HIS:HD2	2.06	0.41
1:A:487:PRO:HB2	1:A:491:CYS:HB2	2.02	0.41
1:A:18:THR:HG21	1:A:183:VAL:HB	2.03	0.41
1:A:132:ASN:HD22	1:A:394:GLY:HA3	1.85	0.41
1:A:168:THR:CG2	6:A:2414:HOH:O	2.60	0.41
1:A:426:LEU:HA	1:A:426:LEU:HD23	1.87	0.41
1:A:515:ARG:HE	1:A:515:ARG:HB2	1.63	0.40
1:A:500:GLU:OE1	1:A:504:ARG:HD2	2.21	0.40
1:A:540[A]:ARG:NH1	1:A:540[A]:ARG:CG	2.85	0.40
1:A:367:ARG:HB3	1:A:374:HIS:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 5 ligands modelled in this entry, 1 is monoatomic and 1 is unknown - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MGD	A	812	3	41,52,52	1.50	3 (7%)	40,81,81	1.74	7 (17%)
2	SF4	A	800	1	0,12,12	-	-	-	-	-
4	MGD	A	811	3	41,52,52	1.44	2 (4%)	40,81,81	1.69	8 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MGD	A	812	3	-	5/18/66/66	0/6/6/6
2	SF4	A	800	1	-	-	0/6/5/5
4	MGD	A	811	3	-	5/18/66/66	0/6/6/6

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	812	MGD	C16-C21	6.57	1.49	1.38
4	A	811	MGD	C16-C21	5.51	1.48	1.38
4	A	811	MGD	O4'-C1'	4.01	1.46	1.41
4	A	812	MGD	C6-N1	-2.45	1.34	1.37
4	A	812	MGD	C17-N18	-2.13	1.34	1.38

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	811	MGD	O17-C17-C16	-4.75	116.36	127.24
4	A	812	MGD	O4'-C4'-C5'	-4.42	94.85	109.37
4	A	812	MGD	O11-C23-N22	-4.03	104.43	108.57
4	A	811	MGD	C19-N20-C21	4.01	120.67	113.43
4	A	812	MGD	C5'-C4'-C3'	-3.92	100.47	115.18
4	A	812	MGD	C19-N20-C21	3.91	120.49	113.43
4	A	811	MGD	C8-N7-C5	3.23	109.14	102.99
4	A	812	MGD	O17-C17-C16	-2.62	121.24	127.24
4	A	811	MGD	O6-C6-N1	-2.56	117.62	120.65
4	A	811	MGD	C16-C17-N18	2.50	119.75	112.31
4	A	811	MGD	C17-C16-N15	2.35	123.07	116.76
4	A	811	MGD	C19-N18-C17	-2.32	120.87	125.10
4	A	812	MGD	C17-C16-N15	2.27	122.85	116.76
4	A	811	MGD	PA-O3B-PB	-2.18	125.33	132.83
4	A	812	MGD	PA-O3B-PB	-2.05	125.80	132.83

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	811	MGD	C5'-O5'-PB-O1B
4	A	811	MGD	C5'-O5'-PB-O3B
4	A	812	MGD	C3'-C4'-C5'-O5'
4	A	812	MGD	O4'-C4'-C5'-O5'
4	A	811	MGD	PA-O3B-PB-O5'
4	A	812	MGD	PA-O3B-PB-O5'
4	A	811	MGD	PA-O3B-PB-O1B
4	A	812	MGD	PB-O3B-PA-O2A
4	A	812	MGD	C5'-O5'-PB-O1B
4	A	811	MGD	O4'-C4'-C5'-O5'


There are no ring outliers.


2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	812	MGD	10	0
4	A	811	MGD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	720/723 (99%)	-0.48	0 100 100	26, 35, 43, 68	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	UNX	A	813	1/1	0.87	0.44	25,25,25,25	0
4	MGD	A	811	47/47	0.97	0.10	21,28,34,37	0
4	MGD	A	812	47/47	0.98	0.11	25,28,30,33	0
2	SF4	A	800	8/8	0.98	0.05	30,32,39,40	0
3	MO	A	810	1/1	0.99	0.13	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.