



Full wwPDB X-ray Structure Validation Report i

Apr 28, 2024 – 01:39 pm BST

PDB ID : 2WTM
Title : Est1E from Butyrivibrio proteoclasticus
Authors : Goldstone, D.C.; Arcus, V.L.
Deposited on : 2009-09-17
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)


There are 4 unique types of molecules in this entry. The entry contains 8709 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EST1E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	250	Total	C 1972	N 1263	O 323	S 373	13	0	4	0
1	B	242	Total	C 1905	N 1219	O 308	S 365	13	0	4	0
1	C	250	Total	C 1979	N 1267	O 323	S 376	13	0	5	0
1	D	243	Total	C 1901	N 1221	O 312	S 355	13	0	4	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O4P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	O 5	P 4	1	0	0
2	B	1	Total	O 5	P 4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	216	Total O 216 216	0	0
4	B	214	Total O 214 214	0	0
4	C	259	Total O 259 259	0	0
4	D	197	Total O 197 197	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: EST1E

Chain A:  92% 8%




- Molecule 1: EST1E

Chain B:  88% 8% • 2%




- Molecule 1: EST1E

Chain C:  92% 8% • %



- Molecule 1: EST1E

Chain D:  85% 11% • • 2%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.03Å 109.07Å 91.65Å 90.00° 102.18° 90.00°	Depositor
Resolution (Å)	89.44 – 1.60 44.79 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.4 (89.44-1.60) 99.4 (44.79-1.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.02 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R , R_{free}	0.156 , 0.194 0.171 , 0.209	Depositor DCC
R_{free} test set	6580 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.6	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8709	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 65.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.5874e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	3/2027 (0.1%)	0.98	5/2748 (0.2%)
1	B	1.02	1/1956 (0.1%)	0.95	4/2655 (0.2%)
1	C	1.05	3/2037 (0.1%)	1.01	6/2763 (0.2%)
1	D	1.00	3/1948 (0.2%)	0.97	5/2647 (0.2%)
All	All	1.02	10/7968 (0.1%)	0.98	20/10813 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	215	CYS	CB-SG	-6.33	1.71	1.82
1	B	215	CYS	CB-SG	-6.20	1.71	1.82
1	C	198	GLU	CB-CG	6.19	1.64	1.52
1	D	203	GLU	CG-CD	6.16	1.61	1.51
1	C	179	GLU	CD-OE1	5.90	1.32	1.25
1	A	198	GLU	CB-CG	5.90	1.63	1.52
1	D	210	LYS	CD-CE	5.32	1.64	1.51
1	A	51	GLU	CB-CG	-5.28	1.42	1.52
1	C	12	ALA	CA-CB	-5.10	1.41	1.52
1	A	240	GLU	C-N	5.04	1.45	1.34

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	139	ARG	NE-CZ-NH2	-9.66	115.47	120.30
1	B	40	ARG	NE-CZ-NH2	-8.38	116.11	120.30
1	B	180	ASP	CB-CG-OD2	-8.19	110.93	118.30
1	C	77	PHE	CB-CG-CD2	-7.85	115.30	120.80
1	D	177	ARG	NE-CZ-NH1	7.40	124.00	120.30
1	B	40	ARG	NE-CZ-NH1	7.31	123.95	120.30
1	D	155	ASP	CB-CG-OD1	6.82	124.44	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	77	PHE	CB-CG-CD1	6.76	125.54	120.80
1	A	195	ASP	CB-CG-OD2	-6.00	112.90	118.30
1	C	163	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	C	180	ASP	CB-CG-OD2	-5.84	113.05	118.30
1	D	139	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	A	117	MET	CG-SD-CE	-5.47	91.44	100.20
1	C	84	LEU	CB-CG-CD2	5.47	120.30	111.00
1	C	228	ASP	CB-CG-OD1	5.39	123.15	118.30
1	A	77	PHE	CB-CG-CD2	-5.17	117.18	120.80
1	A	40	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	D	177	ARG	CG-CD-NE	5.12	122.56	111.80
1	B	73	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	77	PHE	CB-CG-CD1	5.01	124.31	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1972	0	1946	10	0
1	B	1905	0	1879	10	0
1	C	1979	0	1952	10	0
1	D	1901	0	1862	26	0
2	A	5	0	0	0	0
2	B	10	0	0	1	0
2	C	5	0	0	0	0
2	D	10	0	0	1	0
3	A	6	0	8	1	0
3	B	12	0	16	2	0
3	C	6	0	8	0	0
3	D	12	0	16	0	0
4	A	216	0	0	7	0
4	B	214	0	0	2	1
4	C	259	0	0	4	1
4	D	197	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8709	0	7687	57	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:73:ASP:N	4:D:2054:HOH:O	1.78	1.15
1:A:4[B]:ASP:OD1	4:A:2012:HOH:O	1.64	1.15
1:D:34[A]:THR:HG21	1:D:163:ARG:HD3	1.38	1.06
1:D:34[A]:THR:CG2	1:D:163:ARG:HD3	2.00	0.91
1:D:4[B]:ASP:OD1	4:D:2005:HOH:O	1.90	0.86
1:A:119:ARG:O	4:A:2111:HOH:O	1.97	0.81
1:D:166:LYS:HG3	1:D:168:ASN:H	1.51	0.75
1:D:73:ASP:OD1	1:D:166:LYS:HG2	1.85	0.75
1:C:47:GLU:OE1	4:C:2059:HOH:O	2.05	0.74
1:D:151:GLU:OE1	4:D:2117:HOH:O	2.06	0.73
1:D:33[B]:PHE:CD2	1:D:34[B]:THR:HG23	2.25	0.70
1:A:122:ILE:N	4:A:2111:HOH:O	2.24	0.69
1:C:136[B]:GLU:HG2	4:C:2141:HOH:O	1.93	0.69
1:A:203:GLU:OE1	4:A:2183:HOH:O	2.13	0.67
1:C:133:MET:HE3	1:C:136[B]:GLU:HG3	1.81	0.63
1:B:139:ARG:HG2	3:B:1252:GOL:H2	1.80	0.63
1:A:-1:GLY:N	4:A:2003:HOH:O	2.33	0.60
1:D:34[A]:THR:CG2	1:D:163:ARG:CD	2.76	0.60
1:D:166:LYS:HD3	1:D:168:ASN:ND2	2.18	0.57
1:C:240:GLU:O	1:C:244[A]:GLU:HG3	2.06	0.55
1:D:18:LYS:O	1:D:19:ASN:CB	2.56	0.52
1:A:119:ARG:C	4:A:2111:HOH:O	2.46	0.52
1:D:187:LYS:HB3	1:D:188:PRO:CD	2.40	0.51
1:D:143:LEU:HG	1:D:144:LEU:HG	1.92	0.51
1:D:33[B]:PHE:O	1:D:34[B]:THR:HG23	2.11	0.50
1:B:47:GLU:OE1	4:B:2044:HOH:O	2.20	0.50
1:D:166:LYS:HG3	1:D:168:ASN:N	2.22	0.50
1:D:39:GLU:OE2	2:D:1250:PO4:O1	2.31	0.49
1:A:187:LYS:HB3	1:A:188:PRO:CD	2.43	0.48
1:B:236:GLU:O	1:B:240[B]:GLU:HG3	2.14	0.48
1:D:144:LEU:HB3	1:D:159:ALA:HB2	1.95	0.47
1:D:187:LYS:HB3	1:D:188:PRO:HD2	1.96	0.47
1:C:169:TYR:C	1:C:169:TYR:CD2	2.88	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:147:LYS:HE3	1:D:148:PHE:O	2.17	0.45
1:B:109:LEU:HD13	1:B:132:ALA:HA	1.99	0.44
1:D:33[B]:PHE:C	1:D:34[B]:THR:HG23	2.38	0.44
3:A:1250:GOL:O1	4:A:2215:HOH:O	2.15	0.44
1:C:92:LYS:CE	4:C:2120:HOH:O	2.65	0.44
1:A:206:VAL:HG12	1:A:210:LYS:HE2	1.99	0.44
1:B:101:MET:HG3	1:B:122:ILE:HD12	1.99	0.44
1:B:156[B]:GLU:CG	4:B:2132:HOH:O	2.66	0.44
1:C:133:MET:HE1	1:C:136[A]:GLU:OE1	2.19	0.43
1:B:187:LYS:HB3	1:B:188:PRO:CD	2.48	0.43
1:A:184:LYS:HE2	3:B:1252:GOL:H11	1.99	0.43
1:A:80:LEU:HD23	1:A:80:LEU:HA	1.90	0.43
1:D:33[B]:PHE:CE2	1:D:34[B]:THR:CG2	3.02	0.42
1:D:137:ILE:HG22	1:D:142:GLU:HB2	2.01	0.42
1:C:185:TYR:CZ	1:C:187:LYS:HB2	2.54	0.42
1:C:19:ASN:OD1	1:C:21:PRO:HD3	2.19	0.42
1:C:92:LYS:NZ	4:C:2120:HOH:O	2.52	0.42
1:D:137:ILE:CG2	1:D:142:GLU:HB2	2.50	0.41
1:D:166:LYS:CE	4:D:2135:HOH:O	2.68	0.41
1:B:5:CYS:O	1:B:8:ILE:HG13	2.21	0.41
1:B:39:GLU:OE2	2:B:1250:PO4:O1	2.38	0.41
1:B:2:TYR:CD2	1:B:2:TYR:C	2.95	0.41
1:D:40:ARG:NH1	4:D:2028:HOH:O	2.39	0.40
1:D:119:ARG:CZ	4:D:2086:HOH:O	2.69	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:2091:HOH:O	4:C:2212:HOH:O[2_457]	2.13	0.07

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	252/251 (100%)	242 (96%)	10 (4%)	0	100 100
1	B	242/251 (96%)	235 (97%)	7 (3%)	0	100 100
1	C	253/251 (101%)	243 (96%)	10 (4%)	0	100 100
1	D	243/251 (97%)	233 (96%)	9 (4%)	1 (0%)	34 15
All	All	990/1004 (99%)	953 (96%)	36 (4%)	1 (0%)	51 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	19	ASN

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/212 (99%)	209 (100%)	1 (0%)	88 80
1	B	205/212 (97%)	203 (99%)	2 (1%)	76 61
1	C	212/212 (100%)	212 (100%)	0	100 100
1	D	199/212 (94%)	196 (98%)	3 (2%)	65 44
All	All	826/848 (97%)	820 (99%)	6 (1%)	84 73

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	ASP
1	B	146	LEU
1	B	177	ARG
1	D	-2	SER
1	D	8	ILE
1	D	166	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	D	196	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	D	1250	-	4,4,4	0.92	0	6,6,6	0.73	0
2	PO4	C	1248	-	4,4,4	1.06	0	6,6,6	1.44	1 (16%)
2	PO4	B	1249	-	4,4,4	1.23	1 (25%)	6,6,6	0.73	0
2	PO4	D	1249	-	4,4,4	1.03	0	6,6,6	2.16	3 (50%)
3	GOL	A	1250	-	5,5,5	0.42	0	5,5,5	0.42	0
2	PO4	B	1250	-	4,4,4	0.64	0	6,6,6	0.65	0
3	GOL	C	1249	-	5,5,5	0.55	0	5,5,5	0.69	0
3	GOL	D	1252	-	5,5,5	0.22	0	5,5,5	1.29	1 (20%)
2	PO4	A	1249	-	4,4,4	0.79	0	6,6,6	1.60	2 (33%)
3	GOL	D	1251	-	5,5,5	0.54	0	5,5,5	1.09	0
3	GOL	B	1252	-	5,5,5	0.20	0	5,5,5	0.56	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	1251	-	5,5,5	0.74	0	5,5,5	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1250	-	-	2/4/4/4	-
3	GOL	D	1252	-	-	2/4/4/4	-
3	GOL	C	1249	-	-	0/4/4/4	-
3	GOL	D	1251	-	-	2/4/4/4	-
3	GOL	B	1252	-	-	0/4/4/4	-
3	GOL	B	1251	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1249	PO4	P-O2	-2.19	1.48	1.54

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1249	PO4	O4-P-O3	-3.76	95.91	107.97
2	A	1249	PO4	O2-P-O1	-2.65	101.21	110.89
2	A	1249	PO4	O4-P-O2	2.43	115.77	107.97
2	D	1249	PO4	O3-P-O1	-2.33	102.37	110.89
2	D	1249	PO4	O4-P-O1	2.31	119.33	110.89
2	C	1248	PO4	O4-P-O1	-2.22	102.77	110.89
3	D	1252	GOL	O3-C3-C2	-2.08	100.22	110.20

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	1251	GOL	C1-C2-C3-O3
3	D	1252	GOL	C1-C2-C3-O3
3	D	1252	GOL	O2-C2-C3-O3
3	A	1250	GOL	O1-C1-C2-C3
3	D	1251	GOL	O2-C2-C3-O3
3	A	1250	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1250	PO4	1	0
3	A	1250	GOL	1	0
2	B	1250	PO4	1	0
3	B	1252	GOL	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	250/251 (99%)	-0.43	1 (0%)	92	92	8, 16, 27, 41
1	B	242/251 (96%)	-0.29	6 (2%)	57	55	9, 17, 31, 44
1	C	250/251 (99%)	-0.44	2 (0%)	86	86	7, 14, 24, 39
1	D	243/251 (96%)	-0.33	6 (2%)	57	55	8, 16, 29, 39
All	All	985/1004 (98%)	-0.37	15 (1%)	73	73	7, 16, 28, 44

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	-2	SER	3.9
1	B	161	ASP	3.7
1	B	159	ALA	3.6
1	A	248	LYS	3.5
1	D	19	ASN	3.0
1	C	20	ASN	2.9
1	D	-2	SER	2.7
1	B	20	ASN	2.6
1	D	64	HIS	2.5
1	B	160	TRP	2.5
1	D	20	ASN	2.4
1	D	160	TRP	2.4
1	B	162	GLY	2.4
1	B	2	TYR	2.2
1	D	144	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	1252	6/6	0.75	0.33	39,44,44,46	0
3	GOL	D	1252	6/6	0.83	0.14	21,27,32,33	0
3	GOL	A	1250	6/6	0.84	0.15	44,47,47,48	0
3	GOL	B	1251	6/6	0.87	0.14	20,26,30,31	0
3	GOL	D	1251	6/6	0.91	0.10	20,24,29,34	0
3	GOL	C	1249	6/6	0.93	0.14	27,30,31,32	0
2	PO4	A	1249	5/5	0.96	0.10	15,27,29,29	0
2	PO4	C	1248	5/5	0.98	0.06	10,17,20,20	0
2	PO4	D	1250	5/5	0.98	0.05	14,14,15,17	0
2	PO4	B	1249	5/5	0.98	0.06	14,15,17,19	0
2	PO4	B	1250	5/5	0.98	0.05	16,16,18,18	0
2	PO4	D	1249	5/5	0.99	0.05	11,14,20,21	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.