

Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2021 – 02:09 AM EDT

PDB ID : 3ICL

Title: X-Ray Structure of Protein (EAL/GGDEF domain protein) from

M.capsulatus, Northeast Structural Genomics Consortium Target McR174C

Authors: Kuzin, A.; Chen, Y.; Seetharaman, J.; Mao, M.; Xiao, R.; Ciccosanti, C.;

Foote, E.L.; Wang, H.; Everett, J.K.; Nair, R.; Acton, T.B.; Rost, B.; Montelione, G.T.; Tong, L.; Hunt, J.F.; Northeast Structural Genomics Consortium

(NESG)

Deposited on : 2009-07-17

Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.23.2

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	171	70%	25%	5%
1	В	171	68%	20%	• 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Ι	Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
	2	SO4	A	173	_	-	X	-



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2577 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called EAL/GGDEF domain protein.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	Δ	162	Total	С	N	О	S	Se	0	0	0
	11	102	1236	775	224	233	1	3			
1	D	155	Total	С					20	E .	0
1	Б	199	1210	760	219	227	1	3	20)	U

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	initiating methionine	UNP Q60BX6
A	80	LYS	GLU	engineered mutation	UNP Q60BX6
A	164	GLU	-	expression tag	UNP Q60BX6
A	165	HIS	-	expression tag	UNP Q60BX6
A	166	HIS	-	expression tag	UNP Q60BX6
A	167	HIS	-	expression tag	UNP Q60BX6
A	168	HIS	-	expression tag	UNP Q60BX6
A	169	HIS	-	expression tag	UNP Q60BX6
A	170	HIS	-	expression tag	UNP Q60BX6
В	0	MSE	-	initiating methionine	UNP Q60BX6
В	80	LYS	GLU	engineered mutation	UNP Q60BX6
В	164	GLU	-	expression tag	UNP Q60BX6
В	165	HIS	-	expression tag	UNP Q60BX6
В	166	HIS	-	expression tag	UNP Q60BX6
В	167	HIS	-	expression tag	UNP Q60BX6
В	168	HIS	-	expression tag	UNP Q60BX6
В	169	HIS	-	expression tag	UNP Q60BX6
В	170	HIS	-	expression tag	UNP Q60BX6

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).





Mol	Chain	Residues	Ato	ms		ZeroOcc	AltConf	
2	A	1	Total	О	S	0	0	
	A	1	5	4	1	0	U	
2	A	1	Total	О	S	0	0	
	Λ	1	5	4	1	U	U	
2	A	1	Total	Ο	S	0	0	
	Λ	1	5	4	1	U	U	
2	В	1	Total	Ο	S	0	0	
	D	1	5	4	1	U	U	
2	В	1	Total	Ο	S	0	0	
	D	1	5	4	1	0	U	
2	В	1	Total	Ο	S	0	0	
	D	1	5	4	1	O	U	
2	В	1	Total	Ο	S	0	0	
	D	1	5	4	1	0	U	
2	В	1	Total	Ο	S	0	0	
	ע	1	5	4	1			

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	48	Total O 48 48	0	0
3	В	43	Total O 43 43	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EAL/GGDEF domain protein





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	40.67Å 75.95Å 95.20Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.68 - 2.00	Depositor
rtesolution (A)	29.68 - 2.00	EDS
% Data completeness	99.4 (29.68-2.00)	Depositor
(in resolution range)	99.4 (29.68-2.00)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.66 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.4_115, REFMAC	Depositor
D D.	0.208 , 0.222	Depositor
R, R_{free}	0.217 , 0.223	DCC
R_{free} test set	1046 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 49.3	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2577	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.90% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	0.37	0/1251	0.64	0/1689
1	В	0.57	1/1237 (0.1%)	0.62	0/1668
All	All	0.48	$1/2488 \; (0.0\%)$	0.63	0/3357

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	В	136	GLN	C-N	-5.53	1.21	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	7	PRO	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1236	0	1251	26	0
1	В	1210	0	1230	49	0
2	A	15	0	0	2	0
2	В	25	0	0	0	0
3	A	48	0	0	0	0
3	В	43	0	0	3	0
All	All	2577	0	2481	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance} \left(\operatorname{\AA} \right)$	overlap (Å)
1:B:32:LEU:O	1:B:33:LEU:HD23	1.39	1.23
1:A:99:GLN:HE22	1:B:158:PHE:H	1.12	0.93
1:B:34:PHE:HE2	1:B:145[A]:MSE:HE1	1.43	0.82
1:B:30:VAL:HG21	1:B:134:MSE:HE1	1.66	0.78
1:B:35:LEU:HB2	1:B:122:ILE:HG12	1.65	0.78
1:B:34:PHE:CE2	1:B:145[A]:MSE:HE1	2.19	0.77
1:B:32:LEU:O	1:B:33:LEU:CD2	2.30	0.75
1:B:35:LEU:C	1:B:35:LEU:HD23	2.14	0.68
1:A:7:PRO:HB3	1:A:11:LEU:HD23	1.76	0.67
2:A:173:SO4:O2	1:B:153:LYS:HG2	1.95	0.66
1:A:99:GLN:NE2	1:B:158:PHE:H	1.89	0.65
1:B:44:ASN:ND2	3:B:203:HOH:O	2.31	0.63
1:A:0:MSE:HE3	1:A:5:GLY:HA2	1.81	0.62
1:B:119[B]:SER:OG	1:B:153:LYS:HE2	2.00	0.61
1:B:33:LEU:HD22	1:B:124:ILE:HG12	1.82	0.60
1:B:116:ILE:HD13	3:B:195:HOH:O	2.03	0.58
1:B:34:PHE:CE2	1:B:145[A]:MSE:CE	2.87	0.57
1:B:35:LEU:HD12	1:B:122:ILE:HD11	1.87	0.57
1:B:33:LEU:HD21	1:B:124:ILE:HG23	1.87	0.56
1:A:32:LEU:HD21	1:A:138:LEU:HG	1.87	0.56
1:B:15:ARG:HB3	1:B:75:ARG:O	2.07	0.55
1:B:33:LEU:CD2	1:B:124:ILE:HG23	2.37	0.54
1:B:138:LEU:C	1:B:138:LEU:HD23	2.28	0.54
1:A:36:ASP:HB2	1:A:145:MSE:SE	2.58	0.53
1:B:35:LEU:HD12	1:B:122:ILE:CG1	2.39	0.53
1:B:33:LEU:CD2	1:B:124:ILE:HG12	2.38	0.53



 $Continued\ from\ previous\ page...$

Continuea from previo		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap $(Å)$
1:A:47:LEU:HD11	1:A:116:ILE:HD11	1.89	0.53
1:A:39:ASN:O	1:A:43:ILE:HG12	2.09	0.53
1:B:36:ASP:HB3	1:B:145[A]:MSE:SE	2.59	0.52
1:B:33:LEU:HB2	1:B:83:ILE:HB	1.92	0.52
1:B:9:ARG:HG3	1:B:10[B]:GLN:HE21	1.75	0.51
1:B:7:PRO:HD2	1:B:9:ARG:HE	1.76	0.51
1:A:9:ARG:NH2	1:A:76:ILE:HB	2.25	0.51
1:B:75:ARG:HG2	1:B:81:PHE:CE1	2.45	0.51
1:A:61:GLU:O	1:A:65:THR:HG23	2.10	0.51
1:A:0:MSE:CE	1:A:5:GLY:HA2	2.42	0.50
1:B:34:PHE:HD2	1:B:145[A]:MSE:SE	2.44	0.50
1:A:133:THR:OG1	1:A:136:GLN:HG3	2.13	0.49
1:A:7:PRO:HD2	1:A:73:VAL:O	2.12	0.49
1:B:119[B]:SER:HB3	1:B:153:LYS:HB2	1.94	0.49
1:A:88:ALA:HB1	1:A:91:THR:OG1	2.13	0.49
1:A:32:LEU:HD21	1:A:138:LEU:CD2	2.44	0.48
1:B:34:PHE:CD2	1:B:145[A]:MSE:CE	2.97	0.48
1:B:13:CYS:HA	1:B:17:LEU:HB2	1.96	0.47
1:B:126:VAL:O	1:B:130:ASP:HB2	2.14	0.46
1:B:36:ASP:OD1	3:B:219:HOH:O	2.20	0.46
1:A:33:LEU:HB2	1:A:83:ILE:HB	1.97	0.46
1:B:9:ARG:H	1:B:9:ARG:HG2	1.54	0.46
1:B:7:PRO:HG2	1:B:8:ASN:N	2.30	0.46
1:A:122:ILE:O	1:A:156:TYR:HA	2.17	0.45
1:B:107:GLN:NE2	1:B:107:GLN:HA	2.30	0.45
1:B:34:PHE:CE1	1:B:76:ILE:HD11	2.53	0.44
1:A:38:ASP:HB2	1:A:119:SER:OG	2.17	0.44
1:A:59:THR:O	1:A:63:ILE:HG13	2.18	0.44
1:B:35:LEU:HD12	1:B:122:ILE:CD1	2.48	0.44
1:A:33:LEU:HG	1:A:124:ILE:HG12	2.00	0.44
1:A:117:VAL:O	1:A:117:VAL:HG13	2.17	0.44
1:B:35:LEU:C	1:B:35:LEU:CD2	2.86	0.43
1:B:32:LEU:HD23	1:B:33:LEU:N	2.33	0.43
1:B:25:ARG:HH11	1:B:25:ARG:HG3	1.84	0.43
1:B:16:LEU:HB2	1:B:76:ILE:HG22	2.00	0.43
1:B:33:LEU:O	1:B:82:THR:HA	2.19	0.42
1:A:102:LEU:HD13	1:B:151:ARG:HD2	2.02	0.42
1:B:32:LEU:HD22	1:B:138:LEU:HA	2.01	0.42
1:A:153:LYS:HG2	2:A:173:SO4:O3	2.20	0.42
1:B:60:ALA:O	1:B:64:ARG:HG3	2.20	0.41
1:A:110:VAL:HG22	1:A:115:GLN:NE2	2.35	0.41



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)	
1:B:76:ILE:HD12	1:B:76:ILE:C	2.41	0.41	
1:B:7:PRO:HB2	1:B:8:ASN:H	1.56	0.40	
1:A:92:LEU:HD13	1:B:158:PHE:CD2	2.57	0.40	
1:A:148:ALA:HB2	1:A:157:GLN:HG2	2.03	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	160/171 (94%)	157 (98%)	3 (2%)	0	100	100
1	В	158/171 (92%)	148 (94%)	9 (6%)	1 (1%)	25	19
All	All	318/342 (93%)	305 (96%)	12 (4%)	1 (0%)	41	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	26	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/137 (96%)	128 (98%)	3 (2%)	50 53	



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	В	130/137 (95%)	126 (97%)	4 (3%)	40 40		
All	All	261/274~(95%)	254 (97%)	7 (3%)	44 46		

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	PHE
1	A	68	ARG
1	A	151	ARG
1	В	32	LEU
1	В	33	LEU
1	В	36	ASP
1	В	80	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	HIS
1	A	99	GLN
1	A	115	GLN
1	A	140	ASN
1	В	23	HIS
1	В	28	ASN
1	В	44	ASN
1	В	107	GLN
1	В	115	GLN
1	В	140	ASN
1	В	155	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	В	ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	SO4	A	173	-	4,4,4	0.25	0	6,6,6	0.07	0
2	SO4	A	171	-	4,4,4	0.27	0	6,6,6	0.07	0
2	SO4	В	174	-	4,4,4	0.19	0	6,6,6	0.09	0
2	SO4	A	172	-	4,4,4	0.25	0	6,6,6	0.08	0
2	SO4	В	176	-	4,4,4	0.28	0	6,6,6	0.05	0
2	SO4	В	175	-	4,4,4	0.25	0	6,6,6	0.07	0
2	SO4	В	173	-	4,4,4	1.15	0	6,6,6	0.71	0
2	SO4	В	171	-	4,4,4	0.23	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

\mathbf{Mol}	Chain	Res	Type	Clashes	Symm-Clashes
2	A	173	SO4	2	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	159/171 (92%)	0.16	7 (4%) 34 33	14, 23, 46, 69	0
1	В	153/171 (89%)	0.40	18 (11%) 4 4	13, 23, 80, 115	4 (2%)
All	All	312/342 (91%)	0.27	25 (8%) 12 11	13, 23, 55, 115	4 (1%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	В	28	ASN	5.7	
1	В	26	ASP	5.5	
1	В	89	LYS	4.4	
1	В	88	ALA	4.3	
1	A	2	THR	4.2	
1	В	7	PRO	4.2	
1	В	27	GLY	3.8	
1	В	8	ASN	3.7	
1	В	161	PRO	3.6	
1	В	86	ASN	3.1	
1	В	29	PRO	3.1	
1	В	25	ARG	3.1	
1	В	33	LEU	3.0	
1	A	161	PRO	2.6	
1	A	116	ILE	2.6	
1	A	26	ASP	2.5	
1	В	32	LEU	2.4	
1	A	115	GLN	2.4	
1	В	35	LEU	2.4	
1	В	150[A]	SER	2.3	
1	A	25	ARG	2.2	
1	В	31	VAL	2.2	
1	A	98	ALA	2.2	
1	В	87	GLY	2.1	



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	9	ARG	2.0

Non-standard residues in protein, DNA, RNA chains (i) 6.2

There are no non-standard protein/DNA/RNA residues in this entry.

Carbohydrates (i) 6.3

There are no monosaccharides in this entry.

Ligands (i) 6.4

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	SO4	A	173	5/5	0.85	0.48	98,100,102,102	0
2	SO4	A	171	5/5	0.94	0.13	43,47,50,60	0
2	SO4	В	176	5/5	0.94	0.16	55,60,61,63	0
2	SO4	В	173	5/5	0.95	0.11	59,59,60,60	0
2	SO4	A	172	5/5	0.95	0.14	39,41,46,50	0
2	SO4	В	175	5/5	0.96	0.13	37,39,45,46	0
2	SO4	В	171	5/5	0.96	0.39	26,45,46,48	0
2	SO4	В	174	5/5	0.99	0.08	31,32,33,45	0

Other polymers (i) 6.5

There are no such residues in this entry.

