

# Full wwPDB X-ray Structure Validation Report (i)

### Jun 24, 2025 – 04:33 pm BST

PDB ID	:	$5LWQ / pdb_00005lwq$
Title	:	CeuE (H227L variant) a periplasmic protein from Campylobacter jejuni
Authors	:	Wilde, E.J.; Blagova, E.; Hughes, A.; Raines, D.J.; Moroz, O.V.; Turkenburg,
		J.P.; Duhme-Klair, AK.; Wilson, K.S.
Deposited on	:	2016-09-19
Resolution	:	1.52  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	164625	5293 (1.54-1.50)
Clashscore	180529	5759(1.54-1.50)
Ramachandran outliers	177936	5653 (1.54-1.50)
Sidechain outliers	177891	5650(1.54-1.50)
RSRZ outliers	164620	5293(1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	291	85%	14%	••
1	В	291	83%	12%	• ••
1	С	291	83%	13%	••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BR	А	401	-	-	Х	-



#### 5LWQ

# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	280	Total	С	Ν	Ο	$\mathbf{S}$	0	7	0
1	Л	289	2196	1425	352	416	3	0	1	0
1	В	288	Total	С	Ν	Ο	S	0	4	0
1	ГБ	200	2176	1416	343	414	3	0	4	U
1	C	200	Total	С	Ν	0	S	0	4	0
		200	2159	1396	353	407	3	0	4	0

• Molecule 1 is a protein called Enterochelin uptake periplasmic binding protein.

Chain	Residue	Modelled	Actual	Comment	Reference
А	20	GLY	-	expression tag	UNP A0A0W8LI20
А	21	PRO	-	expression tag	UNP A0A0W8LI20
А	22	ALA	-	expression tag	UNP A0A0W8LI20
А	23	MET	-	expression tag	UNP A0A0W8LI20
А	227	LEU	HIS	engineered mutation	UNP A0A0W8LI20
В	20	GLY	-	expression tag	UNP A0A0W8LI20
В	21	PRO	-	expression tag	UNP A0A0W8LI20
В	22	ALA	-	expression tag	UNP A0A0W8LI20
В	23	MET	-	expression tag	UNP A0A0W8LI20
В	227	LEU	HIS	engineered mutation	UNP A0A0W8LI20
С	20	GLY	-	expression tag	UNP A0A0W8LI20
С	21	PRO	-	expression tag	UNP A0A0W8LI20
С	22	ALA	-	expression tag	UNP A0A0W8LI20
С	23	MET	-	expression tag	UNP A0A0W8LI20
С	227	LEU	HIS	engineered mutation	UNP A0A0W8LI20

There are 15 discrepancies between the modelled and reference sequences:

• Molecule 2 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	3	Total Br 3 3	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total Br 1 1	0	0
2	С	1	Total Br 1 1	0	0

• Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	С	1	Total Na 1 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	123	Total O 123 123	0	0
4	В	90	Total O 90 90	0	0
4	С	88	Total         O           88         88	0	0

![](_page_4_Picture_9.jpeg)

# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Enterochelin uptake periplasmic binding protein

![](_page_5_Figure_6.jpeg)

![](_page_5_Picture_7.jpeg)

# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	56.92Å $62.56$ Å $67.79$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$82.21^{\circ}$ $76.99^{\circ}$ $76.18^{\circ}$	Depositor
Bosolution(Å)	65.52 - 1.52	Depositor
Resolution (A)	65.52 - 1.52	EDS
% Data completeness	95.3 (65.52-1.52)	Depositor
(in resolution range)	95.3 (65.52 - 1.52)	EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.49 (at 1.52 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
P. P.	0.185 , $0.221$	Depositor
$n, n_{free}$	0.194 , $0.229$	DCC
$R_{free}$ test set	6332 reflections $(4.89%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	23.2	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31, 25.8	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6838	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

![](_page_6_Picture_8.jpeg)

<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BR, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles	
		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	1.56	16/2246~(0.7%)	1.34	11/3038~(0.4%)
1	В	1.57	19/2220~(0.9%)	1.42	15/3005~(0.5%)
1	С	1.57	18/2194~(0.8%)	1.41	11/2971~(0.4%)
All	All	1.57	53/6660~(0.8%)	1.39	37/9014 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

 $\mathbf{Z}$ Mol Chain Res Type Atoms Observed(Å) -7.81 1 А 246VAL N-CA 1.36N-CA 1 В 29SER 6.64 1.54В 1 146PHE N-CA 6.551.54В ASN CA-CB 1.581 141 6.52 $\overline{\mathbf{C}}$ 208 ILE 1 N-CA -6.471.38 1 В 279ILE C-O 6.33 1.31А 80 ASN CA-C 6.28 1.611 1 В 301 MET SD-CE -6.241.641 В 138 ASP CA-C 6.22 1.60271THR C-O 1 А -6.221.16В 227 1 LEU CA-C 1.606.13

C-0

C-O

N-CA

CA-CB

All (53) bond length outliers are listed below:

 $\overline{\mathbf{C}}$ 

В

С

А

1

1

1

1

109

70

194

229

LYS

ASN

SER

LYS

![](_page_7_Picture_12.jpeg)

1.21

1.17

1.53

1.45

Ideal(Å)

1.46

1.46

1.46

1.53

1.46

1.24

1.52

1.79

1.52

1.23

1.52

1.23

1.24

1.46

1.53

![](_page_7_Picture_13.jpeg)

-6.11

-6.02

5.98

-5.96

5LWQ
------

Conti	Continued from previous page								
Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(A)		
1	С	95	GLY	C-O	-5.93	1.20	1.24		
1	А	274	ALA	N-CA	5.93	1.53	1.46		
1	В	136	GLY	C-O	-5.88	1.17	1.23		
1	С	115	ILE	C-O	-5.88	1.17	1.23		
1	В	109	LYS	CA-CB	5.82	1.56	1.52		
1	А	163	ALA	N-CA	5.80	1.53	1.46		
1	А	84	TYR	N-CA	-5.80	1.38	1.46		
1	А	279	ILE	N-CA	-5.78	1.39	1.46		
1	А	49	ASN	C-O	-5.73	1.21	1.23		
1	В	127	LYS	CA-C	5.67	1.60	1.52		
1	А	135	VAL	C-O	5.65	1.30	1.24		
1	В	215	ILE	CA-C	-5.62	1.46	1.52		
1	В	40	SER	CA-C	-5.50	1.45	1.52		
1	С	258	ARG	CA-C	-5.47	1.46	1.52		
1	С	66	ALA	C-O	5.44	1.30	1.24		
1	С	212	VAL	CA-C	-5.40	1.47	1.52		
1	С	193	ASN	N-CA	-5.39	1.39	1.46		
1	С	105	ILE	N-CA	-5.32	1.40	1.46		
1	С	240	ASN	CA-CB	5.29	1.56	1.52		
1	В	138	ASP	C-O	-5.29	1.17	1.24		
1	В	153	VAL	C-O	5.23	1.30	1.24		
1	А	81	LEU	CB-CG	-5.23	1.43	1.53		
1	С	240	ASN	N-CA	-5.21	1.42	1.46		
1	А	110	PRO	CA-C	-5.19	1.46	1.52		
1	С	152	SER	CA-CB	-5.19	1.45	1.53		
1	С	27	SER	CA-C	5.17	1.58	1.52		
1	А	191	LEU	CA-CB	-5.17	1.46	1.53		
1	А	217	ALA	N-CA	-5.13	1.39	1.46		
1	В	135	VAL	C-O	-5.12	1.18	1.24		
1	С	48	LYS	C-O	5.12	1.29	1.23		
1	А	193	ASN	CA-C	5.09	1.60	1.53		
1	А	305	ILE	C-O	5.07	1.29	1.24		
1	С	67	LEU	CA-C	5.06	1.59	1.52		
1	С	37	VAL	C-O	5.05	1.29	1.24		
1	В	244	ILE	CA-CB	-5.04	1.47	1.54		
1	В	83	LYS	CA-C	-5.03	1.46	1.52		
1	С	283	ASP	CG-OD2	5.03	1.34	1.25		
1	В	75	GLY	N-CA	-5.02	1.40	1.45		

 $\alpha$ 1 0 · · .

All (37) bond angle outliers are listed below:

![](_page_8_Picture_7.jpeg)

5LWQ	
<b>ULIIIIIIIIIIIII</b>	

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	$Ideal(^{o})$
						· · ·	
Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	301	MET	CG-SD-CE	-11.27	76.11	100.90
1	С	46	ILE	CA-C-O	9.51	125.41	119.51
1	В	23	MET	CB-CG-SD	8.11	137.02	112.70
1	А	79	LYS	N-CA-C	7.25	119.86	111.02
1	В	23	MET	CA-CB-CG	7.06	128.22	114.10
1	А	94	VAL	N-CA-C	-6.54	106.42	112.43
1	А	269	ALA	N-CA-C	6.40	119.07	111.71
1	А	301	MET	CG-SD-CE	-6.37	86.89	100.90
1	С	144	SER	N-CA-C	-6.27	104.45	111.28
1	С	42	GLY	N-CA-C	6.26	118.58	110.45
1	А	109	LYS	CA-C-O	6.07	125.15	120.48
1	В	145	SER	N-CA-C	-5.86	104.97	111.36
1	С	28	MET	CG-SD-CE	-5.80	88.15	100.90
1	С	233	SER	N-CA-C	5.74	118.31	111.71
1	В	97	VAL	N-CA-C	-5.73	101.38	109.63
1	А	81	LEU	CA-C-N	-5.67	114.91	120.52
1	А	81	LEU	C-N-CA	-5.67	114.91	120.52
1	В	45	LYS	CA-C-N	-5.63	118.43	123.33
1	В	45	LYS	C-N-CA	-5.63	118.43	123.33
1	В	258	ARG	NE-CZ-NH2	-5.59	114.17	119.20
1	А	28	MET	CG-SD-CE	-5.58	88.61	100.90
1	А	95	GLY	N-CA-C	5.53	117.89	110.43
1	С	26	ILE	CA-C-O	5.48	127.09	120.85
1	А	117	GLY	N-CA-C	5.40	119.42	112.83
1	В	51	SER	N-CA-C	5.38	119.91	113.23
1	В	46	ILE	CA-C-N	-5.37	114.43	119.85
1	В	46	ILE	C-N-CA	-5.37	114.43	119.85
1	В	147	GLU	N-CA-C	-5.32	105.38	111.07
1	В	130	ALA	N-CA-C	5.30	112.79	108.07
1	С	165	GLU	CA-CB-CG	5.21	124.53	114.10
1	В	294	GLY	O-C-N	-5.19	119.51	123.35
1	А	299	LYS	CG-CD-CE	5.11	123.04	111.30
1	С	269	ALA	N-CA-C	5.10	117.58	111.71
1	В	75	GLY	CA-C-O	-5.08	116.48	121.06
1	С	27	SER	O-C-N	5.08	129.48	123.24
1	С	37	VAL	CA-C-O	5.07	125.66	120.39
1	С	118	ARG	NE-CZ-NH2	5.00	123.70	119.20

Continued from previous page...

There are no chirality outliers.

All (1) planarity outliers are listed below:

![](_page_9_Picture_7.jpeg)

Mol	Chain	Res	Type	Group
1	В	80	ASN	Peptide

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2196	0	2216	27	0
1	В	2176	0	2158	36	0
1	С	2159	0	2124	29	0
2	А	3	0	0	2	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
3	С	1	0	0	0	0
4	А	123	0	0	10	1
4	В	90	0	0	13	0
4	C	88	0	0	9	1
All	All	6838	0	6498	92	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:MET:HA	4:B:578:HOH:O	1.13	1.26
1:A:192:THR:O	4:A:501:HOH:O	1.63	1.16
1:B:23:MET:CG	4:B:503:HOH:O	1.89	1.16
1:C:60[B]:ILE:HD12	1:C:114:ILE:HG21	1.31	1.11
1:C:119:GLN:OE1	4:C:501:HOH:O	1.67	1.11
1:C:119:GLN:CD	4:C:501:HOH:O	1.94	1.09
1:A:192:THR:C	4:A:501:HOH:O	1.93	1.09
1:C:258:ARG:NE	4:C:502:HOH:O	1.87	1.08
1:B:23:MET:HG2	4:B:503:HOH:O	1.54	0.98
1:C:119:GLN:NE2	4:C:501:HOH:O	1.98	0.91
1:A:159[B]:LEU:HD22	1:A:159[B]:LEU:N	1.85	0.91
1:C:60[B]:ILE:CD1	1:C:114:ILE:CG2	2.52	0.88
1:C:60[B]:ILE:HD12	1:C:114:ILE:CG2	2.04	0.87

![](_page_10_Picture_11.jpeg)

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:B:60[B]:ILE:HG21	1:B:135:VAL:HG21	1.55	0.87
1:B:227:LEU:O	4:B:502:HOH:O	1.98	0.81
1:A:159[B]:LEU:N	1:A:159[B]:LEU:CD2	2.42	0.81
1:B:23:MET:CB	4:B:503:HOH:O	2.23	0.79
1:A:103:GLU:CG	4:A:623:HOH:O	2.31	0.79
1:B:60[B]:ILE:CD1	1:B:114:ILE:HG21	2.13	0.79
1:A:26:ILE:H	1:A:148:ASN:HD21	1.29	0.79
1:C:26:ILE:H	1:C:148:ASN:HD21	1.32	0.78
1:A:158:GLY:C	1:A:159[B]:LEU:HD22	2.12	0.74
1:A:147:GLU:CG	4:A:567:HOH:O	2.36	0.72
1:B:23:MET:N	4:B:503:HOH:O	2.23	0.72
1:B:55:ILE:HG23	1:B:60[B]:ILE:HD11	1.73	0.70
1:B:23:MET:HA	4:B:584:HOH:O	1.91	0.70
1:B:23:MET:CA	4:B:503:HOH:O	2.40	0.70
1:C:60[B]:ILE:CD1	1:C:114:ILE:HG22	2.21	0.70
1:C:277:LYS:CB	4:C:582:HOH:O	2.39	0.70
1:A:70:ASN:H	1:A:70:ASN:HD22	1.42	0.68
1:B:60[B]:ILE:CD1	1:B:114:ILE:CG2	2.73	0.66
1:C:60[B]:ILE:CD1	1:C:114:ILE:HG21	2.08	0.66
1:B:89:LYS:CD	4:B:546:HOH:O	2.44	0.66
1:C:60[B]:ILE:HG21	1:C:135:VAL:HG21	1.77	0.65
1:B:26:ILE:H	1:B:148:ASN:HD21	1.44	0.64
1:C:258:ARG:CZ	4:C:502:HOH:O	2.36	0.62
1:A:157:TYR:O	1:A:159[B]:LEU:CD2	2.47	0.62
1:C:70:ASN:HD22	1:C:70:ASN:H	1.47	0.62
1:C:99:GLN:CG	4:C:581:HOH:O	2.48	0.62
1:B:194:SER:H	1:B:250:ASN:HD21	1.46	0.61
1:C:57:ASP:OD2	1:C:60[B]:ILE:HG23	2.00	0.60
1:C:258:ARG:NH2	4:C:502:HOH:O	2.36	0.59
1:C:250:ASN:HD22	1:C:255:ASN:HD22	1.51	0.59
1:B:250:ASN:HD22	1:B:255:ASN:HD22	1.51	0.59
1:B:57:ASP:OD2	1:B:60[B]:ILE:HG23	2.03	0.58
1:B:60[B]:ILE:HD13	1:B:114:ILE:CG2	2.34	0.58
1:B:210:HIS:HD2	1:B:215:ILE:O	1.87	0.57
1:B:70:ASN:H	1:B:70:ASN:HD22	1.52	0.57
1:B:60[B]:ILE:HD12	1:B:114:ILE:HG21	1.88	0.55
1:B:60[B]:ILE:HD13	1:B:114:ILE:HG21	1.86	0.55
1:B:23:MET:CA	4:B:578:HOH:O	1.95	0.54
1:B:26:ILE:N	1:B:148:ASN:HD21	2.05	0.54
1:B:194:SER:N	1:B:250:ASN:HD21	2.04	0.54
1:A:234:GLU:CG	4:A:575:HOH:O	2.56	0.53

Continued from previous page...

![](_page_11_Picture_6.jpeg)

Atom-1	Atom-2	Interatomic	Clash
	2100HT 2	distance $(A)$	overlap (Å)
1:C:60[B]:ILE:HD11	1:C:114:ILE:CG2	2.36	0.52
1:B:23:MET:N	4:B:505:HOH:O	2.42	0.51
1:C:210:HIS:HD2	1:C:215:ILE:O	1.94	0.51
1:B:134:PHE:CE2	1:B:136:GLY:HA3	2.46	0.51
1:C:220:GLU:CB	4:C:584:HOH:O	2.58	0.50
1:A:192:THR:CB	4:A:501:HOH:O	2.54	0.49
1:A:192:THR:OG1	4:A:501:HOH:O	1.99	0.48
1:C:194:SER:N	1:C:250:ASN:HD21	2.11	0.47
1:C:60[B]:ILE:HD11	1:C:114:ILE:HG22	1.94	0.47
1:C:142:PHE:C	1:C:142:PHE:CD1	2.92	0.46
1:A:210:HIS:HE1	4:A:530:HOH:O	1.97	0.46
1:C:194:SER:H	1:C:250:ASN:HD21	1.63	0.46
1:A:39:ASP:OD2	2:A:401:BR:BR	2.89	0.46
1:B:23:MET:HB2	1:B:138:ASP:OD2	2.15	0.46
1:B:227:LEU:CB	4:B:504:HOH:O	2.64	0.46
1:B:255:ASN:O	1:B:258:ARG:NH2	2.50	0.45
1:B:55:ILE:O	1:B:77:PRO:HD3	2.17	0.45
1:A:37:VAL:HG21	1:A:46:ILE:HD12	1.98	0.45
1:A:193:ASN:HB3	4:A:521:HOH:O	2.15	0.45
1:C:60[B]:ILE:HD13	1:C:135:VAL:HG21	1.99	0.45
1:C:28:MET:HB3	1:C:28:MET:HE2	1.78	0.44
1:A:180:ILE:C	1:A:180:ILE:HD12	2.42	0.44
1:A:192:THR:CA	4:A:501:HOH:O	2.52	0.44
1:A:26:ILE:N	1:A:148:ASN:HD21	2.06	0.44
1:A:70:ASN:H	1:A:70:ASN:ND2	2.12	0.43
1:B:60[B]:ILE:CG2	1:B:135:VAL:HG21	2.37	0.43
1:A:157:TYR:O	1:A:159[B]:LEU:HD21	2.19	0.42
1:A:248:ASP:OD1	1:A:250:ASN:HB2	2.18	0.42
1:A:37:VAL:HG12	2:A:401:BR:BR	2.75	0.42
1:A:159[B]:LEU:N	1:A:159[B]:LEU:HD23	2.33	0.42
1:B:60[A]:ILE:HG23	1:B:60[A]:ILE:HD12	1.52	0.42
1:A:210:HIS:HD2	1:A:215:ILE:O	2.02	0.42
1:B:80:ASN:OD1	1:B:97:VAL:N	2.46	0.42
1:A:157:TYR:C	1:A:159[B]:LEU:CD2	2.93	0.41
1:B:70:ASN:H	1:B:70:ASN:ND2	2.17	0.41
1:C:57:ASP:HB3	1:C:60[B]:ILE:CD1	2.51	0.40
1:B:210:HIS:HE1	4:B:536:HOH:O	2.04	0.40
1:C:159[A]:LEU:HD12	1:C:159[A]:LEU:HA	1.92	0.40

Continued from previous page..

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

![](_page_12_Picture_6.jpeg)

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:615:HOH:O	4:C:555:HOH:O[1_455]	1.38	0.82

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	294/291~(101%)	284 (97%)	10 (3%)	0	100 100
1	В	290/291~(100%)	282 (97%)	7(2%)	1 (0%)	37 18
1	С	290/291~(100%)	279~(96%)	11 (4%)	0	100 100
All	All	874/873~(100%)	845 (97%)	28 (3%)	1 (0%)	48 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	98	GLN

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	228/252~(90%)	225~(99%)	3~(1%)	65 39		
1	В	218/252~(86%)	209~(96%)	9~(4%)	26 4		
1	С	215/252 (85%)	206 (96%)	9 (4%)	25 4		
All	All	661/756 (87%)	640 (97%)	21 (3%)	41 8		

![](_page_13_Picture_15.jpeg)

Mol	Chain	Res	Type
1	А	70	ASN
1	А	159[A]	LEU
1	А	159[B]	LEU
1	В	23	MET
1	В	24	LEU
1	В	36	LEU
1	В	51	SER
1	В	70	ASN
1	В	159	LEU
1	В	195[A]	ASN
1	В	195[B]	ASN
1	В	258	ARG
1	С	60[A]	ILE
1	С	60[B]	ILE
1	С	70	ASN
1	С	81	LEU
1	С	159[A]	LEU
1	С	159[B]	LEU
1	С	180	ILE
1	С	256	LYS
1	С	271	THR

All (21) residues with a non-rotameric sidechain are listed below:

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	70	ASN
1	А	90	ASN
1	А	148	ASN
1	А	172	ASN
1	А	195	ASN
1	А	203	GLN
1	А	210	HIS
1	А	276	ASN
1	В	70	ASN
1	В	86	GLN
1	В	148	ASN
1	В	210	HIS
1	В	250	ASN
1	В	276	ASN
1	С	49	ASN
1	С	70	ASN

![](_page_14_Picture_8.jpeg)

Mol	Chain	Res	Type
1	С	90	ASN
1	С	98	GLN
1	С	106	ASN
1	С	148	ASN
1	С	210	HIS
1	С	216	ASN
1	С	250	ASN
1	С	276	ASN

Continued from previous page...

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers (i)

There are no such residues in this entry.

![](_page_15_Picture_21.jpeg)

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

![](_page_16_Picture_5.jpeg)

# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	289/291~(99%)	0.20	10 (3%) 47 53	10, 24, 39, 55	10 (3%)
1	В	288/291~(98%)	0.27	12 (4%) 41 46	12, 26, 40, 71	8 (2%)
1	С	288/291~(98%)	0.36	10 (3%) 47 53	11, 27, 41, 59	7 (2%)
All	All	865/873~(99%)	0.28	32 (3%) 45 51	10, 26, 40, 71	25 (2%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	224	VAL	5.7
1	С	224	VAL	4.5
1	В	97	VAL	4.5
1	С	235	PHE	4.4
1	С	222	ILE	4.4
1	В	225	GLY	4.2
1	С	256	LYS	4.1
1	В	80	ASN	4.0
1	С	227	LEU	3.9
1	А	270	LYS	3.8
1	А	222	ILE	3.7
1	А	97	VAL	3.3
1	В	222	ILE	3.2
1	А	224	VAL	2.8
1	С	223	LYS	2.7
1	В	227	LEU	2.7
1	А	254	GLY	2.7
1	С	24	LEU	2.6
1	А	22	ALA	2.6
1	В	23	MET	2.6
1	С	226	THR	2.5
1	С	180	ILE	2.5
1	В	36	LEU	2.5

![](_page_17_Picture_10.jpeg)

Mol	Chain	Res	Type	RSRZ
1	В	218	VAL	2.4
1	В	226	THR	2.4
1	В	184	ASP	2.4
1	В	223	LYS	2.3
1	С	214	GLY	2.2
1	А	184	ASP	2.1
1	А	122	PHE	2.1
1	А	255	ASN	2.1
1	А	253	LEU	2.0

Continued from previous page...

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	BR	В	401	1/1	0.74	0.21	133,133,133,133	0
2	BR	А	401	1/1	0.92	0.36	86,86,86,86	0
3	NA	С	402	1/1	0.97	0.07	23,23,23,23	0
2	BR	С	401	1/1	0.98	0.04	21,21,21,21	1
2	BR	A	403	1/1	0.99	0.02	$25,\!25,\!25,\!25$	1
2	BR	А	402	1/1	0.99	0.03	$25,\!25,\!25,\!25$	1

## 6.5 Other polymers (i)

There are no such residues in this entry.

![](_page_18_Picture_14.jpeg)