



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 01:25 PM EDT

PDB ID : 5WHM
Title : Crystal Structure of IclR Family Transcriptional Regulator from *Brucella abortus*
Authors : Kim, Y.; Wu, R.; Tesar, C.; Endres, M.; Babnigg, G.; Crosson, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2017-07-17
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 94%
1	B	267	 86% 10%
1	C	267	 91% 7%
1	D	267	 87% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACY	D	301[B]	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called IclR family transcriptional regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	263	2076	1305	373	385	4	9	0	3	0
1	B	260	2042	1286	366	377	4	9	0	1	0
1	C	264	2101	1320	381	385	4	11	0	5	0
1	D	260	2144	1354	381	394	4	11	0	15	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	SER	-	expression tag	UNP A0A1S1ZDZ5
A	19	ASN	-	expression tag	UNP A0A1S1ZDZ5
A	20	ALA	-	expression tag	UNP A0A1S1ZDZ5
B	18	SER	-	expression tag	UNP A0A1S1ZDZ5
B	19	ASN	-	expression tag	UNP A0A1S1ZDZ5
B	20	ALA	-	expression tag	UNP A0A1S1ZDZ5
C	18	SER	-	expression tag	UNP A0A1S1ZDZ5
C	19	ASN	-	expression tag	UNP A0A1S1ZDZ5
C	20	ALA	-	expression tag	UNP A0A1S1ZDZ5
D	18	SER	-	expression tag	UNP A0A1S1ZDZ5
D	19	ASN	-	expression tag	UNP A0A1S1ZDZ5
D	20	ALA	-	expression tag	UNP A0A1S1ZDZ5

- Molecule 2 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 8 4 4	0	1
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	123	Total O 123 123	0	0
5	B	50	Total O 50 50	0	0
5	C	91	Total O 91 91	0	0
5	D	51	Total O 51 51	0	0



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IclR family transcriptional regulator





- Molecule 1: IclR family transcriptional regulator



- Molecule 1: IclR family transcriptional regulator



- Molecule 1: IclR family transcriptional regulator



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.15Å 112.46Å 83.65Å 90.00° 115.86° 90.00°	Depositor
Resolution (Å)	33.56 – 1.95 33.56 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.1 (33.56-1.95) 99.1 (33.56-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 1.95Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.177 , 0.212 0.178 , 0.212	Depositor DCC
R_{free} test set	4467 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtrriage
Anisotropy	0.518	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8715	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EDO, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/2105	0.67	0/2837
1	B	0.44	0/2070	0.63	0/2789
1	C	0.50	0/2130	0.62	0/2868
1	D	0.44	0/2171	0.61	0/2922
All	All	0.49	0/8476	0.63	0/11416

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2076	0	2091	11	0
1	B	2042	0	2064	23	0
1	C	2101	0	2123	13	0
1	D	2144	0	2180	25	0
2	A	8	0	6	0	0
2	B	4	0	3	0	0
2	C	8	0	6	1	0
2	D	12	0	9	3	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	4	0	6	1	0
5	A	123	0	0	0	0
5	B	50	0	0	1	0
5	C	91	0	0	0	0
5	D	51	0	0	4	0
All	All	8715	0	8488	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:79:LEU:HD21	2:C:303:ACY:H2	1.74	0.68
2:D:301[B]:ACY:OXT	5:D:401:HOH:O	2.10	0.68
1:C:242:HIS:CD2	1:C:245:TYR:HD2	2.18	0.61
1:B:104:ILE:HG22	1:D:108[A]:GLY:HA2	1.83	0.60
1:B:105:LEU:HD21	1:D:112[A]:LEU:HG	1.84	0.59
1:A:119:THR:CG2	1:D:277:HIS:HE1	2.18	0.56
1:B:115:LEU:HB2	1:B:118:ARG:HD2	1.90	0.54
1:D:214:LEU:O	1:D:218:GLU:HG3	2.08	0.53
1:D:115[B]:LEU:O	1:D:118:ARG:HB3	2.10	0.52
1:B:151:VAL:HG21	1:B:163:GLN:HB2	1.91	0.52
1:C:102:ILE:HB	4:C:301:EDO:H12	1.93	0.50
2:D:301[A]:ACY:O	5:D:402:HOH:O	2.19	0.49
1:B:111:MSE:SE	1:D:105[A]:LEU:HD21	2.63	0.48
1:B:210:LEU:O	1:B:214:LEU:HG	2.12	0.48
1:B:23:THR:OG1	2:D:301[B]:ACY:H1	2.13	0.48
1:D:161:ARG:NH2	5:D:408:HOH:O	2.48	0.47
1:B:116:MSE:HA	1:B:116:MSE:HE2	1.97	0.47
1:B:63:GLN:NE2	1:B:73:LYS:HD2	2.29	0.46
1:D:161:ARG:HG2	1:D:162:ARG:O	2.15	0.46
1:B:105:LEU:HD23	1:B:105:LEU:HA	1.70	0.46
1:B:214:LEU:O	1:B:218:GLU:HG2	2.15	0.46
1:C:232:ASP:OD2	1:D:116[B]:MSE:HE1	2.15	0.46
1:B:104:ILE:HG23	1:D:107[A]:LEU:HB2	1.98	0.45
1:B:105:LEU:HD11	1:D:112[A]:LEU:CD2	2.47	0.45
1:A:90:ARG:NH1	1:A:99:GLN:OE1	2.48	0.45
1:D:116[B]:MSE:HB3	1:D:116[B]:MSE:HE3	1.64	0.45
1:A:119:THR:HG21	1:D:277:HIS:HE1	1.81	0.45
1:D:198:ARG:HG2	1:D:199:GLU:N	2.32	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:24:MSE:HE1	1:D:32:VAL:HG11	1.99	0.44
1:A:119:THR:HG23	1:D:277:HIS:HE1	1.82	0.44
1:C:183:MSE:SE	1:C:191:ILE:HD12	2.68	0.43
1:A:189:THR:O	1:A:193:GLU:HG2	2.19	0.43
1:D:28:LEU:HG	1:D:48:GLU:HG3	2.00	0.43
1:D:147:VAL:HG22	1:D:167:THR:HG22	2.00	0.43
1:C:279:ILE:O	1:C:283:VAL:HG12	2.18	0.43
1:B:141:ARG:O	1:D:102[A]:ILE:HD11	2.19	0.42
1:D:116[B]:MSE:H	1:D:116[B]:MSE:HG2	1.52	0.42
1:B:185:GLU:O	1:B:189:THR:OG1	2.38	0.42
1:B:111:MSE:SE	1:D:105[A]:LEU:CD2	3.18	0.42
1:A:21:LYS:HA	1:A:21:LYS:HD2	1.86	0.42
1:B:172:GLY:N	5:B:403:HOH:O	2.51	0.42
1:C:267:ASP:OD1	1:C:271:ARG:HD2	2.20	0.42
1:B:167:THR:HG21	1:D:106[A]:ARG:HD3	2.02	0.42
1:B:57:GLU:OE1	1:B:68:LYS:NZ	2.35	0.41
1:C:94:LEU:HD13	1:C:94:LEU:HA	1.90	0.41
1:D:242:HIS:HE2	1:D:283:VAL:CG1	2.33	0.41
1:A:35:GLU:OE1	1:A:37:ARG:NH2	2.48	0.41
1:A:113[B]:SER:OG	1:C:55:PRO:HA	2.21	0.41
1:A:102:ILE:H	1:A:102:ILE:HG13	1.69	0.41
1:A:119:THR:HG21	1:D:277:HIS:CE1	2.55	0.41
1:B:51:ARG:HD3	5:D:444:HOH:O	2.20	0.41
1:B:242:HIS:CG	1:B:245:TYR:HD2	2.39	0.41
1:C:283:VAL:O	1:C:283:VAL:HG13	2.21	0.41
1:B:105:LEU:HD21	1:D:112[A]:LEU:CG	2.48	0.41
1:B:141:ARG:CZ	1:B:191:ILE:HD11	2.51	0.41
1:C:115:LEU:HD22	1:C:245:TYR:HE1	1.85	0.41
1:A:102:ILE:HD11	1:C:141:ARG:O	2.22	0.40
1:C:146:MSE:O	1:C:167:THR:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/267 (98%)	256 (98%)	6 (2%)	0	100	100
1	B	257/267 (96%)	253 (98%)	4 (2%)	0	100	100
1	C	267/267 (100%)	261 (98%)	6 (2%)	0	100	100
1	D	271/267 (102%)	262 (97%)	9 (3%)	0	100	100
All	All	1057/1068 (99%)	1032 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/215 (105%)	221 (98%)	4 (2%)	54	49
1	B	221/215 (103%)	213 (96%)	8 (4%)	30	20
1	C	227/215 (106%)	222 (98%)	5 (2%)	47	41
1	D	232/215 (108%)	221 (95%)	11 (5%)	22	11
All	All	905/860 (105%)	877 (97%)	28 (3%)	38	25

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	LYS
1	A	111	MSE
1	A	157	ASN
1	A	194	HIS
1	B	21	LYS
1	B	37	ARG
1	B	111	MSE
1	B	161	ARG
1	B	163	GLN
1	B	189	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	198	ARG
1	B	208	LYS
1	C	21	LYS
1	C	153	GLN
1	C	189	THR
1	C	202	ASN
1	C	242	HIS
1	D	94	LEU
1	D	105[A]	LEU
1	D	105[B]	LEU
1	D	106[A]	ARG
1	D	106[B]	ARG
1	D	111[A]	MSE
1	D	111[B]	MSE
1	D	175	VAL
1	D	188	ARG
1	D	198	ARG
1	D	257	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	153	GLN
1	B	63	GLN
1	B	153	GLN
1	B	163	GLN
1	C	58	ASN
1	D	62	ASN
1	D	277	HIS
1	D	278	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACY	B	301	-	3,3,3	0.94	0	3,3,3	0.55	0
2	ACY	A	302	-	3,3,3	0.95	0	3,3,3	0.35	0
2	ACY	C	303	-	3,3,3	0.71	0	3,3,3	2.33	2 (66%)
2	ACY	A	301	-	3,3,3	0.95	0	3,3,3	1.02	0
2	ACY	D	301[A]	-	3,3,3	0.80	0	3,3,3	0.87	0
2	ACY	D	301[B]	-	3,3,3	0.84	0	3,3,3	0.82	0
2	ACY	D	302	-	3,3,3	0.75	0	3,3,3	1.01	0
2	ACY	C	302	-	3,3,3	0.78	0	3,3,3	0.83	0
4	EDO	C	301	-	3,3,3	0.43	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	301	-	-	0/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	303	ACY	O-C-CH3	-2.92	110.54	122.53
2	C	303	ACY	OXT-C-O	2.74	132.18	122.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	303	ACY	1	0
2	D	301[A]	ACY	1	0
2	D	301[B]	ACY	2	0
4	C	301	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/267 (95%)	-0.09	4 (1%) 70 76	13, 35, 77, 121	3 (1%)
1	B	251/267 (94%)	0.38	8 (3%) 50 57	24, 49, 90, 117	1 (0%)
1	C	254/267 (95%)	0.06	10 (3%) 44 50	13, 42, 100, 133	4 (1%)
1	D	251/267 (94%)	0.32	11 (4%) 39 46	13, 43, 97, 134	13 (5%)
All	All	1010/1068 (94%)	0.16	33 (3%) 49 56	13, 42, 93, 134	21 (2%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	114[A]	ASN	8.2
1	D	110[A]	ALA	4.9
1	D	22	PRO	4.7
1	B	283	VAL	4.5
1	B	257	PHE	4.0
1	D	109[A]	TYR	3.8
1	D	113[A]	SER	3.8
1	A	32	VAL	3.8
1	B	103	GLY	3.5
1	D	257	PHE	3.4
1	D	23	THR	3.3
1	B	22	PRO	3.2
1	C	30	ALA	3.1
1	C	159	THR	3.1
1	C	156	GLY	2.9
1	A	156	GLY	2.9
1	D	283	VAL	2.8
1	C	284	PRO	2.7
1	B	159	THR	2.6
1	D	200	PRO	2.5
1	C	154	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	204	PRO	2.4
1	C	277[A]	HIS	2.3
1	B	94	LEU	2.3
1	B	205	SER	2.3
1	B	187	GLU	2.2
1	C	32	VAL	2.2
1	C	202	ASN	2.1
1	C	94	LEU	2.1
1	C	194	HIS	2.1
1	D	33	GLU	2.1
1	A	277[A]	HIS	2.0
1	A	283	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACY	D	301[A]	4/4	0.59	0.17	45,45,47,48	4
2	ACY	D	301[B]	4/4	0.59	0.17	46,47,47,48	4
2	ACY	A	302	4/4	0.80	0.16	45,55,57,60	0
2	ACY	C	303	4/4	0.87	0.23	49,57,62,62	0
4	EDO	C	301	4/4	0.87	0.18	57,61,65,67	0
3	CA	A	303	1/1	0.90	0.17	104,104,104,104	0
2	ACY	B	301	4/4	0.91	0.09	36,41,42,43	0
2	ACY	D	302	4/4	0.94	0.10	53,55,55,56	0
2	ACY	C	302	4/4	0.97	0.06	35,36,38,40	0
2	ACY	A	301	4/4	0.99	0.04	21,23,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.