

Full wwPDB X-ray Structure Validation Report (i)

Jun 22, 2024 – 11:18 AM EDT

:	6QKI
:	Native structure of EgtB from Chloracidobacterium thermophilum, a type II
	sulfoxide synthase
:	Stampfli, A.R.; Badri, B.N.; Schirmer, T.; Seebeck, F.P.
:	2019-01-29
:	2.04 Å(reported)
	::

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Matria	Whole archive	Similar resolution		
Metric	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$		
R_{free}	130704	1692 (2.04-2.04)		
Clashscore	141614	1773 (2.04-2.04)		
Ramachandran outliers	138981	1752 (2.04-2.04)		
Sidechain outliers	138945	1752 (2.04-2.04)		
RSRZ outliers	127900	1672(2.04-2.04)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	437	80%	9%	•	10%
1	В	437	4% 81%	8%		10%
1	С	437	3% 81%	7%	•	10%
1	D	437	80%	8%	•	10%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 12935 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A 905		Total	С	Ν	0	\mathbf{S}	7	0	0
	A 395	- 390	3193	2049	573	561	10	1	0	U
1	р	202	Total	С	Ν	0	S	7	0	0
1	D	392	3170	2037	569	554	10	1	0	U
1	C	202	Total	С	Ν	0	S	0	0	0
1		595	3180	2042	571	557	10	0	0	0
1	П	302	Total	С	Ν	0	S	6	0	0
		592	3167	2034	569	554	10	0	0	0

• Molecule 1 is a protein called Uncharacterized protein.

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-2	GLY	-	expression tag	UNP G2LET6
А	-1	SER	-	expression tag	UNP G2LET6
А	0	HIS	-	expression tag	UNP G2LET6
В	-2	GLY	-	expression tag	UNP G2LET6
В	-1	SER	-	expression tag	UNP G2LET6
В	0	HIS	-	expression tag	UNP G2LET6
С	-2	GLY	-	expression tag	UNP G2LET6
С	-1	SER	-	expression tag	UNP G2LET6
С	0	HIS	-	expression tag	UNP G2LET6
D	-2	GLY	-	expression tag	UNP G2LET6
D	-1	SER	-	expression tag	UNP G2LET6
D	0	HIS	-	expression tag	UNP G2LET6

• Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Fe 1 1	0	0
2	В	1	Total Fe 1 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	101	Total O 101 101	0	0
3	В	65	Total O 65 65	0	0
3	С	28	TotalO2828	0	0
3	D	27	Total O 27 27	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Uncharacterized protein







• Molecule 1: Uncharacterized protein





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	84.97Å 127.46Å 89.54Å	Depositor
a, b, c, α , β , γ	90.00° 113.00° 90.00°	Depositor
Bosolution(A)	30.00 - 2.04	Depositor
Resolution (A)	39.21 - 2.04	EDS
% Data completeness	98.5 (30.00-2.04)	Depositor
(in resolution range)	98.5 (39.21-2.04)	EDS
R_{merge}	0.11	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.94 (at 2.05 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
P. P.	0.209 , 0.247	Depositor
n, n_{free}	0.217 , 0.213	DCC
R_{free} test set	5280 reflections $(4.84%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	28.5	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , 30.1	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12935	wwPDB-VP
Average B, all atoms $(Å^2)$	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	ond lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.98	7/3306~(0.2%)	1.07	7/4514~(0.2%)	
1	В	1.07	4/3283~(0.1%)	1.09	7/4484~(0.2%)	
1	С	0.93	2/3293~(0.1%)	1.06	10/4498~(0.2%)	
1	D	0.96	7/3280~(0.2%)	1.08	7/4480~(0.2%)	
All	All	0.98	20/13162~(0.2%)	1.08	31/17976~(0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	1	1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	213	ARG	CA-CB	-24.75	0.99	1.53
1	А	213	ARG	CA-CB	-8.07	1.36	1.53
1	В	242	GLU	CD-OE2	-6.92	1.18	1.25
1	В	324	GLU	CD-OE2	-6.78	1.18	1.25
1	А	272	GLU	CD-OE1	6.49	1.32	1.25
1	D	148	GLU	CD-OE2	-6.33	1.18	1.25
1	А	202	GLU	CD-OE2	6.19	1.32	1.25
1	В	300	GLU	CD-OE2	6.07	1.32	1.25
1	D	41	GLU	CD-OE2	-5.93	1.19	1.25
1	А	431	GLU	CD-OE2	-5.90	1.19	1.25
1	А	242	GLU	CD-OE2	-5.72	1.19	1.25
1	D	409	ARG	C-O	5.64	1.34	1.23
1	D	209	GLY	C-O	5.62	1.32	1.23
1	D	43	GLU	CD-OE2	-5.48	1.19	1.25



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	324	GLU	CD-OE2	-5.48	1.19	1.25
1	С	386	ASN	C-O	5.41	1.33	1.23
1	А	292	GLY	C-O	5.27	1.32	1.23
1	D	328	GLU	CD-OE1	5.13	1.31	1.25
1	С	421	ARG	C-O	5.10	1.33	1.23
1	А	159	GLU	CD-OE1	5.01	1.31	1.25

Continued from previous page...

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	213	ARG	CB-CA-C	12.79	135.98	110.40
1	А	179	ARG	NE-CZ-NH1	9.79	125.19	120.30
1	В	421	ARG	CG-CD-NE	-8.69	93.55	111.80
1	С	421	ARG	CG-CD-NE	-7.93	95.15	111.80
1	С	213	ARG	CG-CD-NE	7.34	127.22	111.80
1	А	421	ARG	CG-CD-NE	-7.14	96.81	111.80
1	С	213	ARG	NE-CZ-NH2	6.99	123.80	120.30
1	D	421	ARG	CG-CD-NE	-6.98	97.14	111.80
1	А	179	ARG	NE-CZ-NH2	-6.93	116.84	120.30
1	D	229	ARG	NE-CZ-NH1	6.92	123.76	120.30
1	С	106	ARG	NE-CZ-NH2	-6.89	116.85	120.30
1	D	278	ARG	NE-CZ-NH2	-6.69	116.95	120.30
1	В	284	THR	CA-CB-OG1	-6.55	95.25	109.00
1	D	205	ARG	NE-CZ-NH1	6.13	123.36	120.30
1	С	386	ASN	C-N-CA	6.10	135.12	122.30
1	D	229	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	С	205	ARG	NE-CZ-NH2	-6.02	117.29	120.30
1	А	421	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	С	284	THR	CA-CB-OG1	-5.93	96.55	109.00
1	В	320	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	D	333	ARG	CB-CG-CD	5.71	126.46	111.60
1	В	213	ARG	CA-CB-CG	5.63	125.79	113.40
1	С	179	ARG	NE-CZ-NH1	5.53	123.07	120.30
1	А	284	THR	CA-CB-CG2	5.47	120.06	112.40
1	В	315	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	А	205	ARG	NE-CZ-NH2	-5.35	117.62	120.30
1	С	213	ARG	CD-NE-CZ	5.31	131.03	123.60
1	С	303	CYS	CA-CB-SG	-5.09	104.83	114.00
1	А	208	ILE	CA-CB-CG2	5.05	120.99	110.90
1	В	284	THR	CA-CB-CG2	5.03	119.45	112.40
1	D	208	ILE	CA-CB-CG2	5.03	120.95	110.90



All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	В	213	ARG	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	212	GLY	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3193	0	3032	17	0
1	В	3170	0	3011	16	0
1	С	3180	0	3022	14	0
1	D	3167	0	3015	11	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	А	101	0	0	2	0
3	В	65	0	0	2	0
3	С	28	0	0	0	0
3	D	27	0	0	0	0
All	All	12935	0	12080	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:ARG:HH11	1:B:179:ARG:HG2	1.30	0.97
1:B:294:GLN:HG3	3:B:665:HOH:O	1.75	0.87
1:C:218:ASP:HB2	1:C:386:ASN:ND2	1.98	0.78
1:A:218:ASP:HB2	1:A:386:ASN:ND2	1.99	0.78



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:136:ASP:OD1	1:C:139:THR:HG23	1.85	0.76
1:B:179:ARG:NH1	3:B:601:HOH:O	2.09	0.74
1:D:136:ASP:OD1	1:D:139:THR:HG23	1.93	0.68
1:B:179:ARG:HH11	1:B:179:ABG:CG	2.06	0.67
1:B:180:PRO:O	1:B:182:PRO:HD3	1.95	0.65
1:B:23:GLN:OE1	1:B:129:GLU:OE1	2.15	0.64
1:C:170:ALA:HA	1:C:288:LEU:HD22	1.80	0.63
1:B:179:ARG:HG2	1:B:179:ARG:NH1	2.05	0.62
1:A:200:ILE:O	1:A:333:ARG:NH2	2.36	0.58
1:C:181:GLN:NE2	1:C:345:VAL:HG13	2.18	0.58
1:A:218:ASP:HB2	1:A:386:ASN:HD22	1.68	0.57
1:B:23:GLN:HG2	1:B:132:ILE:HG21	1.86	0.57
1:A:148:GLU:OE2	3:A:601:HOH:O	2.17	0.56
1:C:218:ASP:HB2	1:C:386:ASN:HD21	1.69	0.54
1:D:208:ILE:HG22	1:D:224:HIS:CE1	2.42	0.54
1:B:229:ARG:CZ	1:B:229:ARG:HB2	2.37	0.54
1:B:92:SER:O	1:B:100:ARG:NH1	2.39	0.54
1:A:156:GLN:NE2	1:A:385:TYR:OH	2.41	0.53
1:B:156:GLN:NE2	1:B:385:TYR:OH	2.42	0.53
1:C:208:ILE:HG22	1:C:224:HIS:CE1	2.44	0.53
1:A:181:GLN:NE2	1:A:345:VAL:HG13	2.24	0.52
1:A:208:ILE:HG22	1:A:224:HIS:CE1	2.45	0.52
1:B:208:ILE:HG22	1:B:224:HIS:CE1	2.44	0.52
1:A:282:ASP:OD1	1:A:284:THR:HB	2.09	0.51
1:C:156:GLN:NE2	1:C:385:TYR:OH	2.44	0.51
1:D:282:ASP:OD1	1:D:284:THR:HB	2.11	0.51
1:D:65:TRP:CZ2	1:D:91:ASN:HB2	2.46	0.51
1:A:177:VAL:HG11	3:A:694:HOH:O	2.10	0.50
1:A:65:TRP:CZ2	1:A:91:ASN:HB2	2.47	0.49
1:B:282:ASP:OD1	1:B:284:THR:HB	2.13	0.49
1:A:23:GLN:HG3	1:A:132:ILE:HG21	1.95	0.49
1:D:181:GLN:NE2	1:D:345:VAL:O	2.47	0.48
1:C:282:ASP:OD1	1:C:284:THR:HB	2.12	0.48
1:D:208:ILE:HD11	1:D:329:VAL:HG23	1.95	0.48
1:C:23:GLN:HG3	1:C:132:ILE:HG21	1.97	0.47
1:A:111:ARG:HA	1:A:112:PRO:C	2.36	0.45
1:B:238:VAL:HB	1:B:427:VAL:HG13	1.97	0.45
1:B:65:TRP:CZ2	1:B:91:ASN:HB2	2.52	0.45
1:D:179:ARG:O	1:D:179:ARG:HG3	2.15	0.45
1:A:387:GLY:HA2	1:A:390:MET:SD	2.56	0.45
1:C:210:HIS:CE1	1:C:213:ARG:HB3	2.52	0.45



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:C:111:ARG:HA	1:C:112:PRO:C	2.37	0.44
1:A:229:ARG:CZ	1:A:229:ARG:HB2	2.48	0.43
1:A:384:GLU:HG2	1:A:385:TYR:N	2.33	0.43
1:D:111:ARG:HA	1:D:112:PRO:C	2.40	0.43
1:C:73:LYS:HD2	1:C:73:LYS:HA	1.81	0.42
1:D:347:HIS:CD2	1:D:348:PRO:HD2	2.55	0.42
1:D:170:ALA:HA	1:D:288:LEU:HD22	2.01	0.41
1:B:111:ARG:HA	1:B:112:PRO:C	2.40	0.41
1:A:73:LYS:HD3	1:A:73:LYS:HA	1.83	0.41
1:C:65:TRP:CZ2	1:C:91:ASN:HB2	2.56	0.41
1:A:347:HIS:CD2	1:A:348:PRO:HD2	2.55	0.41
1:D:53:VAL:HG13	1:D:164:ASP:HB3	2.04	0.40
1:C:275:LEU:HB3	1:C:304:HIS:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	387/437~(89%)	374 (97%)	11 (3%)	2(0%)	29	18
1	В	384/437~(88%)	370~(96%)	11 (3%)	3(1%)	19	10
1	С	385/437~(88%)	372 (97%)	9(2%)	4 (1%)	15	6
1	D	384/437~(88%)	368 (96%)	14 (4%)	2(0%)	29	18
All	All	1540/1748 (88%)	1484 (96%)	45 (3%)	11 (1%)	22	12

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	388	LYS
1	А	77	ALA



J F F F F F F F F F F F F F F F F F F F							
\mathbf{Mol}	Chain	\mathbf{Res}	Type				
1	А	388	LYS				
1	В	77	ALA				
1	С	77	ALA				
1	С	213	ARG				
1	D	77	ALA				
1	В	388	LYS				
1	D	388	LYS				
1	С	18	ASP				
1	В	386	ASN				

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	324/358~(90%)	314 (97%)	10 (3%)	40 33
1	В	321/358~(90%)	314 (98%)	7 (2%)	52 46
1	С	323/358~(90%)	310 (96%)	13 (4%)	31 24
1	D	322/358~(90%)	305~(95%)	17 (5%)	22 14
All	All	1290/1432~(90%)	1243 (96%)	47 (4%)	35 28

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	73	LYS
1	А	179	ARG
1	А	195	THR
1	А	208	ILE
1	А	213	ARG
1	А	279	GLN
1	А	288	LEU
1	А	354	SER
1	А	384	GLU
1	А	422	TRP
1	В	24	ARG
1	В	73	LYS



Mol	Chain	Res	Type
1	В	92	SER
1	В	179	ARG
1	В	208	ILE
1	В	288	LEU
1	В	422	TRP
1	С	20	LYS
1	С	73	LYS
1	С	139	THR
1	С	179	ARG
1	С	208	ILE
1	С	213	ARG
1	С	279	GLN
1	С	288	LEU
1	С	294	GLN
1	С	353	VAL
1	С	376	ARG
1	С	388	LYS
1	С	422	TRP
1	D	20	LYS
1	D	56	PRO
1	D	73	LYS
1	D	78	ASP
1	D	139	THR
1	D	179	ARG
1	D	181	GLN
1	D	195	THR
1	D	208	ILE
1	D	213	ARG
1	D	229	ARG
1	D	279	GLN
1	D	288	LEU
1	D	294	GLN
1	D	388	LYS
1	D	391	CYS
1	D	422	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	156	GLN
1	А	181	GLN
1	А	279	GLN



Mol	Chain	Res	Type
1	А	386	ASN
1	А	392	ASN
1	В	156	GLN
1	В	279	GLN
1	С	156	GLN
1	С	181	GLN
1	С	210	HIS
1	С	386	ASN
1	D	181	GLN
1	D	279	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	395/437~(90%)	-0.16	15 (3%) 40 44	21, 32, 65, 91	1 (0%)
1	В	392/437~(89%)	-0.12	16 (4%) 37 40	20, 31, 70, 98	1 (0%)
1	С	393/437~(89%)	-0.17	14 (3%) 42 46	23, 36, 70, 95	0
1	D	392/437~(89%)	-0.07	8 (2%) 65 69	21, 33, 67, 103	1 (0%)
All	All	1572/1748~(89%)	-0.13	53 (3%) 45 49	20, 33, 69, 103	3 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	353	VAL	6.5
1	D	194	PRO	5.4
1	С	386	ASN	4.7
1	В	99	ALA	4.6
1	D	99	ALA	4.4
1	D	434	SER	4.3
1	А	80	ARG	4.1
1	А	434	SER	4.1
1	D	78	ASP	4.1
1	С	281	ALA	4.0
1	А	78	ASP	3.6
1	В	212	GLY	3.5
1	В	80	ARG	3.5
1	С	17	SER	3.4
1	А	77	ALA	3.3
1	С	213	ARG	3.3
1	С	195	THR	3.2
1	С	100	ARG	3.1
1	С	104	PRO	3.1
1	В	20	LYS	3.0
1	С	102	PRO	3.0



Mol	Chain	Res	Type	RSRZ
1	С	434	SER	3.0
1	В	73	LYS	2.9
1	А	212	GLY	2.9
1	В	434	SER	2.9
1	А	17	SER	2.8
1	А	137	THR	2.8
1	С	376	ARG	2.8
1	D	281	ALA	2.7
1	А	281	ALA	2.5
1	В	138	ARG	2.5
1	А	386	ASN	2.5
1	D	195	THR	2.4
1	В	17	SER	2.4
1	В	376	ARG	2.4
1	В	354	SER	2.4
1	D	353	VAL	2.3
1	А	138	ARG	2.3
1	А	376	ARG	2.3
1	А	388	LYS	2.2
1	А	283	GLY	2.2
1	В	196	GLY	2.2
1	А	194	PRO	2.2
1	А	280	ALA	2.2
1	С	99	ALA	2.2
1	В	352	SER	2.1
1	D	134	HIS	2.1
1	В	23	GLN	2.1
1	С	212	GLY	2.1
1	С	138	ARG	2.0
1	В	133	ALA	2.0
1	С	353	VAL	2.0
1	В	22	TRP	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	FE	А	501	1/1	1.00	0.10	$22,\!22,\!22,\!22$	0
2	FE	В	501	1/1	1.00	0.08	30,30,30,30	0
2	FE	С	501	1/1	1.00	0.09	27,27,27,27	0
2	FE	D	501	1/1	1.00	0.06	27,27,27,27	0

6.5 Other polymers (i)

There are no such residues in this entry.

