

Full wwPDB X-ray Structure Validation Report (i)

Jan 14, 2024 – 02:15 am GMT

PDB ID : 6SYQ

Title : ASR Alternansucrase in complex with isomaltotriose Authors : Cioci, G.; Molina, M.; Moulis, C.; Remaud-Simeon, M.

Deposited on : 2019-09-30

Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	1275	93%	6%			
2	В	1185	92%	7%			
3	С	3	10	0%			
4	D	2	50%	50%			
4	Е	2	50%	50%			



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 19347 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Alternansucrase.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
1	A	1275	Total 10000	C 6248	N 1703	O 2023	S 26	0	0	0

• Molecule 2 is a protein called Alternansucrase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	1180	Total 9242	C 5776	N 1571	O 1869	S 26	0	0	0

• Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
3	С	3	Total 34	C 18	O 16	0	0	0

• Molecule 4 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	D	2	Total C O 23 12 11	0	0	0
4	Е	2	Total C O 23 12 11	0	0	0



• Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Ca 1 1	0	0
5	В	1	Total Ca 1 1	0	0

• Molecule 6 is alpha-D-glucopyranose (three-letter code: GLC) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 11 6 5	0	0
6	В	1	Total C O 12 6 6	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





 \bullet Molecule 3: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose e



Chain C:	100	9%
GLC2 GLC2 GLC3		
• Molecule	4: alpha-D-glucopyranose-(1-6)	-alpha-D-glucopyranose
Chain D:	50%	50%
GLC1 GLC2		
• Molecule	4: alpha-D-glucopyranose-(1-6)	-alpha-D-glucopyranose
Chain E:	Fox	500/
Cham E.	50%	50%
GLC2		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	100.72Å 135.65Å 238.94Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	119.47 - 3.00	Depositor
Resolution (A)	49.28 - 3.00	EDS
% Data completeness	92.4 (119.47-3.00)	Depositor
(in resolution range)	92.5 (49.28-3.00)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.87 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
D D.	0.201 , 0.237	Depositor
R, R_{free}	0.204 , 0.237	DCC
R_{free} test set	3044 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	86.2	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.29, 46.2	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	19347	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.13% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.54	0/10212	0.67	0/13845	
2	В	0.52	0/9437	0.66	0/12798	
All	All	0.53	0/19649	0.66	0/26643	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10000	0	9417	56	0
2	В	9242	0	8696	55	0
3	С	34	0	30	0	0
4	D	23	0	20	1	0
4	Ε	23	0	21	1	0
5	A	1	0	0	0	0
5	В	1	0	0	0	0
6	A	11	0	10	0	0
6	В	12	0	12	3	0
All	All	19347	0	18206	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 3.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A + a m 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap(Å)
1:A:152:PRO:HA	1:A:180:ASN:O	1.79	0.81
2:B:700:GLN:OE1	4:E:2:GLC:O4	1.99	0.80
1:A:544:LEU:HD13	1:A:592:LEU:HD21	1.64	0.80
2:B:566:ASN:O	2:B:597:ASP:OD2	2.02	0.76
2:B:358:GLU:N	2:B:358:GLU:OE1	2.19	0.76
1:A:553:ASP:OD1	1:A:563:SER:OG	2.05	0.72
1:A:290:ARG:O	1:A:318:GLY:HA2	1.90	0.72
2:B:241:TYR:CD2	6:B:1504:GLC:H4	2.27	0.70
1:A:156:GLY:O	1:A:167:TYR:CE1	2.45	0.69
2:B:271:TYR:HD2	2:B:279:ILE:HG13	1.57	0.69
1:A:472:SER:HB3	1:A:479:THR:HA	1.76	0.68
2:B:371:ASN:O	2:B:372:ASN:HB3	1.93	0.67
1:A:249:THR:HG21	4:D:2:GLC:O2	1.97	0.64
2:B:241:TYR:CE2	6:B:1504:GLC:H4	2.33	0.64
2:B:733:ILE:N	2:B:733:ILE:HD13	2.13	0.63
1:A:543:TRP:CZ3	2:B:287:ASN:HA	2.33	0.63
2:B:874:GLY:HA3	2:B:901:GLY:O	2.02	0.60
2:B:262:LEU:HD23	2:B:286:VAL:CG1	2.32	0.59
1:A:417:PHE:HB2	1:A:1287:ILE:HD13	1.84	0.59
1:A:737:ASN:O	1:A:737:ASN:ND2	2.22	0.59
2:B:290:ARG:O	2:B:318:GLY:HA2	2.03	0.59
2:B:786:ASN:O	2:B:789:ASP:HB2	2.03	0.58
1:A:786:ASN:O	1:A:789:ASP:HB2	2.04	0.58
2:B:358:GLU:CD	2:B:358:GLU:H	2.07	0.58
1:A:544:LEU:CD1	1:A:592:LEU:HD21	2.33	0.57
2:B:271:TYR:CD2	2:B:279:ILE:HG13	2.39	0.57
1:A:543:TRP:HZ3	2:B:287:ASN:HA	1.69	0.57
2:B:892:THR:HG23	2:B:934:LEU:HD21	1.86	0.57
2:B:553:ASP:OD1	2:B:563:SER:OG	2.16	0.57
1:A:892:THR:HG23	1:A:934:LEU:HD21	1.86	0.56
2:B:417:PHE:HB2	2:B:1287:ILE:HD13	1.85	0.56
1:A:163:GLY:O	2:B:352:LYS:HE3	2.05	0.56
2:B:262:LEU:H	2:B:262:LEU:HD12	1.72	0.55
1:A:553:ASP:CG	1:A:563:SER:HG	2.05	0.55
2:B:262:LEU:HD12	2:B:262:LEU:N	2.21	0.55
1:A:541:LEU:HD13	1:A:591:LEU:HD13	1.90	0.54
2:B:544:LEU:HD13	2:B:592:LEU:HD21	1.89	0.54
1:A:231:ASP:HB3	1:A:237:LEU:HD21	1.90	0.54

Continued on next page...



Continued from previous page...

Continued from preva	Atom-1 Atom-2		Clash
Atom-1	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
2:B:469:ASN:HD21	2:B:482:TYR:H	1.54	0.53
2:B:271:TYR:HD2	2:B:279:ILE:CG1	2.21	0.53
1:A:156:GLY:O	1:A:167:TYR:CZ	2.62	0.52
1:A:1222:GLY:O	1:A:1255:LYS:HE2	2.10	0.52
2:B:1222:GLY:O	2:B:1255:LYS:HE2	2.11	0.51
1:A:544:LEU:CD1	1:A:592:LEU:CD2	2.88	0.51
1:A:251:TYR:HD2	1:A:259:LEU:HD12	1.76	0.50
1:A:325:PHE:CE1	1:A:349:ILE:HD11	2.47	0.50
1:A:544:LEU:HD13	1:A:592:LEU:CD2	2.38	0.49
2:B:325:PHE:CE1	2:B:349:ILE:HD11	2.46	0.49
1:A:268:ASN:HD22	1:A:268:ASN:N	2.10	0.49
1:A:541:LEU:HD13	1:A:591:LEU:CD1	2.42	0.49
2:B:812:GLY:HA3	2:B:1010:LEU:HD23	1.95	0.49
2:B:522:PHE:O	2:B:525:THR:HB	2.13	0.48
1:A:290:ARG:O	1:A:318:GLY:CA	2.61	0.48
1:A:558:THR:HG21	1:A:1185:THR:OG1	2.13	0.48
2:B:1097:GLU:HA	2:B:1167:ILE:HB	1.96	0.48
1:A:330:ALA:O	1:A:1410:ALA:HA	2.14	0.48
2:B:262:LEU:HD23	2:B:286:VAL:HG11	1.94	0.48
1:A:163:GLY:O	2:B:352:LYS:CE	2.61	0.47
1:A:1097:GLU:HA	1:A:1167:ILE:HB	1.96	0.47
1:A:522:PHE:O	1:A:525:THR:HB	2.14	0.47
1:A:812:GLY:HA3	1:A:1010:LEU:HD23	1.95	0.47
2:B:732:LYS:C	2:B:733:ILE:HD13	2.33	0.47
2:B:248:GLN:HB3	2:B:277:VAL:HG13	1.95	0.47
2:B:558:THR:HG21	2:B:1185:THR:OG1	2.15	0.46
2:B:330:ALA:O	2:B:1410:ALA:HA	2.15	0.46
2:B:553:ASP:CG	2:B:563:SER:HG	2.18	0.46
1:A:219:ASN:OD1	1:A:244:ASP:HA	2.15	0.46
1:A:271:TYR:CD1	1:A:286:VAL:HG21	2.51	0.46
1:A:175:VAL:CG1	1:A:179:GLN:HG3	2.46	0.45
2:B:1373:MET:HB3	2:B:1374:PRO:CD	2.47	0.45
1:A:238:LYS:HG2	1:A:254:GLY:O	2.16	0.45
1:A:175:VAL:HG11	1:A:179:GLN:HG3	1.99	0.45
1:A:591:LEU:HG	1:A:592:LEU:CD2	2.47	0.45
2:B:996:LYS:HE3	2:B:1002:ILE:HD11	1.99	0.45
1:A:153:ILE:HD11	1:A:181:ILE:HG12	1.99	0.44
1:A:944:LEU:HB3	1:A:946:MET:HE3	1.99	0.44
1:A:1373:MET:HB3	1:A:1374:PRO:CD	2.48	0.44
2:B:731:LYS:O	2:B:733:ILE:HD13	2.17	0.44
2:B:1116:LEU:HA	2:B:1119:PHE:CE2	2.53	0.44

Continued on next page...



Continued from previous page...

Continued from press		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	$overlap (\AA)$
1:A:1063:PHE:CG	1:A:1099:ALA:HB2	2.53	0.43
1:A:205:LYS:HE3	1:A:235:ASN:OD1	2.18	0.43
2:B:470:PHE:CE2	2:B:522:PHE:HB2	2.54	0.43
2:B:659:ASP:OD1	2:B:660:LYS:HE3	2.19	0.43
2:B:283:PHE:HB2	2:B:290:ARG:NH2	2.34	0.43
2:B:894:LEU:HD23	2:B:942:ILE:HD13	2.00	0.43
1:A:284:GLN:HG3	1:A:286:VAL:HG23	2.01	0.43
1:A:1116:LEU:HA	1:A:1119:PHE:CE2	2.54	0.43
1:A:572:GLN:HB2	1:A:573:PRO:HD2	2.01	0.42
1:A:358:GLU:CD	1:A:358:GLU:H	2.22	0.42
2:B:618:GLY:N	2:B:628:ASN:OD1	2.52	0.42
1:A:766:HIS:HD2	1:A:1065:ASN:HD21	1.67	0.42
1:A:894:LEU:HD23	1:A:942:ILE:HD13	2.00	0.42
2:B:563:SER:HB3	2:B:567:ARG:CZ	2.50	0.42
1:A:868:ARG:HA	1:A:872:VAL:HG23	2.02	0.41
1:A:1061:GLU:HA	1:A:1097:GLU:HB3	2.02	0.41
2:B:572:GLN:HB2	2:B:573:PRO:HD2	2.02	0.41
1:A:794:ASP:OD2	1:A:794:ASP:N	2.53	0.41
1:A:566:ASN:HD22	1:A:566:ASN:HA	1.71	0.41
2:B:766:HIS:HD2	2:B:1065:ASN:HD21	1.68	0.41
1:A:553:ASP:CG	1:A:563:SER:OG	2.56	0.41
2:B:771:GLN:HE21	2:B:771:GLN:HB2	1.74	0.41
2:B:898:VAL:HG22	2:B:946:MET:HE1	2.02	0.41
2:B:262:LEU:H	2:B:262:LEU:CD1	2.25	0.40
2:B:406:HIS:HB3	2:B:442:TRP:CZ3	2.57	0.40
1:A:618:GLY:N	1:A:628:ASN:OD1	2.54	0.40
2:B:280:LYS:NZ	6:B:1504:GLC:O2	2.53	0.40
2:B:943:THR:CG2	2:B:985:ASP:HB3	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentile	\mathbf{s}
1	A	1273/1275 (100%)	1228 (96%)	45 (4%)	0	100 100	
2	В	1176/1185 (99%)	1131 (96%)	45 (4%)	0	100 100	
All	All	2449/2460 (100%)	2359 (96%)	90 (4%)	0	100 100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Rotameric Outliers	
1	A	1068/1076 (99%)	1052 (98%)	16 (2%)	65 87
2	В	985/1003 (98%)	967 (98%)	18 (2%)	59 85
All	All	2053/2079 (99%)	2019 (98%)	34 (2%)	60 85

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	268	ASN
1	A	358	GLU
1	A	372	ASN
1	A	460	ASP
1	A	481	THR
1	A	737	ASN
1	A	756	VAL
1	A	789	ASP
1	A	792	THR
1	A	794	ASP
1	A	934	LEU
1	A	998	ASP
1	A	1000	THR
1	A	1169	ASP
1	A	1201	PHE
1	A	1310	SER
2	В	245	SER
2	В	262	LEU

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
2	В	264	THR
2	В	275	GLN
2	В	358	GLU
2	В	419	ASN
2	В	460	ASP
2	В	725	LEU
2	В	743	THR
2	В	756	VAL
2	В	771	GLN
2	В	789	ASP
2	В	792	THR
2	В	794	ASP
2	В	934	LEU
2	В	1000	THR
2	В	1169	ASP
2	В	1201	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	703	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths		Bond ang		gles	
WIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLC	С	1	3	12,12,12	0.61	0	17,17,17	1.06	1 (5%)
3	GLC	С	2	3	11,11,12	0.69	0	15,15,17	1.20	1 (6%)
3	GLC	С	3	3	11,11,12	0.79	0	15,15,17	1.17	2 (13%)
4	GLC	D	1	4	12,12,12	0.64	0	17,17,17	1.85	4 (23%)
4	GLC	D	2	4	11,11,12	0.72	0	15,15,17	1.60	1 (6%)
4	GLC	Е	1	4	12,12,12	0.88	0	17,17,17	1.26	1 (5%)
4	GLC	Е	2	4	11,11,12	0.54	0	15,15,17	1.21	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	С	1	3	-	1/2/22/22	0/1/1/1
3	GLC	С	2	3	-	0/2/19/22	0/1/1/1
3	GLC	С	3	3	-	0/2/19/22	0/1/1/1
4	GLC	D	1	4	-	0/2/22/22	0/1/1/1
4	GLC	D	2	4	-	0/2/19/22	0/1/1/1
4	GLC	Е	1	4	-	0/2/22/22	0/1/1/1
4	GLC	Е	2	4	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
4	D	2	GLC	O5-C5-C6	5.20	115.36	107.20
4	D	1	GLC	C1-O5-C5	-4.62	104.94	113.66
4	D	1	GLC	O1-C1-O5	3.17	119.90	110.38
4	D	1	GLC	O1-C1-C2	3.02	117.54	109.03
4	D	1	GLC	C1-C2-C3	-2.94	104.20	110.31
4	Е	1	GLC	O5-C5-C4	-2.62	104.94	109.69
4	Е	2	GLC	C1-O5-C5	2.45	115.51	112.19
3	С	3	GLC	C2-C3-C4	-2.40	106.74	110.89
3	С	2	GLC	C2-C3-C4	-2.37	106.80	110.89
3	С	3	GLC	C1-O5-C5	2.35	115.37	112.19
3	С	1	GLC	O2-C2-C3	2.06	115.12	110.35



There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	1	GLC	O5-C5-C6-O6
4	Е	2	GLC	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Е	2	GLC	1	0
4	D	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GLC	В	1504	-	12,12,12	0.52	0	17,17,17	1.96	4 (23%)
6	GLC	A	1505	-	11,11,12	0.98	1 (9%)	15,15,17	1.42	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GLC	В	1504	-	-	1/2/22/22	0/1/1/1
6	GLC	A	1505	-	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(ext{\AA})$
6	A	1505	GLC	C2-C3	2.09	1.55	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
6	В	1504	GLC	O1-C1-C2	4.40	121.41	109.03
6	В	1504	GLC	C1-O5-C5	-4.01	106.09	113.66
6	A	1505	GLC	C2-C3-C4	-2.96	105.77	110.89
6	В	1504	GLC	O5-C1-C2	2.72	115.14	110.28
6	В	1504	GLC	O1-C1-O5	2.63	118.27	110.38
6	A	1505	GLC	O3-C3-C2	2.47	114.73	109.99
6	A	1505	GLC	O3-C3-C4	2.30	115.66	110.35

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	В	1504	GLC	O5-C5-C6-O6

There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	В	1504	GLC	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient



equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		-2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	A	$1275/1275\ (100\%)$	-0.24	10 (0%)	86	65	59, 87, 149, 189	0
2	В	$1180/1185\ (99\%)$	-0.22	16 (1%)	75	49	64, 100, 160, 195	0
All	All	$2455/2460\ (99\%)$	-0.23	26 (1%)	80	56	59, 93, 155, 195	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	305	GLU	4.7
2	В	910	THR	3.8
1	A	150	SER	3.2
2	В	245	SER	3.0
2	В	982	ASP	3.0
1	A	326	SER	3.0
1	A	321	VAL	3.0
2	В	941	THR	2.9
2	В	246	SER	2.8
1	A	316	ALA	2.7
1	A	162	ASN	2.6
1	A	172	GLY	2.6
2	В	316	ALA	2.6
2	В	942	ILE	2.6
2	В	286	VAL	2.6
1	A	740	ASN	2.5
2	В	250	TYR	2.5
2	В	1418	VAL	2.4
1	A	362	LEU	2.3
2	В	285	ASP	2.3
2	В	265	ILE	2.3
2	В	309	GLY	2.2
1	A	174	GLN	2.1
1	A	191	ASN	2.1

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	В	1041	SER	2.0
2	В	289	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
4	GLC	D	1	12/12	0.85	0.20	132,148,154,154	0
4	GLC	D	2	11/12	0.85	0.15	122,130,141,164	0
3	GLC	С	1	12/12	0.89	0.16	124,139,147,148	0
4	GLC	Е	1	12/12	0.90	0.13	89,102,106,116	0
4	GLC	Е	2	11/12	0.91	0.17	105,115,119,120	0
3	GLC	С	2	11/12	0.95	0.12	93,99,102,104	0
3	GLC	С	3	11/12	0.96	0.17	84,96,100,105	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around Chain D:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
6	GLC	В	1504	12/12	0.71	0.20	135,147,152,160	0
6	GLC	A	1505	11/12	0.87	0.19	128,143,148,152	0
5	CA	В	1501	1/1	0.96	0.19	88,88,88,88	0
5	CA	A	1501	1/1	0.97	0.23	79,79,79,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

