

Full wwPDB X-ray Structure Validation Report (i)

Oct 7, 2023 – 08:51 AM EDT

PDB ID : 6VP3

Title: Artificial Metalloproteins with Dinuclear Iron Centers

Authors: Miller, K.R.; Follmer, A.H.; Jasniewski, A.J.; Sabuncu, S.; Biswas, S.; Albert,

T.; Hendrich, M.P.; Moenne-Loccoz, P.; Borovik, A.S.

Deposited on : 2020-02-01

Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	159	70%	5%		23%	
1	В	159	70%	6%	•	23%	
1	С	159	69%	6%	•	23%	
1	D	159	68%	8%	•	23%	



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Streptavidin.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace
1	A	122	Total	С	N	О	0	3	0
1	A	122	923	573	161	189	0	3	0
1	В	122	Total	С	N	О	0	3	0
1	Ъ		923	573	161	189		3	
1	С	122	Total	С	N	О	0	3	0
1		122	923	573	161	189	U	9	U
1	D	D 100	Total	С	N	О	0	3	0
	122	923	573	161	189	U	3	U	

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P22629
A	2	ALA	-	- expression tag	
A	3	SER	-	expression tag	UNP P22629
A	4	MET	-	expression tag	UNP P22629
A	5	THR	-	expression tag	UNP P22629
A	6	GLY	-	expression tag	UNP P22629
A	7	GLY	-	expression tag	UNP P22629
A	8	GLN	-	expression tag	UNP P22629
A	9	GLN	-	expression tag	UNP P22629
A	10	MET	-	expression tag	UNP P22629
A	11	GLY	-	expression tag	UNP P22629
A	12	ARG	-	expression tag	UNP P22629
A	13	ASP	-	expression tag	UNP P22629
A	121	ALA	LYS	engineered mutation	UNP P22629
A	124	TYR	LEU	engineered mutation	UNP P22629
В	1	MET	-	initiating methionine	UNP P22629
В	2	ALA	-	expression tag	UNP P22629
В	3	SER	-	expression tag	UNP P22629
В	4	MET	-	expression tag	UNP P22629
В	5	THR	-	expression tag	UNP P22629
В	6	GLY	-	expression tag	UNP P22629

Continued on next page...




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	7	GLY	-	expression tag	UNP P22629
В	8	GLN	-	expression tag	UNP P22629
В	9	GLN	-	expression tag	UNP P22629
В	10	MET	-	expression tag	UNP P22629
В	11	GLY	-	expression tag	UNP P22629
В	12	ARG	-	expression tag	UNP P22629
В	13	ASP	-	expression tag	UNP P22629
В	121	ALA	LYS	engineered mutation	UNP P22629
В	124	TYR	LEU	engineered mutation	UNP P22629
С	1	MET	-	initiating methionine	UNP P22629
С	2	ALA	-	expression tag	UNP P22629
С	3	SER	-	expression tag	UNP P22629
С	4	MET	-	expression tag	UNP P22629
С	5	THR	-	expression tag	UNP P22629
С	6	GLY	-	expression tag	UNP P22629
С	7	GLY	-	expression tag	UNP P22629
С	8	GLN	-	expression tag	UNP P22629
С	9	GLN	-	expression tag	UNP P22629
С	10	MET	-	expression tag	UNP P22629
С	11	GLY	-	expression tag	UNP P22629
С	12	ARG	-	expression tag	UNP P22629
С	13	ASP	-	expression tag	UNP P22629
С	121	ALA	LYS	engineered mutation	UNP P22629
С	124	TYR	LEU	engineered mutation	UNP P22629
D	1	MET	-	initiating methionine	UNP P22629
D	2	ALA	-	expression tag	UNP P22629
D	3	SER	-	expression tag	UNP P22629
D	4	MET	-	expression tag	UNP P22629
D	5	THR	-	expression tag	UNP P22629
D	6	GLY	-	expression tag	UNP P22629
D	7	GLY	-	expression tag	UNP P22629
D	8	GLN	-	expression tag	UNP P22629
D	9	GLN	-	expression tag	UNP P22629
D	10	MET	-	expression tag	UNP P22629
D	11	GLY	-	expression tag	UNP P22629
D	12	ARG	-	expression tag	UNP P22629
D	13	ASP	-	expression tag	UNP P22629
D	121	ALA	LYS	engineered mutation	UNP P22629
D	124	TYR	LEU	engineered mutation	UNP P22629


• Molecule 2 is $\{N-(4-\{bis[(pyridin-2-yl-kappaN)methyl]amino-kappaN\}butyl)-5-[(3aS,4S,6aR)-2-oxohexahydro-1H-thieno[3,4-d]imidazol-4-yl]pentanamide<math>\}iron(3+)$ (three-letter code: KM3) (formula: $C_{26}H_{36}FeN_6O_2S$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	С	Fe	N	О	S	0	1
	Λ	1	36	26	1	6	2	1	0	1
2	В	1	Total	С	Fe	N	О	S	0	1
	Ъ	1	36	26	1	6	2	1		1
2	С	1	Total	С	Fe	N	О	S	0	1
		1	36	26	1	6	2	1	0	1
9	9 D	1	Total	С	Fe	N	О	S	0	1
	1	36	26	1	6	2	1		1	

 \bullet Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0


\bullet Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	125	Total O 125 125	0	0
4	В	115	Total O 115 115	0	0
4	С	127	Total O 127 127	0	0
4	D	132	Total O 132 132	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	193.50Å 57.86Å 57.86Å	Donogiton
a, b, c, α , β , γ	90.00° 107.40° 90.00°	Depositor
Resolution (Å)	42.19 - 1.65	Depositor
Resolution (A)	42.16 - 1.65	EDS
% Data completeness	64.6 (42.19-1.65)	Depositor
(in resolution range)	$64.6 \ (42.16 - 1.65)$	EDS
R_{merge}	0.07	Depositor
R_{sum}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.41 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
P. P.	0.176 , 0.221	Depositor
R, R_{free}	0.188 , 0.228	DCC
R_{free} test set	2442 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35 , 37.1	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.34$	Xtriage
	0.448 for -h-k-l,l,k	
Estimated twinning fraction	$0.459 \ { m for -h+k-l,-l,-k}$	Xtriage
	0.458 for -h-2*l,-k,l	
F_o, F_c correlation	0.96	EDS
Total number of atoms	4343	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.19% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: KM3, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.76	0/963	1.10	2/1318~(0.2%)	
1	В	0.77	0/963	1.18	$2/1318 \; (0.2\%)$	
1	С	0.80	0/963	1.24	2/1318~(0.2%)	
1	D	0.79	0/963	1.10	2/1318 (0.2%)	
All	All	0.78	0/3852	1.16	8/5272 (0.2%)	

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	С	53[A]	ARG	CG-CD-NE	-21.15	67.39	111.80
1	С	53[B]	ARG	CG-CD-NE	-21.15	67.39	111.80
1	В	53[A]	ARG	CG-CD-NE	-17.78	74.46	111.80
1	В	53[B]	ARG	CG-CD-NE	-17.78	74.46	111.80
1	D	53[A]	ARG	CG-CD-NE	-15.72	78.78	111.80
1	D	53[B]	ARG	CG-CD-NE	-15.72	78.78	111.80
1	A	53[A]	ARG	CG-CD-NE	-15.39	79.48	111.80
1	A	53[B]	ARG	CG-CD-NE	-15.39	79.48	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	923	0	848	10	0
1	В	923	0	848	11	0
1	С	923	0	848	14	0
1	D	923	0	848	9	0
2	A	36	0	0	0	0
2	В	36	0	0	0	0
2	С	36	0	0	0	0
2	D	36	0	0	0	0
3	A	4	0	3	0	0
3	В	4	0	3	0	0
4	A	125	0	0	2	0
4	В	115	0	0	4	0
4	С	127	0	0	2	0
4	D	132	0	0	2	0
All	All	4343	0	3398	39	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:40[A]:THR:HG22	4:B:304:HOH:O	1.73	0.86
1:B:40[B]:THR:HG22	4:B:304:HOH:O	1.73	0.86
1:B:133:VAL:O	1:B:134:LYS:HB2	1.80	0.80
1:B:134:LYS:HG2	4:B:379:HOH:O	1.81	0.79
1:C:103:ARG:CG	1:C:103:ARG:HH21	2.07	0.66
1:C:40[A]:THR:HG22	4:C:367:HOH:O	1.94	0.66
1:C:40[B]:THR:HG22	4:C:367:HOH:O	1.94	0.66
1:B:133:VAL:O	1:B:134:LYS:CB	2.44	0.65
1:A:40[A]:THR:HG22	4:A:368:HOH:O	2.00	0.61
1:A:40[B]:THR:HG22	4:A:368:HOH:O	2.00	0.61
1:A:52:SER:HG	1:A:53[B]:ARG:HH11	1.46	0.61
1:C:110:LEU:HD22	1:C:124:TYR:HD2	1.68	0.58
1:C:110:LEU:C	1:C:110:LEU:HD23	2.26	0.56
1:C:103:ARG:CG	1:C:103:ARG:NH2	2.67	0.55
1:D:110:LEU:C	1:D:110:LEU:HD23	2.27	0.55
1:C:103:ARG:NH2	1:C:103:ARG:HG2	2.22	0.54
1:A:52:SER:OG	1:A:53[B]:ARG:NH1	2.37	0.54
1:A:133:VAL:O	1:A:134:LYS:HB2	2.08	0.53
1:A:133:VAL:O	1:A:134:LYS:CB	2.58	0.52
1:A:91:THR:HB	1:C:91:THR:HB	1.91	0.52

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance} (\mathrm{\AA})$	overlap (Å)
1:C:107[B]:GLN:NE2	1:D:126:GLY:HA2	2.25	0.51
1:C:126:GLY:HA2	1:D:107[B]:GLN:NE2	2.27	0.49
1:D:13:ASP:OD2	1:D:71:THR:OG1	2.18	0.49
1:B:103:ARG:HH21	1:B:103:ARG:HG3	1.77	0.49
1:B:30:ILE:CD1	1:B:53[A]:ARG:NH1	2.76	0.49
1:B:91:THR:HB	1:D:91:THR:HB	1.95	0.48
1:B:110:LEU:C	1:B:110:LEU:HD23	2.34	0.48
1:B:97:VAL:HG23	4:B:307:HOH:O	2.15	0.47
1:A:103:ARG:HG3	1:A:103:ARG:HH21	1.81	0.46
1:A:110:LEU:C	1:A:110:LEU:HD23	2.36	0.46
1:C:110:LEU:HD22	1:C:124:TYR:CD2	2.51	0.45
1:D:53[B]:ARG:CZ	4:D:360:HOH:O	2.64	0.45
1:D:82:ASN:ND2	4:D:302:HOH:O	2.51	0.42
1:D:23:ASN:HB3	1:D:130:PHE:CE1	2.54	0.42
1:C:103:ARG:HH21	1:C:103:ARG:HG3	1.84	0.42
1:C:23:ASN:HB3	1:C:130:PHE:CE2	2.55	0.41
1:A:107[B]:GLN:NE2	1:B:126:GLY:HA2	2.36	0.41
1:D:22:TYR:O	1:D:130:PHE:HA	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	123/159 (77%)	120 (98%)	3 (2%)	0	100	100
1	В	123/159 (77%)	120 (98%)	3 (2%)	0	100	100
1	С	123/159 (77%)	118 (96%)	5 (4%)	0	100	100
1	D	123/159 (77%)	120 (98%)	3 (2%)	0	100	100
All	All	492/636 (77%)	478 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	92/114 (81%)	90 (98%)	2 (2%)	52 27
1	В	92/114 (81%)	92 (100%)	0	100 100
1	С	92/114 (81%)	89 (97%)	3 (3%)	38 12
1	D	92/114 (81%)	89 (97%)	3 (3%)	38 12
All	All	368/456 (81%)	360 (98%)	8 (2%)	52 27

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	LEU
1	A	133	VAL
1	С	103	ARG
1	С	133	VAL
1	С	134	LYS
1	D	73	LEU
1	D	133	VAL
1	D	134	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	KM3	D	201[A]	4,3	30,41,41	2.53	8 (26%)	41,59,59	2.71	19 (46%)
2	KM3	A	201[D]	4,1,3	30,41,41	2.67	6 (20%)	41,59,59	2.50	14 (34%)
3	ACT	A	202	2	3,3,3	0.88	0	3,3,3	0.67	0
2	KM3	С	201[C]	1,3	30,41,41	2.65	6 (20%)	41,59,59	2.25	10 (24%)
2	KM3	В	201[B]	1,3	30,41,41	2.46	6 (20%)	41,59,59	2.73	15 (36%)
3	ACT	В	202	2	3,3,3	0.44	0	3,3,3	1.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KM3	В	201[B]	1,3	-	1/13/61/61	0/6/6/6
2	KM3	D	201[A]	4,3	-	1/13/61/61	0/6/6/6
2	KM3	A	201[D]	4,1,3	-	2/13/61/61	0/6/6/6
2	KM3	С	201[C]	1,3	-	1/13/61/61	0/6/6/6

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
2	С	201[C]	KM3	C14-C13	-8.99	1.20	1.38
2	A	201[D]	KM3	C14-C13	-8.62	1.20	1.38
2	D	201[A]	KM3	C14-C13	-7.76	1.22	1.38
2	В	201[B]	KM3	C14-C13	-7.64	1.22	1.38
2	A	201[D]	KM3	C13-C12	-7.02	1.27	1.39
2	С	201[C]	KM3	C13-C12	-6.39	1.28	1.39
2	В	201[B]	KM3	C13-C12	-6.20	1.28	1.39
2	С	201[C]	KM3	C14-C15	-6.09	1.21	1.38
2	D	201[A]	KM3	C14-C15	-5.98	1.22	1.38

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	D	201[A]	KM3	C13-C12	-5.94	1.29	1.39
2	В	201[B]	KM3	C14-C15	-5.80	1.22	1.38
2	A	201[D]	KM3	C14-C15	-5.77	1.22	1.38
2	С	201[C]	KM3	C16-C15	-4.22	1.29	1.38
2	A	201[D]	KM3	C16-C15	-4.20	1.29	1.38
2	A	201[D]	KM3	C6-S1	-3.85	1.76	1.82
2	В	201[B]	KM3	C01-N6	3.75	1.41	1.33
2	В	201[B]	KM3	C16-C15	-3.74	1.30	1.38
2	D	201[A]	KM3	C6-S1	-3.63	1.76	1.82
2	D	201[A]	KM3	C16-C15	-3.34	1.31	1.38
2	С	201[C]	KM3	C01-N6	3.09	1.40	1.33
2	A	201[D]	KM3	C01-N6	3.03	1.40	1.33
2	D	201[A]	KM3	C01-N6	2.82	1.39	1.33
2	D	201[A]	KM3	C8-N1	2.41	1.49	1.46
2	С	201[C]	KM3	C6-S1	-2.27	1.79	1.82
2	D	201[A]	KM3	C2-C01	2.23	1.55	1.51
2	В	201[B]	KM3	O1-C01	2.17	1.27	1.23

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}({}^o)$	$\operatorname{Ideal}({}^{o})$
2	В	201[B]	KM3	C7-C8-N1	-8.88	101.74	113.03
2	D	201[A]	KM3	C16-N4-C12	-8.48	112.16	118.23
2	С	201[C]	KM3	C16-N4-C12	-7.71	112.71	118.23
2	A	201[D]	KM3	C10-C6-S1	7.19	112.05	105.20
2	В	201[B]	KM3	C6-C10-N2	-6.94	106.91	113.13
2	A	201[D]	KM3	C15-C16-N4	-5.63	118.76	122.93
2	A	201[D]	KM3	C6-C10-N2	-5.56	108.14	113.13
2	D	201[A]	KM3	C8-C7-S1	5.32	110.86	106.31
2	D	201[A]	KM3	C7-C8-C10	-5.11	104.22	108.66
2	A	201[D]	KM3	C16-N4-C12	-5.09	114.59	118.23
2	С	201[C]	KM3	C8-C7-S1	4.93	110.53	106.31
2	В	201[B]	KM3	C19-C18-N5	-4.78	118.46	121.78
2	D	201[A]	KM3	C10-C6-S1	4.76	109.74	105.20
2	A	201[D]	KM3	C6-C10-C8	-4.71	103.47	108.94
2	D	201[A]	KM3	C19-C18-N5	-4.64	118.56	121.78
2	В	201[B]	KM3	C10-C6-S1	4.30	109.30	105.20
2	В	201[B]	KM3	C15-C16-N4	-4.28	119.76	122.93
2	С	201[C]	KM3	C17-C18-C19	4.18	131.40	122.66
2	В	201[B]	KM3	C17-C18-C19	4.16	131.36	122.66
2	С	201[C]	KM3	C7-C8-C10	-4.11	105.09	108.66
2	D	201[A]	KM3	C17-C18-C19	3.89	130.79	122.66

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
2	В	201[B]	KM3	C8-C7-S1	3.73	109.50	106.31
2	С	201[C]	KM3	C15-C16-N4	-3.58	120.28	122.93
2	С	201[C]	KM3	C6-C10-N2	-3.54	109.95	113.13
2	D	201[A]	KM3	C6-C10-N2	-3.51	109.98	113.13
2	A	201[D]	KM3	C17-C18-C19	3.49	129.96	122.66
2	В	201[B]	KM3	C16-N4-C12	-3.46	115.75	118.23
2	D	201[A]	KM3	C11-C12-C13	3.33	129.63	122.66
2	D	201[A]	KM3	C8-C10-N2	-3.30	99.12	102.67
2	D	201[A]	KM3	O3-C9-N1	-3.27	121.24	125.94
2	С	201[C]	KM3	C7-C8-N1	-3.12	109.07	113.03
2	С	201[C]	KM3	C16-C15-C14	3.10	123.48	118.85
2	С	201[C]	KM3	C10-C6-S1	3.03	108.09	105.20
2	С	201[C]	KM3	C19-C18-N5	-2.99	119.70	121.78
2	A	201[D]	KM3	C16-C15-C14	2.98	123.31	118.85
2	A	201[D]	KM3	C21-C22-N5	-2.96	120.74	122.93
2	В	201[B]	KM3	C11-C12-C13	2.88	128.69	122.66
2	В	201[B]	KM3	C13-C12-N4	-2.86	119.79	121.78
2	В	201[B]	KM3	C6-C10-C8	-2.74	105.76	108.94
2	В	201[B]	KM3	C22-N5-C18	2.64	120.12	118.23
2	D	201[A]	KM3	N2-C9-N1	2.61	111.21	108.76
2	A	201[D]	KM3	C7-S1-C6	-2.58	84.59	89.89
2	D	201[A]	KM3	C7-S1-C6	-2.57	84.61	89.89
2	D	201[A]	KM3	C22-N5-C18	2.54	120.05	118.23
2	D	201[A]	KM3	C6-C10-C8	-2.49	106.05	108.94
2	A	201[D]	KM3	C4-C5-C6	2.38	118.70	113.86
2	В	201[B]	KM3	C10-C8-N1	-2.37	99.91	102.43
2	D	201[A]	KM3	O1-C01-N6	-2.34	118.59	123.01
2	D	201[A]	KM3	C7-C8-N1	-2.32	110.09	113.03
2	A	201[D]	KM3	C11-C12-C13	2.22	127.30	122.66
2	A	201[D]	KM3	C8-C7-S1	2.09	108.10	106.31
2	D	201[A]	KM3	C14-C13-C12	2.07	121.76	119.18
2	D	201[A]	KM3	C10-N2-C9	2.07	114.54	112.62
2	A	201[D]	KM3	O1-C01-N6	-2.05	119.14	123.01
2	В	201[B]	KM3	C16-C15-C14	2.05	121.92	118.85
2	D	201[A]	KM3	C16-C15-C14	2.03	121.89	118.85
2	A	201[D]	KM3	O3-C9-N2	-2.03	123.02	125.94
2	В	201[B]	KM3	C10-N2-C9	-2.00	110.75	112.62

There are no chirality outliers.

All (5) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
2	A	201[D]	KM3	C4-C5-C6-C10
2	В	201[B]	KM3	C4-C5-C6-C10
2	A	201[D]	KM3	C24-C25-C26-C23
2	D	201[A]	KM3	C24-C25-C26-C23
2	С	201[C]	KM3	C24-C25-C26-C23


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	122/159~(76%)	0.13	1 (0%) 86 88	8, 14, 32, 54	3 (2%)
1	В	122/159~(76%)	0.15	1 (0%) 86 88	8, 14, 34, 50	3 (2%)
1	С	122/159 (76%)	0.08	3 (2%) 57 58	8, 14, 32, 50	3 (2%)
1	D	122/159 (76%)	0.12	2 (1%) 72 75	8, 14, 32, 46	3 (2%)
All	All	488/636 (76%)	0.12	7 (1%) 75 79	8, 14, 34, 54	12 (2%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	100	ALA	3.1
1	С	100	ALA	2.7
1	D	66	THR	2.7
1	С	124	TYR	2.3
1	A	99	GLY	2.3
1	D	100	ALA	2.1
1	С	15	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	ACT	В	202	4/4	0.81	0.16	22,26,27,31	0
3	ACT	A	202	4/4	0.85	0.14	22,30,31,33	0
2	KM3	D	201[A]	36/36	0.95	0.11	8,17,44,46	0
2	KM3	A	201[D]	36/36	0.96	0.11	8,18,39,41	0
2	KM3	В	201[B]	36/36	0.96	0.10	8,21,40,43	0
2	KM3	С	201[C]	36/36	0.96	0.10	9,19,39,40	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around KM3 A 201 (D):

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around KM3 B 201 (B):


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

