



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 03:49 PM EDT

PDB ID : 7A1A  
Title : 2,3-Dihydroxybenzoate Decarboxylase of *Aspergillus oryzae*  
Authors : Hofer, G.; Keller, W.  
Deposited on : 2020-08-12  
Resolution : 1.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	
1	B	358	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11561 atoms, of which 5303 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amidohydrolase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	339	Total	C	H	N	O	S	0	2	0
			5420	1772	2653	470	512	13			
1	B	339	Total	C	H	N	O	S	0	3	0
			5423	1776	2650	468	516	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A1S9DW14
A	-18	GLY	-	expression tag	UNP A0A1S9DW14
A	-17	SER	-	expression tag	UNP A0A1S9DW14
A	-16	SER	-	expression tag	UNP A0A1S9DW14
A	-15	HIS	-	expression tag	UNP A0A1S9DW14
A	-14	HIS	-	expression tag	UNP A0A1S9DW14
A	-13	HIS	-	expression tag	UNP A0A1S9DW14
A	-12	HIS	-	expression tag	UNP A0A1S9DW14
A	-11	HIS	-	expression tag	UNP A0A1S9DW14
A	-10	HIS	-	expression tag	UNP A0A1S9DW14
A	-9	SER	-	expression tag	UNP A0A1S9DW14
A	-8	SER	-	expression tag	UNP A0A1S9DW14
A	-7	GLY	-	expression tag	UNP A0A1S9DW14
A	-6	LEU	-	expression tag	UNP A0A1S9DW14
A	-5	VAL	-	expression tag	UNP A0A1S9DW14
A	-4	PRO	-	expression tag	UNP A0A1S9DW14
A	-3	ARG	-	expression tag	UNP A0A1S9DW14
A	-2	GLY	-	expression tag	UNP A0A1S9DW14
A	-1	SER	-	expression tag	UNP A0A1S9DW14
A	0	HIS	-	expression tag	UNP A0A1S9DW14
B	-19	MET	-	initiating methionine	UNP A0A1S9DW14
B	-18	GLY	-	expression tag	UNP A0A1S9DW14
B	-17	SER	-	expression tag	UNP A0A1S9DW14
B	-16	SER	-	expression tag	UNP A0A1S9DW14
B	-15	HIS	-	expression tag	UNP A0A1S9DW14

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP A0A1S9DW14
B	-13	HIS	-	expression tag	UNP A0A1S9DW14
B	-12	HIS	-	expression tag	UNP A0A1S9DW14
B	-11	HIS	-	expression tag	UNP A0A1S9DW14
B	-10	HIS	-	expression tag	UNP A0A1S9DW14
B	-9	SER	-	expression tag	UNP A0A1S9DW14
B	-8	SER	-	expression tag	UNP A0A1S9DW14
B	-7	GLY	-	expression tag	UNP A0A1S9DW14
B	-6	LEU	-	expression tag	UNP A0A1S9DW14
B	-5	VAL	-	expression tag	UNP A0A1S9DW14
B	-4	PRO	-	expression tag	UNP A0A1S9DW14
B	-3	ARG	-	expression tag	UNP A0A1S9DW14
B	-2	GLY	-	expression tag	UNP A0A1S9DW14
B	-1	SER	-	expression tag	UNP A0A1S9DW14
B	0	HIS	-	expression tag	UNP A0A1S9DW14

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	363	Total O 363 363	0	0
4	B	352	Total O 352 352	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amidohydrolase 2



- Molecule 1: Amidohydrolase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.89Å 99.89Å 129.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.95 – 1.53 49.95 – 1.53	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.95-1.53) 99.6 (49.95-1.53)	Depositor EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.31 (at 1.53Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.192 , 0.225 0.193 , 0.226	Depositor DCC
$R_{free}$ test set	4845 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.4	Xtriage
Anisotropy	0.172	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 36.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11561	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/2849	0.56	0/3863
1	B	0.38	0/2861	0.55	0/3880
All	All	0.39	0/5710	0.56	0/7743

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2767	2653	2656	8	0
1	B	2773	2650	2657	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	1	0	0	0	0
4	A	363	0	0	2	3
4	B	352	0	0	2	4
All	All	6258	5303	5313	15	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285[A]:SER:OG	4:A:501:HOH:O	2.16	0.62
1:B:164:PHE:HB3	1:B:218:ILE:HD13	1.91	0.52
1:A:253:LYS:HE2	4:A:600:HOH:O	2.11	0.49
1:B:81:GLU:OE2	4:B:501:HOH:O	2.20	0.48
1:B:167:HIS:CG	1:B:168:PRO:HD2	2.53	0.44
1:A:266:THR:HA	1:A:289:LEU:O	2.19	0.43
1:A:8:GLU:O	1:A:293:ASP:HA	2.18	0.43
1:B:317:LEU:HG	4:B:652:HOH:O	2.19	0.43
1:A:188:GLY:HA2	1:A:193:PHE:CE2	2.54	0.42
1:A:225:GLU:O	1:A:226:HIS:HB2	2.19	0.42
1:A:158:THR:HB	1:A:215:LYS:HE3	2.00	0.42
1:B:292:ILE:O	1:B:293:ASP:CB	2.69	0.41
1:B:318:LYS:HA	1:B:322:GLU:HB2	2.03	0.41
1:A:271:PHE:CZ	1:A:302:ALA:HB2	2.56	0.40
1:B:189:PRO:N	1:B:190:PRO:CD	2.84	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:758:HOH:O	4:B:659:HOH:O[3_654]	2.08	0.12
4:A:743:HOH:O	4:A:808:HOH:O[8_667]	2.10	0.10
4:B:528:HOH:O	4:B:528:HOH:O[8_667]	2.10	0.10
4:A:717:HOH:O	4:B:555:HOH:O[6_577]	2.16	0.04
4:B:675:HOH:O	4:B:712:HOH:O[8_667]	2.19	0.01

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/358 (95%)	324 (96%)	14 (4%)	1 (0%)	41	19
1	B	340/358 (95%)	325 (96%)	14 (4%)	1 (0%)	41	19
All	All	679/716 (95%)	649 (96%)	28 (4%)	2 (0%)	41	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	191	LEU
1	B	191	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/307 (95%)	290 (99%)	3 (1%)	76	55
1	B	294/307 (96%)	293 (100%)	1 (0%)	92	84
All	All	587/614 (96%)	583 (99%)	4 (1%)	84	68

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	40	ASP
1	A	165	TYR
1	B	7	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	277	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	339/358 (94%)	-0.39	0 100 100	10, 15, 29, 58	0
1	B	339/358 (94%)	-0.38	0 100 100	11, 16, 29, 55	0
All	All	678/716 (94%)	-0.39	0 100 100	10, 16, 29, 58	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	401	1/1	0.95	0.05	16,16,16,16	1
2	MG	A	401	1/1	0.99	0.06	16,16,16,16	0
3	CA	B	402	1/1	1.00	0.04	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.




**Electron density around MG B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MG A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.