



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2022 – 09:17 am GMT

PDB ID : 7A6T
Title : Crystal Structure of Asn173Ser variant of Human Deoxyhypusine Synthase in complex with NAD and spermidine
Authors : Wator, E.; Wilk, P.; Grudnik, P.
Deposited on : 2020-08-26
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.27
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	
1	B	369	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SPD	B	1602	-	-	X	-
5	BME	A	406	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 11380 atoms, of which 5500 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Deoxyhypusine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	336	5381	1720	2685	456	503	17	0	17	0
1	B	328	5343	1705	2681	446	494	17	0	20	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	173	SER	ASN	engineered mutation	UNP P49366
B	173	SER	ASN	engineered mutation	UNP P49366

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	A	1	70	21	26	7	14	2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	B	1	70	21	26	7	14	2	0	0

- Molecule 3 is SPERMIDINE (three-letter code: SPD) (formula: C₇H₁₉N₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	N		
3	A	1	29	7	19	3	0	0
3	B	1	29	7	19	3	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	10	2	6	2	0	0
4	A	1	10	2	6	2	0	0
4	A	1	10	2	6	2	0	0
4	A	1	10	2	6	2	0	0
4	A	1	10	2	6	2	0	0
4	B	1	10	2	6	2	0	0

- Molecule 5 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	S		
5	A	1	10	2	6	1	1	0	0

- Molecule 6 is OXAMIC ACID (three-letter code: OXM) (formula: $C_2H_3NO_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
6	B	1	8	2	2	1	3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	191	Total 191	O 191	0	0
7	B	189	Total 189	O 189	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deoxyhypusine synthase



- Molecule 1: Deoxyhypusine synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.03Å 105.03Å 160.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.01 – 1.66 46.01 – 1.66	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.01-1.66) 99.8 (46.01-1.66)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 1.66Å)	Xtrriage
Refinement program	PHENIX 1.17.1	Depositor
R, R_{free}	0.171 , 0.193 0.171 , 0.193	Depositor DCC
R_{free} test set	2099 reflections (1.74%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtrriage
Anisotropy	0.538	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	11380	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SPD, BME, EDO, CSS, NAD, OXM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	2/2798 (0.1%)	0.69	2/3792 (0.1%)
1	B	0.55	1/2770 (0.0%)	0.71	1/3754 (0.0%)
All	All	0.54	3/5568 (0.1%)	0.70	3/7546 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	40	VAL	CB-CG1	-11.34	1.29	1.52
1	A	284	GLY	C-N	-8.31	1.15	1.34
1	A	285	VAL	CA-C	6.04	1.68	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	40	VAL	CA-CB-CG1	-11.58	93.54	110.90
1	A	243	ASP	CB-CG-OD2	-6.12	112.79	118.30
1	A	284	GLY	C-N-CA	5.84	136.30	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2696	2685	2645	23	0
1	B	2662	2681	2641	16	0
2	A	44	26	26	2	0
2	B	44	26	26	0	0
3	A	10	19	19	5	0
3	B	10	19	19	8	0
4	A	20	30	30	5	0
4	B	4	6	6	0	0
5	A	4	6	6	3	0
6	B	6	2	2	0	0
7	A	191	0	0	3	0
7	B	189	0	0	5	0
All	All	5880	5500	5420	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:HIS:NE2	3:A:402:SPD:H51	1.84	0.93
1:A:320:ARG:HD2	4:A:405:EDO:H21	1.55	0.89
1:A:261:GLU:OE2	5:A:406:BME:S2	2.44	0.75
1:A:329:LYS:NZ	3:A:402:SPD:H52	2.05	0.72
1:B:137[B]:GLU:OE1	7:B:1701:HOH:O	2.12	0.66
1:A:43:ARG:NH1	1:A:357:GLN:CD	2.53	0.62
7:A:595:HOH:O	1:B:296:MET:HE2	2.02	0.60
1:B:288:HIS:NE2	3:B:1602:SPD:H52	2.16	0.60
1:A:320:ARG:HH11	4:A:405:EDO:H21	1.67	0.60
1:B:329:LYS:HZ1	3:B:1602:SPD:H51	1.67	0.58
1:B:329:LYS:NZ	3:B:1602:SPD:H51	2.19	0.57
1:A:329:LYS:HZ1	3:A:402:SPD:H52	1.71	0.55
1:B:358:LYS:HG3	7:B:1863:HOH:O	2.06	0.55
1:A:77:LEU:HD13	1:A:78:SER:N	2.24	0.53
1:A:164:ASN:HB2	1:A:171[B]:VAL:HG23	1.90	0.52
4:A:407:EDO:H22	7:A:549:HOH:O	2.10	0.51
1:A:329:LYS:HZ3	3:A:402:SPD:H52	1.76	0.50
4:A:408:EDO:H21	7:B:1758:HOH:O	2.11	0.50
1:A:307:ASN:HA	2:A:401:NAD:H2A	1.96	0.48
4:A:408:EDO:O2	3:B:1602:SPD:N10	2.47	0.48
1:B:92:ARG:NH1	1:B:123:ASN:OD1	2.30	0.48
1:B:327:TRP:CZ3	3:B:1602:SPD:H72	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:327:TRP:CZ2	3:B:1602:SPD:H31	2.50	0.47
1:A:83:GLN:HA	1:A:83:GLN:OE1	2.16	0.46
1:B:329:LYS:HZ3	3:B:1602:SPD:C5	2.29	0.45
1:A:43:ARG:HH11	1:A:357:GLN:NE2	2.14	0.45
1:B:329:LYS:NZ	3:B:1602:SPD:C5	2.79	0.45
1:B:283:GLY:O	1:B:286[B]:VAL:HG22	2.16	0.45
1:B:137[B]:GLU:OE2	7:B:1702:HOH:O	2.21	0.44
1:A:268:THR:HG21	5:A:406:BME:H12	2.00	0.44
1:A:137:GLU:OE2	2:A:401:NAD:H4N	2.17	0.44
1:B:92:ARG:HD3	7:B:1821:HOH:O	2.18	0.43
1:A:43:ARG:NH1	1:A:357:GLN:NE2	2.66	0.42
1:A:123:ASN:ND2	1:A:227:ASN:HB3	2.33	0.42
1:A:283:GLY:O	1:A:286[B]:VAL:HG22	2.19	0.42
1:B:268[B]:THR:HG23	1:B:272:PHE:CD2	2.54	0.41
1:A:32:ARG:NH1	7:A:510:HOH:O	2.53	0.41
1:A:123:ASN:HD21	1:A:227:ASN:HB3	1.85	0.41
1:A:327:TRP:CE2	1:B:164:ASN:HB3	2.55	0.41
1:A:265:LEU:HD23	5:A:406:BME:H22	2.03	0.40
1:A:323:GLU:OE1	3:A:402:SPD:H91	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/369 (95%)	345 (99%)	5 (1%)	0	100	100
1	B	343/369 (93%)	337 (98%)	6 (2%)	0	100	100
All	All	693/738 (94%)	682 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/309 (95%)	292 (100%)	1 (0%)	92	88
1	B	293/309 (95%)	293 (100%)	0	100	100
All	All	586/618 (95%)	585 (100%)	1 (0%)	93	89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSS	A	177	1	4,6,7	0.63	0	1,6,8	0.14	0
1	CSS	B	177	1	4,6,7	1.14	0	1,6,8	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	177	1	-	0/1/5/7	-
1	CSS	B	177	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SPD	B	1602	-	9,9,9	0.19	0	8,8,8	0.32	0
4	EDO	B	1604	-	3,3,3	0.45	0	2,2,2	0.50	0
6	OXM	B	1603	-	2,5,5	0.47	0	2,6,6	2.72	1 (50%)
2	NAD	B	1601	-	42,48,48	1.74	12 (28%)	50,73,73	1.54	11 (22%)
3	SPD	A	402	-	9,9,9	0.79	0	8,8,8	0.96	0
4	EDO	A	407	-	3,3,3	0.33	0	2,2,2	0.37	0
5	BME	A	406	-	3,3,3	0.47	0	1,2,2	1.11	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	401	-	42,48,48	0.70	0	50,73,73	1.08	4 (8%)
4	EDO	A	408	-	3,3,3	0.32	0	2,2,2	0.18	0
4	EDO	A	405	-	3,3,3	0.40	0	2,2,2	0.18	0
4	EDO	A	403	-	3,3,3	0.44	0	2,2,2	0.30	0
4	EDO	A	404	-	3,3,3	0.39	0	2,2,2	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SPD	B	1602	-	-	4/7/7/7	-
4	EDO	B	1604	-	-	0/1/1/1	-
6	OXM	B	1603	-	-	0/0/4/4	-
2	NAD	B	1601	-	-	6/26/62/62	0/5/5/5
3	SPD	A	402	-	-	1/7/7/7	-
4	EDO	A	407	-	-	1/1/1/1	-
5	BME	A	406	-	-	0/1/1/1	-
2	NAD	A	401	-	-	5/26/62/62	0/5/5/5
4	EDO	A	408	-	-	0/1/1/1	-
4	EDO	A	405	-	-	1/1/1/1	-
4	EDO	A	403	-	-	0/1/1/1	-
4	EDO	A	404	-	-	0/1/1/1	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1601	NAD	C4A-N3A	-3.63	1.30	1.35
2	B	1601	NAD	PN-O2N	-3.51	1.38	1.55
2	B	1601	NAD	O7N-C7N	-3.34	1.17	1.24
2	B	1601	NAD	PA-O2A	-3.27	1.40	1.55
2	B	1601	NAD	C2N-N1N	-2.63	1.31	1.35
2	B	1601	NAD	PN-O1N	-2.57	1.41	1.50
2	B	1601	NAD	C5A-N7A	-2.43	1.30	1.39
2	B	1601	NAD	PA-O1A	-2.38	1.42	1.50
2	B	1601	NAD	O4B-C4B	-2.12	1.40	1.45
2	B	1601	NAD	O5D-C5D	-2.09	1.36	1.44
2	B	1601	NAD	C2N-C3N	-2.06	1.35	1.39
2	B	1601	NAD	O4D-C4D	-2.06	1.40	1.45

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1601	NAD	C4A-C5A-N7A	-3.87	105.36	109.40
2	B	1601	NAD	O3D-C3D-C4D	3.86	122.20	111.05
6	B	1603	OXM	C2-C1-N1	3.63	121.89	115.85
2	B	1601	NAD	N3A-C2A-N1A	-3.30	123.53	128.68
2	B	1601	NAD	O4D-C1D-C2D	-3.16	102.31	106.93
2	A	401	NAD	O4D-C1D-C2D	-3.01	102.53	106.93
2	B	1601	NAD	C2D-C3D-C4D	2.96	108.40	102.64
2	B	1601	NAD	O5D-PN-O1N	2.71	119.66	109.07
2	B	1601	NAD	O3D-C3D-C2D	2.68	120.49	111.82
2	B	1601	NAD	O4D-C4D-C5D	2.45	117.43	109.37
2	B	1601	NAD	C3D-C2D-C1D	-2.26	97.57	100.98
2	A	401	NAD	O2B-C2B-C1B	2.25	119.15	110.85
2	B	1601	NAD	O5D-C5D-C4D	2.22	116.64	108.99
2	A	401	NAD	O2A-PA-O1A	2.14	122.83	112.24
2	A	401	NAD	C5A-C6A-N6A	2.14	123.60	120.35
2	B	1601	NAD	C2A-N1A-C6A	2.09	122.33	118.75

There are no chirality outliers.

All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	NAD	PA-O3-PN-O5D
2	A	401	NAD	C5D-O5D-PN-O1N
2	A	401	NAD	C5D-O5D-PN-O2N
2	B	1601	NAD	PA-O3-PN-O5D
2	B	1601	NAD	C5D-O5D-PN-O1N
2	B	1601	NAD	C5D-O5D-PN-O2N
3	B	1602	SPD	N6-C7-C8-C9
3	B	1602	SPD	C3-C4-C5-N6
3	A	402	SPD	N6-C7-C8-C9
3	B	1602	SPD	N1-C2-C3-C4
2	B	1601	NAD	C5D-O5D-PN-O3
4	A	405	EDO	O1-C1-C2-O2
3	B	1602	SPD	C7-C8-C9-N10
2	A	401	NAD	C5D-O5D-PN-O3
2	A	401	NAD	PN-O3-PA-O1A
2	B	1601	NAD	PN-O3-PA-O1A
2	B	1601	NAD	PN-O3-PA-O2A
4	A	407	EDO	O1-C1-C2-O2


There are no ring outliers.

7 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1602	SPD	8	0
3	A	402	SPD	5	0
4	A	407	EDO	1	0
5	A	406	BME	3	0
2	A	401	NAD	2	0
4	A	408	EDO	2	0
4	A	405	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	284:GLY	C	285:VAL	N	1.14

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/369 (90%)	0.39	39 (11%) 4 4	23, 33, 74, 129	0
1	B	327/369 (88%)	0.36	31 (9%) 8 7	22, 32, 58, 122	1 (0%)
All	All	662/738 (89%)	0.38	70 (10%) 6 5	22, 33, 65, 129	1 (0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	87	LEU	8.9
1	A	77	LEU	7.6
1	A	80	ASP	7.5
1	B	364	HIS	7.3
1	B	94	LEU	7.2
1	A	78	SER	6.8
1	A	83	GLN	6.5
1	A	84	HIS	6.1
1	A	82	ASP	6.0
1	B	79	GLN	5.8
1	B	77	LEU	5.7
1	B	93	PRO	5.4
1	A	86[A]	ASP	5.3
1	B	78	SER	5.2
1	B	91	ARG	5.1
1	A	76	PRO	5.1
1	A	91	ARG	4.5
1	A	94	LEU	4.4
1	A	332[A]	VAL	4.4
1	B	80	ASP	4.4
1	A	79	GLN	4.3
1	B	76	PRO	4.3
1	A	28	SER	4.3
1	B	81	GLU	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	81	GLU	4.0
1	A	85	ALA	3.8
1	B	250	TYR	3.6
1	B	28	SER	3.4
1	A	362	PHE	3.3
1	B	198	GLY	3.3
1	A	272	PHE	3.2
1	A	93	PRO	3.1
1	B	286[A]	VAL	3.0
1	A	281	LEU	3.0
1	A	363	MET	2.9
1	B	312	PHE	2.9
1	B	340	TYR	2.8
1	A	283	GLY	2.8
1	A	89	GLN	2.8
1	A	308	THR	2.7
1	A	88	THR	2.7
1	B	308	THR	2.7
1	B	196	THR	2.6
1	A	235	ALA	2.5
1	A	234	PRO	2.5
1	B	235	ALA	2.5
1	A	282	GLY	2.4
1	B	346[A]	VAL	2.4
1	B	104	THR	2.4
1	A	340	TYR	2.4
1	B	341	ALA	2.4
1	B	75[A]	GLU	2.3
1	A	341	ALA	2.3
1	A	333	ASP	2.3
1	A	285	VAL	2.3
1	A	306	ILE	2.2
1	B	309	ALA	2.2
1	A	29	THR	2.2
1	A	286[A]	VAL	2.1
1	B	95	THR	2.1
1	B	32	ARG	2.1
1	A	312	PHE	2.1
1	B	186	ILE	2.1
1	A	129	VAL	2.1
1	A	309	ALA	2.0
1	B	159	ARG	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	174	GLU	2.0
1	A	95	THR	2.0
1	B	285	VAL	2.0
1	B	131	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSS	B	177	7/8	0.95	0.05	40,50,68,82	0
1	CSS	A	177	7/8	0.97	0.05	28,36,51,61	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	405	4/4	0.60	0.23	63,76,87,87	0
4	EDO	A	404	4/4	0.78	0.20	74,89,100,102	0
5	BME	A	406	4/4	0.79	0.53	180,217,219,219	0
4	EDO	A	407	4/4	0.83	0.22	114,136,137,137	0
3	SPD	A	402	10/10	0.86	0.14	36,47,55,57	0
6	OXM	B	1603	6/6	0.90	0.09	47,53,67,67	0
2	NAD	B	1601	44/44	0.93	0.15	22,32,42,45	0
3	SPD	B	1602	10/10	0.94	0.16	29,39,52,53	0
4	EDO	A	408	4/4	0.94	0.11	32,44,51,53	10
4	EDO	A	403	4/4	0.95	0.16	32,38,40,44	0
4	EDO	B	1604	4/4	0.96	0.21	22,27,29,33	10
2	NAD	A	401	44/44	0.97	0.18	21,30,38,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.