

Full wwPDB X-ray Structure Validation Report (i)

Jun 24, 2024 – 08:45 AM EDT

PDB ID : 7D79

Title : The structure of DcsB complex with its substrate analogue

Authors: Tang, Y.; Zhou, J.H.; Wang, G.Q.

Deposited on : 2020-10-03

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.20.1

EDS : 2.37.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	322	86%	5%	9%
1	В	322	85%	5%	10%
1	С	322	84%	6%	10%
1	D	322	83%	9%	8%



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 9646 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DltD domain-containing protein.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	D	295	Total	С	N	О	S	0	9	0
1	ש	290	2290	1462	385	429	14	0		U
1	Λ	293	Total	С	N	О	S	0	0	0
1	A	290	2246	1434	375	423	14		U	
1	В	290	Total	С	N	О	S	0	1	0
1	Б	290	2241	1433	373	421	14	0	1	0
1	С	200	Total	С	N	О	S	0	0	0
1		C 290	2228	1424	372	418	14			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	MET	-	initiating methionine	UNP J4WAT9
D	-18	GLY	-	expression tag	UNP J4WAT9
D	-17	SER	-	expression tag	UNP J4WAT9
D	-16	SER	-	expression tag	UNP J4WAT9
D	-15	HIS	-	expression tag	UNP J4WAT9
D	-14	HIS	-	expression tag	UNP J4WAT9
D	-13	HIS	-	expression tag	UNP J4WAT9
D	-12	HIS	-	expression tag	UNP J4WAT9
D	-11	HIS	-	expression tag	UNP J4WAT9
D	-10	HIS	-	expression tag	UNP J4WAT9
D	-9	SER	-	expression tag	UNP J4WAT9
D	-8	SER	-	expression tag	UNP J4WAT9
D	-7	GLY	-	expression tag	UNP J4WAT9
D	-6	LEU	-	expression tag	UNP J4WAT9
D	-5	VAL	-	expression tag	UNP J4WAT9
D	-4	PRO	-	expression tag	UNP J4WAT9
D	-3	ARG	-	expression tag	UNP J4WAT9
D	-2	GLY	=	expression tag	UNP J4WAT9
D	-1	SER	=	expression tag	UNP J4WAT9
D	0	HIS	-	expression tag	UNP J4WAT9
A	-19	MET	-	initiating methionine	UNP J4WAT9



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	expression tag	UNP J4WAT9
A	-17	SER	-	expression tag	UNP J4WAT9
A	-16	SER	-	expression tag	UNP J4WAT9
A	-15	HIS	_	expression tag	UNP J4WAT9
A	-14	HIS	_	expression tag	UNP J4WAT9
A	-13	HIS	-	expression tag	UNP J4WAT9
A	-12	HIS	-	expression tag	UNP J4WAT9
A	-11	HIS	-	expression tag	UNP J4WAT9
A	-10	HIS	-	expression tag	UNP J4WAT9
A	-9	SER	-	expression tag	UNP J4WAT9
A	-8	SER	-	expression tag	UNP J4WAT9
A	-7	GLY	-	expression tag	UNP J4WAT9
A	-6	LEU	-	expression tag	UNP J4WAT9
A	-5	VAL	-	expression tag	UNP J4WAT9
A	-4	PRO	-	expression tag	UNP J4WAT9
A	-3	ARG	-	expression tag	UNP J4WAT9
A	-2	GLY	_	expression tag	UNP J4WAT9
A	-1	SER	-	expression tag	UNP J4WAT9
A	0	HIS	-	expression tag	UNP J4WAT9
В	-19	MET	_	initiating methionine	UNP J4WAT9
В	-18	GLY	-	expression tag	UNP J4WAT9
В	-17	SER	-	expression tag	UNP J4WAT9
В	-16	SER	-	expression tag	UNP J4WAT9
В	-15	HIS	-	expression tag	UNP J4WAT9
В	-14	HIS	-	expression tag	UNP J4WAT9
В	-13	HIS	-	expression tag	UNP J4WAT9
В	-12	HIS	-	expression tag	UNP J4WAT9
В	-11	HIS	-	expression tag	UNP J4WAT9
В	-10	HIS	_	expression tag	UNP J4WAT9
В	-9	SER	_	expression tag	UNP J4WAT9
В	-8	SER	_	expression tag	UNP J4WAT9
В	-7	GLY	_	expression tag	UNP J4WAT9
В	-6	LEU	-	expression tag	UNP J4WAT9
В	-5	VAL	-	expression tag	UNP J4WAT9
В	-4	PRO	-	expression tag	UNP J4WAT9
В	-3	ARG	-	expression tag	UNP J4WAT9
В	-2	GLY	-	expression tag	UNP J4WAT9
В	-1	SER	-	expression tag	UNP J4WAT9
В	0	HIS	-	expression tag	UNP J4WAT9
С	-19	MET	-	initiating methionine	UNP J4WAT9
С	-18	GLY	-	expression tag	UNP J4WAT9
С	-17	SER	-	expression tag	UNP J4WAT9



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
С	-16	SER	-	expression tag	UNP J4WAT9
С	-15	HIS	-	expression tag	UNP J4WAT9
С	-14	HIS	-	expression tag	UNP J4WAT9
С	-13	HIS	-	expression tag	UNP J4WAT9
С	-12	HIS	-	expression tag	UNP J4WAT9
С	-11	HIS	1	expression tag	UNP J4WAT9
С	-10	HIS	-	expression tag	UNP J4WAT9
С	-9	SER	-	expression tag	UNP J4WAT9
С	-8	SER	-	expression tag	UNP J4WAT9
С	-7	GLY	-	expression tag	UNP J4WAT9
С	-6	LEU	-	expression tag	UNP J4WAT9
С	-5	VAL	-	expression tag	UNP J4WAT9
С	-4	PRO	-	expression tag	UNP J4WAT9
С	-3	ARG	-	expression tag	UNP J4WAT9
С	-2	GLY	-	expression tag	UNP J4WAT9
С	-1	SER	-	expression tag	UNP J4WAT9
С	0	HIS	-	expression tag	UNP J4WAT9

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Cl 1 1	0	0
2	A	2	Total Cl 2 2	0	0
2	С	1	Total Cl 1 1	0	0

• Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0


• Molecule 4 is methyl 3-[(E,3R,9R)-3,9-bis(oxidanyl)dec-4-enoyl]sulfanylpropanoate (three-letter code: GY0) (formula: $C_{14}H_{24}O_5S$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total 20	C 14	O 5	S 1	0	0


• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	С	1	Total C O 6 3 3	0	0



• Molecule 6 is methyl 3-sulfanylpropanoate (three-letter code: GYF) (formula: $C_4H_8O_2S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	В	1	Total 7	C 4	O 2	S 1	0	0

• Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	С	1	Total Na 1 1	0	0

• Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	185	Total O 185 185	0	0
8	A	171	Total O 171 171	0	0
8	В	122	Total O 122 122	0	0
8	С	95	Total O 95 95	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DltD domain-containing protein









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	47.26Å 59.01Å 113.99Å	Donositor
a, b, c, α , β , γ	84.85° 78.55° 84.01°	Depositor
Resolution (Å)	39.00 - 2.10	Depositor
Resolution (A)	46.14 - 2.10	EDS
% Data completeness	86.0 (39.00-2.10)	Depositor
(in resolution range)	86.0 (46.14-2.10)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.23 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
D D.	0.182 , 0.227	Depositor
R, R_{free}	0.183 , 0.228	DCC
R_{free} test set	2980 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 43.2	EDS
L-test for twinning ²	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9646	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GY0, NA, EDO, CL, GYF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.25	0/2294	0.42	0/3102	
1	В	0.25	0/2289	0.41	0/3096	
1	С	0.25	0/2275	0.41	0/3075	
1	D	0.25	0/2339	0.42	0/3164	
All	All	0.25	0/9197	0.41	0/12437	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2246	0	2251	11	0
1	В	2241	0	2250	9	0
1	С	2228	0	2236	11	0
1	D	2290	0	2303	18	0
2	A	2	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	1	0
3	A	8	0	12	0	0
3	В	4	0	6	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	12	0	18	0	0
4	D	20	0	0	0	0
5	A	6	0	8	0	0
5	С	6	0	8	0	0
6	В	7	0	0	0	0
7	С	1	0	0	0	0
8	A	171	0	0	1	0
8	В	122	0	0	0	0
8	С	95	0	0	1	0
8	D	185	0	0	0	0
All	All	9646	0	9092	46	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:C:76:SER:OG	1:C:78:LYS:NZ	2.27	0.68
1:A:175:ASN:HD21	1:A:179:GLU:HB2	1.65	0.60
1:D:11:GLU:HG2	1:D:21:ARG:HG2	1.85	0.58
1:D:46:MET:HE3	1:D:198:PHE:HE1	1.70	0.57
1:B:107:LYS:HD2	1:B:300:VAL:HG13	1.87	0.56
1:A:175:ASN:ND2	1:A:179:GLU:HB2	2.22	0.55
1:D:52:ALA:HB2	1:D:64:ILE:HD11	1.90	0.54
1:C:77:PRO:HG2	1:C:80:LEU:HD12	1.90	0.53
1:B:52:ALA:HB2	1:B:64:ILE:HD11	1.92	0.52
1:B:279:VAL:HG13	1:B:280:LEU:HG	1.91	0.52
1:D:45:GLU:HB2	1:D:207:ALA:HB2	1.91	0.51
1:B:177:LYS:NZ	1:B:179:GLU:OE1	2.35	0.51
1:C:282:GLY:O	1:C:284:GLY:N	2.44	0.51
1:C:164:ARG:NH1	8:C:502:HOH:O	2.42	0.50
1:C:52:ALA:HB2	1:C:64:ILE:HD11	1.93	0.50
1:A:175:ASN:OD1	1:A:178:GLY:N	2.45	0.49
1:A:52:ALA:HB2	1:A:64:ILE:HD11	1.92	0.49
1:D:125:ALA:HB2	1:D:234:VAL:HG22	1.96	0.48
1:D:18:ILE:HG23	1:D:74:ASP:HB2	1.95	0.48
1:B:58:GLN:O	1:B:297:ARG:NH2	2.45	0.47
1:B:260:GLU:HB2	1:C:78:LYS:CE	2.46	0.46
1:D:139[B]:ILE:HG12	1:D:250:SER:HB2	1.97	0.46
1:B:296:ILE:O	1:B:300:VAL:N	2.47	0.45



Continued from previous page...

A	A. 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:B:114:SER:HA	1:B:137:CYS:O	2.16	0.45
1:C:32:PRO:HB3	1:C:107:LYS:HB3	1.99	0.45
1:D:296:ILE:O	1:D:300:VAL:N	2.51	0.44
1:D:114:SER:HA	1:D:137:CYS:O	2.17	0.43
1:D:173:PRO:HG3	1:D:214:ILE:HG23	2.00	0.43
1:D:274:LYS:NZ	1:D:281:SER:O	2.45	0.43
1:D:226:GLN:HG3	1:A:226:GLN:HB2	2.00	0.43
1:A:114:SER:HA	1:A:137:CYS:O	2.19	0.42
1:A:279:VAL:HG13	1:A:280:LEU:HG	2.01	0.42
1:C:69:SER:HB2	1:C:78:LYS:HA	2.01	0.42
1:D:226:GLN:HB2	1:A:226:GLN:HG3	2.01	0.42
1:D:13:LYS:HE2	1:D:13:LYS:HB3	1.92	0.42
1:D:33:ALA:HB3	1:D:108:ILE:HD13	2.01	0.42
1:A:183:GLY:HA2	1:A:187:SER:HB3	2.01	0.42
1:C:240:LEU:HD13	1:C:267:LYS:HD3	2.01	0.42
1:C:274:LYS:NZ	1:C:281:SER:O	2.48	0.42
1:D:14:THR:OG1	1:D:16:ASP:OD1	2.29	0.41
1:A:56:GLN:NE2	8:A:502:HOH:O	2.27	0.41
1:C:114:SER:HA	1:C:137:CYS:O	2.20	0.41
1:A:259:PHE:CE2	1:A:268:PHE:HB2	2.56	0.41
1:D:211:ARG:HD3	2:D:401:CL:CL	2.59	0.40
1:D:46:MET:HE3	1:D:198:PHE:CE1	2.54	0.40
1:B:235:ASP:OD1	1:B:236:LYS:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	A	289/322~(90%)	279 (96%)	10 (4%)	0	100	100
1	В	287/322 (89%)	277 (96%)	10 (4%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	C	284/322 (88%)	272 (96%)	12 (4%)	0	100	100
1	D	295/322~(92%)	284 (96%)	11 (4%)	0	100	100
All	All	1155/1288 (90%)	1112 (96%)	43 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Percentiles		
1	A	243/268 (91%)	243 (100%)	0	100	100
1	В	$244/268 \ (91\%)$	242 (99%)	2 (1%)	81	86
1	С	242/268 (90%)	240 (99%)	2 (1%)	81	86
1	D	$249/268 \; (93\%)$	249 (100%)	0	100	100
All	All	978/1072 (91%)	974 (100%)	4 (0%)	91	94

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	8	GLN
1	В	300	VAL
1	С	171	LEU
1	С	187	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 15 ligands modelled in this entry, 5 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	403	-	5,5,5	0.87	0	5,5,5	1.10	0
5	GOL	С	402	-	5,5,5	0.90	0	5,5,5	1.08	0
3	EDO	В	401	_	3,3,3	0.43	0	2,2,2	0.34	0
4	GY0	D	405	-	19,19,19	1.23	3 (15%)	21,22,22	1.92	6 (28%)
3	EDO	D	403	-	3,3,3	0.42	0	2,2,2	0.38	0
3	EDO	A	404	_	3,3,3	0.41	0	2,2,2	0.37	0
6	GYF	В	402	-	5,6,6	1.60	2 (40%)	5,6,6	1.29	1 (20%)
3	EDO	D	402	-	3,3,3	0.43	0	2,2,2	0.36	0
3	EDO	A	405	-	3,3,3	0.41	0	2,2,2	0.31	0
3	EDO	D	404	-	3,3,3	0.43	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	403	-	-	0/4/4/4	-
5	GOL	С	402	-	-	0/4/4/4	-
3	EDO	В	401	-	-	1/1/1/1	-
4	GY0	D	405	-	-	8/20/20/20	-



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	403	-	-	0/1/1/1	-
3	EDO	A	404	-	-	0/1/1/1	-
6	GYF	В	402	-	-	2/5/5/5	-
3	EDO	D	402	-	-	0/1/1/1	-
3	EDO	A	405	-	-	1/1/1/1	-
3	EDO	D	404	-	-	0/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
4	D	405	GY0	C10-S11	2.43	1.82	1.76
4	D	405	GY0	O18-C17	2.40	1.40	1.33
6	В	402	GYF	O01-C02	2.39	1.40	1.33
6	В	402	GYF	O01-C07	-2.19	1.40	1.45
4	D	405	GY0	O18-C19	-2.12	1.40	1.45

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
4	D	405	GY0	C09-C10-S11	5.81	120.89	113.56
4	D	405	GY0	O16-C10-S11	-3.10	118.74	122.68
4	D	405	GY0	C03-C04-C05	-2.25	109.59	114.47
4	D	405	GY0	O16-C10-C09	-2.15	120.36	123.74
6	В	402	GYF	O01-C02-C03	2.14	120.22	112.14
4	D	405	GY0	C12-S11-C10	2.11	108.09	101.84
4	D	405	GY0	O18-C17-C13	2.10	120.09	112.14

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	405	GY0	C14-C02-C03-C04
4	D	405	GY0	C08-C09-C10-S11
6	В	402	GYF	C03-C02-O01-C07
4	D	405	GY0	O15-C08-C09-C10
6	В	402	GYF	O06-C02-O01-C07
4	D	405	GY0	O01-C02-C03-C04
4	D	405	GY0	C08-C09-C10-O16
3	A	405	EDO	O1-C1-C2-O2
4	D	405	GY0	C04-C05-C06-C07
4	D	405	GY0	C13-C12-S11-C10




Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	В	401	EDO	O1-C1-C2-O2
4	D	405	GY0	C07-C08-C09-C10


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	293/322 (90%)	-0.21	0 100 100	15, 23, 39, 57	0
1	В	290/322~(90%)	-0.18	2 (0%) 87 89	17, 28, 41, 64	0
1	С	290/322~(90%)	-0.04	4 (1%) 75 78	19, 32, 49, 60	0
1	D	295/322~(91%)	-0.25	1 (0%) 94 94	15, 23, 36, 51	0
All	All	1168/1288 (90%)	-0.17	7 (0%) 89 91	15, 26, 44, 64	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	208	PRO	3.1
1	С	246	LEU	2.5
1	С	269	LEU	2.3
1	В	196	TYR	2.2
1	С	286	VAL	2.2
1	С	268	PHE	2.2
1	D	207	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	EDO	D	402	4/4	0.87	0.16	27,32,36,43	0
3	EDO	A	405	4/4	0.90	0.14	30,32,32,41	0
5	GOL	A	403	6/6	0.91	0.20	25,26,32,37	0
5	GOL	С	402	6/6	0.91	0.21	32,39,44,52	0
3	EDO	В	401	4/4	0.92	0.32	36,36,37,45	0
2	CL	A	402	1/1	0.93	0.19	52,52,52,52	0
3	EDO	D	403	4/4	0.93	0.15	27,29,35,37	0
4	GY0	D	405	20/20	0.93	0.14	19,27,37,37	0
3	EDO	D	404	4/4	0.93	0.17	29,31,35,44	0
3	EDO	A	404	4/4	0.93	0.20	29,42,45,45	0
6	GYF	В	402	7/7	0.96	0.08	25,34,40,53	0
7	NA	С	403	1/1	0.96	0.15	35,35,35,35	0
2	CL	D	401	1/1	0.97	0.21	46,46,46,46	0
2	CL	С	401	1/1	0.98	0.17	42,42,42,42	0
2	CL	A	401	1/1	0.98	0.09	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









Electron density around EDO A 405: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)













Electron density around CL A 402:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO D 403: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around EDO D 404: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




















Electron density around CL D 401:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














6.5 Other polymers (i)

There are no such residues in this entry.

