



Full wwPDB X-ray Structure Validation Report i

Jun 24, 2024 – 10:31 AM EDT

PDB ID : 7NCT
Title : Glutathione-S-transferase GliG mutant K127G
Authors : Groll, M.; Huber, E.M.
Deposited on : 2021-01-29
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain		
1	F	253	6%	89%	5% 7%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 11545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione S-transferase GliG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total 1918	C 1232	N 330	O 351	S 5	0	0	0
1	B	237	Total 1918	C 1232	N 330	O 351	S 5	0	0	0
1	C	236	Total 1909	C 1227	N 329	O 348	S 5	0	0	0
1	D	238	Total 1924	C 1235	N 331	O 353	S 5	0	0	0
1	E	237	Total 1918	C 1232	N 330	O 351	S 5	0	0	0
1	F	236	Total 1909	C 1227	N 329	O 348	S 5	0	0	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	initiating methionine	UNP B0Y813
A	-11	SER	-	expression tag	UNP B0Y813
A	-10	GLY	-	expression tag	UNP B0Y813
A	-9	SER	-	expression tag	UNP B0Y813
A	-8	HIS	-	expression tag	UNP B0Y813
A	-7	HIS	-	expression tag	UNP B0Y813
A	-6	HIS	-	expression tag	UNP B0Y813
A	-5	HIS	-	expression tag	UNP B0Y813
A	-4	HIS	-	expression tag	UNP B0Y813
A	-3	HIS	-	expression tag	UNP B0Y813
A	-2	SER	-	expression tag	UNP B0Y813
A	-1	GLY	-	expression tag	UNP B0Y813
A	0	SER	-	expression tag	UNP B0Y813
A	127	GLY	LYS	engineered mutation	UNP B0Y813
B	-12	MET	-	initiating methionine	UNP B0Y813
B	-11	SER	-	expression tag	UNP B0Y813
B	-10	GLY	-	expression tag	UNP B0Y813

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	SER	-	expression tag	UNP B0Y813
B	-8	HIS	-	expression tag	UNP B0Y813
B	-7	HIS	-	expression tag	UNP B0Y813
B	-6	HIS	-	expression tag	UNP B0Y813
B	-5	HIS	-	expression tag	UNP B0Y813
B	-4	HIS	-	expression tag	UNP B0Y813
B	-3	HIS	-	expression tag	UNP B0Y813
B	-2	SER	-	expression tag	UNP B0Y813
B	-1	GLY	-	expression tag	UNP B0Y813
B	0	SER	-	expression tag	UNP B0Y813
B	127	GLY	LYS	engineered mutation	UNP B0Y813
C	-12	MET	-	initiating methionine	UNP B0Y813
C	-11	SER	-	expression tag	UNP B0Y813
C	-10	GLY	-	expression tag	UNP B0Y813
C	-9	SER	-	expression tag	UNP B0Y813
C	-8	HIS	-	expression tag	UNP B0Y813
C	-7	HIS	-	expression tag	UNP B0Y813
C	-6	HIS	-	expression tag	UNP B0Y813
C	-5	HIS	-	expression tag	UNP B0Y813
C	-4	HIS	-	expression tag	UNP B0Y813
C	-3	HIS	-	expression tag	UNP B0Y813
C	-2	SER	-	expression tag	UNP B0Y813
C	-1	GLY	-	expression tag	UNP B0Y813
C	0	SER	-	expression tag	UNP B0Y813
C	127	GLY	LYS	engineered mutation	UNP B0Y813
D	-12	MET	-	initiating methionine	UNP B0Y813
D	-11	SER	-	expression tag	UNP B0Y813
D	-10	GLY	-	expression tag	UNP B0Y813
D	-9	SER	-	expression tag	UNP B0Y813
D	-8	HIS	-	expression tag	UNP B0Y813
D	-7	HIS	-	expression tag	UNP B0Y813
D	-6	HIS	-	expression tag	UNP B0Y813
D	-5	HIS	-	expression tag	UNP B0Y813
D	-4	HIS	-	expression tag	UNP B0Y813
D	-3	HIS	-	expression tag	UNP B0Y813
D	-2	SER	-	expression tag	UNP B0Y813
D	-1	GLY	-	expression tag	UNP B0Y813
D	0	SER	-	expression tag	UNP B0Y813
D	127	GLY	LYS	engineered mutation	UNP B0Y813
E	-12	MET	-	initiating methionine	UNP B0Y813
E	-11	SER	-	expression tag	UNP B0Y813
E	-10	GLY	-	expression tag	UNP B0Y813

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-9	SER	-	expression tag	UNP B0Y813
E	-8	HIS	-	expression tag	UNP B0Y813
E	-7	HIS	-	expression tag	UNP B0Y813
E	-6	HIS	-	expression tag	UNP B0Y813
E	-5	HIS	-	expression tag	UNP B0Y813
E	-4	HIS	-	expression tag	UNP B0Y813
E	-3	HIS	-	expression tag	UNP B0Y813
E	-2	SER	-	expression tag	UNP B0Y813
E	-1	GLY	-	expression tag	UNP B0Y813
E	0	SER	-	expression tag	UNP B0Y813
E	127	GLY	LYS	engineered mutation	UNP B0Y813
F	-12	MET	-	initiating methionine	UNP B0Y813
F	-11	SER	-	expression tag	UNP B0Y813
F	-10	GLY	-	expression tag	UNP B0Y813
F	-9	SER	-	expression tag	UNP B0Y813
F	-8	HIS	-	expression tag	UNP B0Y813
F	-7	HIS	-	expression tag	UNP B0Y813
F	-6	HIS	-	expression tag	UNP B0Y813
F	-5	HIS	-	expression tag	UNP B0Y813
F	-4	HIS	-	expression tag	UNP B0Y813
F	-3	HIS	-	expression tag	UNP B0Y813
F	-2	SER	-	expression tag	UNP B0Y813
F	-1	GLY	-	expression tag	UNP B0Y813
F	0	SER	-	expression tag	UNP B0Y813
F	127	GLY	LYS	engineered mutation	UNP B0Y813


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	16	Total O 16 16	0	0
2	B	3	Total O 3 3	0	0
2	C	8	Total O 8 8	0	0
2	D	8	Total O 8 8	0	0
2	E	11	Total O 11 11	0	0
2	F	3	Total O 3 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutathione S-transferase GliG




- Molecule 1: Glutathione S-transferase GliG




- Molecule 1: Glutathione S-transferase GliG




- Molecule 1: Glutathione S-transferase GliG





- Molecule 1: Glutathione S-transferase GliG





- Molecule 1: Glutathione S-transferase GliG



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	139.43Å 81.23Å 126.98Å 90.00° 90.88° 90.00°	Depositor
Resolution (Å)	30.00 – 2.65 47.30 – 2.65	Depositor EDS
% Data completeness (in resolution range)	94.4 (30.00-2.65) 92.2 (47.30-2.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.59 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R , R_{free}	0.214 , 0.281 0.218 , 0.281	Depositor DCC
R_{free} test set	1959 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	76.3	Xtriage
Anisotropy	0.476	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.014 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.018 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.248 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.237 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.030 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11545	wwPDB-VP
Average B, all atoms (Å ²)	105.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.65	0/1968	0.72	0/2673
1	B	0.64	0/1968	0.69	0/2673
1	C	0.65	0/1959	0.70	0/2661
1	D	0.65	0/1974	0.69	0/2681
1	E	0.65	0/1968	0.69	0/2673
1	F	0.65	0/1959	0.70	0/2661
All	All	0.65	0/11796	0.70	0/16022

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1918	0	1911	15	0
1	B	1918	0	1911	10	0
1	C	1909	0	1905	18	0
1	D	1924	0	1916	6	0
1	E	1918	0	1911	8	0
1	F	1909	0	1905	7	0
2	A	16	0	0	0	0
2	B	3	0	0	0	0
2	C	8	0	0	0	0
2	D	8	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	11	0	0	0	0
2	F	3	0	0	0	0
All	All	11545	0	11459	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:ALA:HB2	1:C:232:GLU:HA	1.69	0.74
1:A:22:ALA:CB	1:C:232:GLU:HA	2.19	0.73
1:C:32:LEU:HD22	1:C:225:VAL:HG23	1.76	0.67
1:A:96:LYS:HD2	1:B:201:LYS:HB3	1.80	0.63
1:E:141:PRO:HG2	1:F:22:ALA:HB2	1.79	0.63
1:C:25:THR:HB	1:C:225:VAL:CG1	2.28	0.63
1:B:47:VAL:HG21	1:B:55:PHE:CE1	2.34	0.62
1:B:184:LEU:O	1:B:188:MET:HG2	2.00	0.61
1:C:25:THR:HB	1:C:225:VAL:HG11	1.82	0.61
1:D:25:THR:HG22	1:D:225:VAL:HG22	1.83	0.60
1:E:46:VAL:HG23	1:E:238:LEU:HD11	1.87	0.57
1:A:4:ARG:HG2	1:A:5:PRO:HD2	1.87	0.56
1:A:78:LEU:HD22	1:D:106:GLN:HA	1.91	0.53
1:A:230:HIS:CE1	1:C:230:HIS:CE1	2.97	0.53
1:C:32:LEU:CD2	1:C:225:VAL:HG23	2.38	0.53
1:E:161:ARG:O	1:E:164:GLU:HG3	2.09	0.52
1:A:71:SER:HB2	1:A:72:PRO:HD2	1.91	0.52
1:C:19:LYS:HG2	1:C:230:HIS:CE1	2.46	0.51
1:C:148:ARG:HH11	1:C:148:ARG:HG3	1.75	0.50
1:E:83:SER:O	1:E:86:THR:OG1	2.26	0.49
1:A:96:LYS:HA	1:B:201:LYS:HD3	1.95	0.48
1:C:140:LEU:HD12	1:C:140:LEU:N	2.30	0.47
1:B:58:ILE:O	1:B:79:ARG:HD2	2.14	0.47
1:F:12:ARG:HD3	1:F:41:PRO:HB2	1.96	0.47
1:C:70:ARG:HG2	1:C:74:GLY:HA2	1.95	0.46
1:F:184:LEU:HB3	1:F:185:PRO:HD3	1.97	0.46
1:D:25:THR:O	1:D:29:VAL:HB	2.17	0.45
1:A:71:SER:HB2	1:A:72:PRO:CD	2.47	0.45
1:C:44:ILE:HG12	1:C:228:PHE:CZ	2.52	0.45
1:D:58:ILE:O	1:D:79:ARG:HD2	2.16	0.45
1:C:148:ARG:HD3	1:C:148:ARG:C	2.37	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:184:LEU:HB3	1:E:185:PRO:HD3	1.99	0.45
1:A:13:LEU:C	1:A:13:LEU:HD23	2.38	0.44
1:E:58:ILE:O	1:E:79:ARG:HD2	2.16	0.44
1:A:78:LEU:CD2	1:D:106:GLN:HA	2.48	0.43
1:A:223:GLN:OE1	1:C:223:GLN:NE2	2.50	0.43
1:F:238:LEU:C	1:F:238:LEU:HD13	2.39	0.43
1:A:22:ALA:HB1	1:C:232:GLU:HA	1.97	0.42
1:C:131:TYR:HD1	1:C:135:LEU:HD23	1.83	0.42
1:B:214:GLU:HA	1:B:219:LYS:HE2	2.02	0.41
1:B:184:LEU:HB3	1:B:185:PRO:HD3	2.02	0.41
1:B:23:THR:HG22	1:B:130:LEU:HD11	2.02	0.41
1:B:67:ILE:O	1:B:79:ARG:HA	2.21	0.41
1:E:14:VAL:O	1:E:67:ILE:HA	2.19	0.41
1:C:78:LEU:HD23	1:C:78:LEU:N	2.35	0.41
1:F:67:ILE:O	1:F:79:ARG:HA	2.20	0.41
1:B:113:ASN:OD1	1:F:80:ALA:HA	2.21	0.41
1:F:148:ARG:HD2	1:F:197:LEU:HD21	2.03	0.41
1:A:118:HIS:CD2	1:A:183:THR:HA	2.57	0.40
1:E:5:PRO:HB2	1:E:45:TYR:OH	2.21	0.40
1:D:162:LEU:CD1	1:D:205:LEU:HA	2.51	0.40
1:A:68:LEU:HA	1:A:68:LEU:HD23	1.76	0.40
1:C:78:LEU:HD23	1:C:78:LEU:H	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	235/253 (93%)	228 (97%)	6 (3%)	1 (0%)	34 48
1	B	235/253 (93%)	227 (97%)	7 (3%)	1 (0%)	34 48
1	C	234/253 (92%)	229 (98%)	4 (2%)	1 (0%)	34 48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	D	236/253 (93%)	226 (96%)	9 (4%)	1 (0%)	34 48
1	E	235/253 (93%)	226 (96%)	8 (3%)	1 (0%)	34 48
1	F	234/253 (92%)	227 (97%)	6 (3%)	1 (0%)	34 48
All	All	1409/1518 (93%)	1363 (97%)	40 (3%)	6 (0%)	34 48

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	GLU
1	C	82	GLU
1	D	82	GLU
1	E	82	GLU
1	B	82	GLU
1	F	82	GLU

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	205/218 (94%)	201 (98%)	4 (2%)	55 73
1	B	205/218 (94%)	201 (98%)	4 (2%)	55 73
1	C	204/218 (94%)	203 (100%)	1 (0%)	88 94
1	D	206/218 (94%)	205 (100%)	1 (0%)	88 94
1	E	205/218 (94%)	204 (100%)	1 (0%)	88 94
1	F	204/218 (94%)	204 (100%)	0	100 100
All	All	1229/1308 (94%)	1218 (99%)	11 (1%)	78 87

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	VAL
1	A	70	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	73	ASP
1	A	223	GLN
1	B	21	THR
1	B	188	MET
1	B	190	SER
1	B	223	GLN
1	C	148	ARG
1	D	106	GLN
1	E	83	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	GLN
1	C	223	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/253 (93%)	-0.12	3 (1%) 77 75	52, 88, 138, 182	0
1	B	237/253 (93%)	0.06	5 (2%) 63 59	58, 93, 145, 168	0
1	C	236/253 (93%)	-0.04	3 (1%) 77 75	58, 92, 131, 160	0
1	D	238/253 (94%)	0.31	16 (6%) 17 14	54, 112, 162, 193	0
1	E	237/253 (93%)	0.33	20 (8%) 11 8	62, 113, 157, 191	0
1	F	236/253 (93%)	0.41	16 (6%) 17 14	58, 119, 169, 200	0
All	All	1421/1518 (93%)	0.16	63 (4%) 34 31	52, 102, 157, 200	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	141	PRO	6.1
1	E	232	GLU	5.8
1	A	137	PRO	5.3
1	D	133	TYR	5.0
1	B	3	GLU	4.9
1	D	137	PRO	4.7
1	D	232	GLU	4.6
1	E	228	PHE	4.3
1	D	198	GLU	4.0
1	E	143	THR	4.0
1	E	100	PHE	3.9
1	E	3	GLU	3.8
1	D	193	GLU	3.8
1	E	158	LEU	3.7
1	E	229	GLY	3.6
1	B	232	GLU	3.6
1	F	234	GLU	3.6
1	E	147	LEU	3.5
1	D	141	PRO	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	75	ARG	3.2
1	A	132	PHE	3.1
1	D	32	LEU	3.1
1	D	90	ILE	3.1
1	F	133	TYR	3.0
1	B	143	THR	3.0
1	A	138	GLU	3.0
1	F	232	GLU	2.9
1	D	127	GLY	2.8
1	F	4	ARG	2.8
1	F	184	LEU	2.8
1	E	178	ILE	2.8
1	F	137	PRO	2.8
1	E	212	MET	2.8
1	D	200	GLU	2.7
1	E	16	PHE	2.7
1	C	130	LEU	2.7
1	F	140	LEU	2.7
1	D	147	LEU	2.6
1	B	137	PRO	2.6
1	F	132	PHE	2.6
1	F	229	GLY	2.6
1	E	130	LEU	2.6
1	E	67	ILE	2.5
1	F	152	THR	2.5
1	E	89	TYR	2.5
1	D	158	LEU	2.5
1	D	38	LEU	2.4
1	F	16	PHE	2.4
1	D	126	ALA	2.4
1	D	210	VAL	2.4
1	C	168	GLN	2.4
1	F	142	LYS	2.4
1	C	129	TRP	2.4
1	B	132	PHE	2.3
1	E	170	LEU	2.2
1	D	130	LEU	2.2
1	E	38	LEU	2.2
1	E	197	LEU	2.2
1	F	9	VAL	2.1
1	E	195	PHE	2.1
1	F	195	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	9	VAL	2.1
1	F	198	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.