



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2023 – 01:24 pm BST

PDB ID : 7OU1
Title : Crystal structure of Rhizobium etli inducible L-asparaginase ReAV (monoclinic form MP2)
Authors : Imiolczyk, B.; Loch, J.I.; Gilski, M.; Jaskolski, M.
Deposited on : 2021-06-10
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.32.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	373	 91% 6%
1	BBB	373	 91% 6%
1	CCC	373	 89% 5% 6%
1	DDD	373	 92% 6%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 11942 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-asparaginase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	350	Total 2619	C 1617	N 481	O 499	S 22	0	5	0
1	BBB	350	Total 2635	C 1626	N 484	O 503	S 22	0	8	0
1	CCC	350	Total 2641	C 1632	N 483	O 503	S 23	0	9	0
1	DDD	350	Total 2636	C 1627	N 485	O 502	S 22	0	8	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-5	GLY	-	expression tag	UNP Q2K0Z2
AAA	-4	ILE	-	expression tag	UNP Q2K0Z2
AAA	-3	ASP	-	expression tag	UNP Q2K0Z2
AAA	-2	PRO	-	expression tag	UNP Q2K0Z2
AAA	-1	PHE	-	expression tag	UNP Q2K0Z2
AAA	0	THR	-	expression tag	UNP Q2K0Z2
BBB	-5	GLY	-	expression tag	UNP Q2K0Z2
BBB	-4	ILE	-	expression tag	UNP Q2K0Z2
BBB	-3	ASP	-	expression tag	UNP Q2K0Z2
BBB	-2	PRO	-	expression tag	UNP Q2K0Z2
BBB	-1	PHE	-	expression tag	UNP Q2K0Z2
BBB	0	THR	-	expression tag	UNP Q2K0Z2
CCC	-5	GLY	-	expression tag	UNP Q2K0Z2
CCC	-4	ILE	-	expression tag	UNP Q2K0Z2
CCC	-3	ASP	-	expression tag	UNP Q2K0Z2
CCC	-2	PRO	-	expression tag	UNP Q2K0Z2
CCC	-1	PHE	-	expression tag	UNP Q2K0Z2
CCC	0	THR	-	expression tag	UNP Q2K0Z2
DDD	-5	GLY	-	expression tag	UNP Q2K0Z2
DDD	-4	ILE	-	expression tag	UNP Q2K0Z2
DDD	-3	ASP	-	expression tag	UNP Q2K0Z2

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
DDD	-2	PRO	-	expression tag	UNP Q2K0Z2
DDD	-1	PHE	-	expression tag	UNP Q2K0Z2
DDD	0	THR	-	expression tag	UNP Q2K0Z2

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total Zn 1 1	0	0
2	BBB	1	Total Zn 1 1	0	0
2	CCC	1	Total Zn 1 1	0	0
2	DDD	1	Total Zn 1 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	DDD	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	336	Total 336	O 336	0	0
4	BBB	364	Total 364	O 364	0	0
4	CCC	373	Total 373	O 373	0	0
4	DDD	330	Total 330	O 330	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: L-asparaginase




- Molecule 1: L-asparaginase



- Molecule 1: L-asparaginase



- Molecule 1: L-asparaginase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.92Å 91.31Å 114.16Å 90.00° 97.11° 90.00°	Depositor
Resolution (Å)	45.65 – 1.65 45.65 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.0 (45.65-1.65) 99.1 (45.65-1.65)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 1.64Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.190 , 0.222 0.202 , 0.235	Depositor DCC
R_{free} test set	1000 reflections (0.53%)	wwPDB-VP
Wilson B-factor (Å ²)	14.4	Xtrriage
Anisotropy	0.107	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11942	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.45 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.9855e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, ZN, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.84	0/2658	0.83	0/3589
1	BBB	0.87	0/2683	0.84	0/3622
1	CCC	0.86	0/2686	0.82	0/3627
1	DDD	0.83	0/2681	0.82	0/3619
All	All	0.85	0/10708	0.83	0/14457

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2619	0	2592	9	0
1	BBB	2635	0	2612	9	0
1	CCC	2641	0	2620	14	0
1	DDD	2636	0	2614	5	0
2	AAA	1	0	0	0	0
2	BBB	1	0	0	0	0
2	CCC	1	0	0	0	0
2	DDD	1	0	0	0	0
3	DDD	4	0	6	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	AAA	336	0	0	5	0
4	BBB	364	0	0	6	0
4	CCC	373	0	0	5	0
4	DDD	330	0	0	2	0
All	All	11942	0	10444	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:249[A]:CSO:OD	4:AAA:501:HOH:O	1.62	1.15
1:BBB:249[A]:CSO:OD	4:BBB:501:HOH:O	1.80	0.97
1:DDD:249[A]:CSO:OD	4:DDD:501:HOH:O	1.80	0.96
1:AAA:249[B]:CSO:OD	4:AAA:502:HOH:O	1.98	0.82
1:CCC:249[A]:CSO:OD	4:CCC:501:HOH:O	1.98	0.82
1:CCC:249[B]:CSO:OD	4:CCC:502:HOH:O	2.07	0.71
1:BBB:249[B]:CSO:OD	4:BBB:502:HOH:O	2.11	0.68
1:DDD:249[B]:CSO:OD	4:DDD:502:HOH:O	2.13	0.65
1:CCC:331:PRO:HD2	1:CCC:344[A]:VAL:HG11	1.80	0.63
1:DDD:310:LEU:HD11	1:DDD:350:LEU:HD21	1.81	0.61
1:DDD:68:GLY:HA3	3:DDD:402:EDO:H22	1.88	0.56
1:CCC:10:THR:HB	4:CCC:718:HOH:O	2.06	0.56
1:AAA:37:LEU:HD11	1:AAA:310:LEU:HD21	1.90	0.54
1:CCC:310:LEU:HD11	1:CCC:350:LEU:HD21	1.90	0.53
1:BBB:330:HIS:HE1	4:BBB:644:HOH:O	1.91	0.52
1:CCC:237:ARG:HG2	1:CCC:238:HIS:CE1	2.45	0.51
1:CCC:10:THR:HG22	1:CCC:344[B]:VAL:HG22	1.96	0.47
1:AAA:10:THR:HB	4:AAA:675:HOH:O	2.14	0.47
1:BBB:268:ALA:HA	1:BBB:303:LEU:HD22	1.97	0.46
1:BBB:223[B]:THR:HG22	4:BBB:541:HOH:O	2.16	0.45
1:CCC:331:PRO:HD2	1:CCC:344[A]:VAL:CG1	2.46	0.45
1:CCC:20:HIS:CD2	1:CCC:302:MET:HG3	2.54	0.43
1:CCC:265:GLY:HA3	1:CCC:269[B]:SER:OG	2.19	0.43
1:CCC:268:ALA:HA	1:CCC:303:LEU:HD22	2.01	0.43
1:AAA:43:MET:HE1	4:AAA:567:HOH:O	2.19	0.42
1:AAA:20:HIS:CD2	1:AAA:302:MET:HG3	2.55	0.42
1:AAA:322:ARG:HB3	4:AAA:753:HOH:O	2.20	0.42
1:DDD:94:MET:HB3	1:DDD:140:VAL:HG11	2.02	0.42
1:BBB:223[B]:THR:CG2	4:BBB:541:HOH:O	2.67	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:230:HIS:HA	1:BBB:233:ARG:NH2	2.34	0.41
1:CCC:94:MET:HB3	1:CCC:140:VAL:HG11	2.01	0.41
1:BBB:221:GLN:HA	4:BBB:702:HOH:O	2.21	0.41
1:CCC:254:ARG:NH1	4:CCC:503:HOH:O	2.26	0.41
1:AAA:138:LYS:O	1:AAA:142[B]:MET:HG2	2.21	0.41
1:AAA:268:ALA:HA	1:AAA:303:LEU:HD22	2.03	0.41
1:CCC:178:ASP:HB3	4:CCC:686:HOH:O	2.21	0.40
1:BBB:10:THR:HG22	1:BBB:344:VAL:HG22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	351/373 (94%)	343 (98%)	8 (2%)	0	100	100
1	BBB	354/373 (95%)	344 (97%)	10 (3%)	0	100	100
1	CCC	355/373 (95%)	347 (98%)	8 (2%)	0	100	100
1	DDD	354/373 (95%)	345 (98%)	9 (2%)	0	100	100
All	All	1414/1492 (95%)	1379 (98%)	35 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	267/282 (95%)	267 (100%)	0	100	100
1	BBB	270/282 (96%)	270 (100%)	0	100	100
1	CCC	271/282 (96%)	271 (100%)	0	100	100
1	DDD	270/282 (96%)	270 (100%)	0	100	100
All	All	1078/1128 (96%)	1078 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	AAA	249[B]	-	3,6,7	0.76	0	0,6,8	-	-
1	CSO	BBB	249[A]	-	3,6,7	0.57	0	0,6,8	-	-
1	CSO	AAA	249[A]	-	3,6,7	0.74	0	0,6,8	-	-
1	CSO	DDD	249[B]	-	3,6,7	0.88	0	0,6,8	-	-
1	CSO	CCC	249[B]	1	3,6,7	0.75	0	0,6,8	-	-
1	CSO	DDD	249[A]	-	3,6,7	0.86	0	0,6,8	-	-
1	CSO	BBB	249[B]	-	3,6,7	0.72	0	0,6,8	-	-
1	CSO	CCC	249[A]	1	3,6,7	0.74	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	AAA	249[B]	-	-	0/1/5/7	-
1	CSO	BBB	249[A]	-	-	0/1/5/7	-
1	CSO	AAA	249[A]	-	-	0/1/5/7	-
1	CSO	DDD	249[B]	-	-	0/1/5/7	-
1	CSO	CCC	249[B]	1	-	0/1/5/7	-
1	CSO	DDD	249[A]	-	-	0/1/5/7	-
1	CSO	BBB	249[B]	-	-	0/1/5/7	-
1	CSO	CCC	249[A]	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	AAA	249[B]	CSO	1	0
1	BBB	249[A]	CSO	1	0
1	AAA	249[A]	CSO	1	0
1	DDD	249[B]	CSO	1	0
1	CCC	249[B]	CSO	1	0
1	DDD	249[A]	CSO	1	0
1	BBB	249[B]	CSO	1	0
1	CCC	249[A]	CSO	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	DDD	402	-	3,3,3	0.21	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	DDD	402	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	DDD	402	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	349/373 (93%)	-0.16	4 (1%) 80 83	9, 16, 33, 48	0
1	BBB	349/373 (93%)	-0.25	4 (1%) 80 83	8, 13, 29, 39	0
1	CCC	349/373 (93%)	-0.27	3 (0%) 84 86	8, 13, 28, 41	0
1	DDD	349/373 (93%)	-0.08	11 (3%) 47 48	9, 16, 35, 65	0
All	All	1396/1492 (93%)	-0.19	22 (1%) 72 75	8, 15, 32, 65	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	3	PRO	3.7
1	DDD	126	PHE	3.3
1	DDD	124	GLN	3.1
1	DDD	220	GLY	3.1
1	DDD	3	PRO	3.0
1	DDD	116	MET	2.9
1	BBB	3	PRO	2.6
1	DDD	221	GLN	2.5
1	AAA	122	ILE	2.5
1	AAA	286	ASP	2.5
1	DDD	218	GLY	2.5
1	AAA	220	GLY	2.4
1	DDD	352	GLY	2.4
1	CCC	154	ASP	2.4
1	DDD	119	ARG	2.4
1	BBB	151	ALA	2.3
1	DDD	216	ASP	2.2
1	CCC	352	GLY	2.1
1	BBB	119	ARG	2.1
1	CCC	116	MET	2.1
1	BBB	116	MET	2.1
1	DDD	217	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	DDD	249[A]	7/8	0.97	0.09	12,14,17,17	4
1	CSO	DDD	249[B]	7/8	0.97	0.09	12,13,15,16	4
1	CSO	BBB	249[A]	7/8	0.98	0.08	11,11,13,13	4
1	CSO	BBB	249[B]	7/8	0.98	0.08	11,11,14,15	4
1	CSO	CCC	249[A]	7/8	0.98	0.08	12,12,15,17	7
1	CSO	CCC	249[B]	7/8	0.98	0.08	12,12,13,14	7
1	CSO	AAA	249[A]	7/8	0.98	0.09	13,14,15,17	4
1	CSO	AAA	249[B]	7/8	0.98	0.09	13,13,14,15	4

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	DDD	402	4/4	0.81	0.12	43,43,44,45	0
2	ZN	DDD	401	1/1	0.99	0.04	16,16,16,16	1
2	ZN	AAA	401	1/1	0.99	0.03	22,22,22,22	0
2	ZN	BBB	401	1/1	1.00	0.04	20,20,20,20	0
2	ZN	CCC	401	1/1	1.00	0.02	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN DDD 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN AAA 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN BBB 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.