

Full wwPDB NMR Structure Validation Report (i)

Dec 25, 2024 – 06:36 PM EST

PDB ID	:	7RC7
BMRB ID	:	30933
Title	:	Solution NMR Structure of [Ala19]Crp4
Authors	:	Conibear, A.C.; Rosengren, K.J.
Deposited on	:	2021-07-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI	:	$v_1n_11_5_13_A$ (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 77%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain	
1	А	32	84%	16%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model				
1 A:4-A:30 (27) 0.43 10				

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	3, 4, 7, 10, 12, 14, 15, 18, 20
2	1, 5, 6, 8, 13, 16, 17, 19
3	2, 9
Single-model clusters	11



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 527 atoms, of which 267 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Alpha-defensin 4.

Mol	Chain	Residues	Atoms			Trace			
1	٨	20	Total	С	Η	Ν	0	S	0
	А	32	527	159	267	57	38	6	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	19	ALA	GLY	engineered mutation	UNP P28311




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Alpha-defensin 4



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

• Molecule 1: Alpha-defensin 4

4.2.1 Score per residue for model 1

Chain A: 81% · 16%

4.2.2 Score per residue for model 2

• Molecule 1: Alpha-defensin 4

Chain A: 78% 6% 16%



4.2.3 Score per residue for model 3

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16%

4.2.4 Score per residue for model 4

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16%

4.2.5 Score per residue for model 5

• Molecule 1: Alpha-defensin 4

Chain A: 78% 6% 16%

4.2.6 Score per residue for model 6

• Molecule 1: Alpha-defensin 4

Chain A: 75% 9% 16%

4.2.7 Score per residue for model 7

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16%



4.2.8 Score per residue for model 8

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16%

4.2.9 Score per residue for model 9

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16%

4.2.10 Score per residue for model 10 (medoid)

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16%

- 4.2.11 Score per residue for model 11
- Molecule 1: Alpha-defensin 4

Chain A: 75% 9% 16%

- 4.2.12 Score per residue for model 12
- Molecule 1: Alpha-defensin 4

Chain A: 84% 16%



4.2.13 Score per residue for model 13

• Molecule 1: Alpha-defensin 4

Chain A: 81% • 16%

4.2.14 Score per residue for model 14

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16% ទងន រីខ្លី

4.2.15 Score per residue for model 15

• Molecule 1: Alpha-defensin 4


Chain A: 78% 6% 16%

4.2.16 Score per residue for model 16

 \bullet Molecule 1: Alpha-defensin 4

Chain A: 84% 16%

- 4.2.17 Score per residue for model 17
- Molecule 1: Alpha-defensin 4





4.2.18 Score per residue for model 18

• Molecule 1: Alpha-defensin 4

Chain A: 84% 16%

4.2.19 Score per residue for model 19

• Molecule 1: Alpha-defensin 4

Chain A:	72%	12%	16%
G1 L2 H10 C28 C28			

4.2.20 Score per residue for model 20

• Molecule 1: Alpha-defensin 4

Chain A:	72%	12%	16%
01 13 13 13 15 15 15 15 15 15 15 15 15 15 15 15 15			



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing, simulated annealing, simulated annealing.*

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	
CNS	structure calculation	
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	354
Number of shifts mapped to atoms	354
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	77%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	217	214	214	0±1
All	All	4340	4280	4280	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:8:LYS:HG2	1:A:25:PHE:CE2	0.69	2.23	20	1
1:A:18:ARG:HD2	1:A:29:CYS:SG	0.49	2.47	2	2
1:A:17:VAL:HG22	1:A:28:CYS:SG	0.49	2.48	19	1
1:A:20:THR:HB	1:A:22:GLY:O	0.48	2.09	17	1
1:A:7:ARG:HB2	1:A:10:HIS:O	0.47	2.10	6	2
1:A:12:LYS:HG2	1:A:15:GLU:CD	0.41	2.35	20	1
1:A:14:GLY:O	1:A:30:PRO:HA	0.40	2.16	15	1

All unique clashes are listed below, sorted by their clash magnitude.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	27/32~(84%)	$27 \pm 1 (99 \pm 2\%)$	$0\pm1~(1\pm2\%)$	0±0 (0±0%)	100 100
All	All	540/640~(84%)	533 (99%)	7 (1%)	0 (0%)	100 100

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

There are no Ramachandran outliers.

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the side chain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentile	\mathbf{s}
1	А	23/27~(85%)	23±0 (100±1%)	0±0 (0±1%)	91 98	
All	All	460/540~(85%)	459 (100%)	1 (0%)	91 98	

All 1 unique residues with a non-rotameric sidechain are listed below.

Mol	Chain	Res	Type	Models (Total)
1	А	10	HIS	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.



6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 77% for the well-defined parts and 76% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: $crp4G19A_CS.str$

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	354
Number of shifts mapped to atoms	354
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction}\pm{\rm precision},ppm$	Suggested action
$^{13}C_{\alpha}$	32	0.47 ± 0.45	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	28	0.11 ± 0.22	None needed (< 0.5 ppm)
$^{13}C'$	0		None (insufficient data)
¹⁵ N	29	0.03 ± 0.42	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 77%, i.e. 290 atoms were assigned a chemical shift out of a possible 379. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	109/136~(80%)	56/56~(100%)	27/54~(50%)	26/26~(100%)
Sidechain	166/208~(80%)	113/133~(85%)	52/58~(90%)	1/17~(6%)

Continued on next page...



	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Aromatic	15/35~(43%)	15/17~(88%)	0/17~(0%)	0/1~(0%)
Overall	290/379~(77%)	184/206~(89%)	79/129~(61%)	27/44~(61%)

Continued from previous page...

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 76%, i.e. 354 atoms were assigned a chemical shift out of a possible 467. 0 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	127/162~(78%)	66/67~(99%)	32/64~(50%)	29/31~(94%)
Sidechain	212/270 (79%)	145/173~(84%)	66/74~(89%)	1/23~(4%)
Aromatic	15/35~(43%)	15/17~(88%)	0/17~(0%)	0/1~(0%)
Overall	354/467~(76%)	226/257~(88%)	98/155~(63%)	30/55~(55%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	А	7	ARG	NE	124.70	76.53 - 92.65	24.9

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:







8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	409
Intra-residue (i-j =0)	162
Sequential (i-j =1)	105
Medium range ($ i-j >1$ and $ i-j <5$)	29
Long range $(i-j \ge 5)$	93
Inter-chain	0
Hydrogen bond restraints	20
Disulfide bond restraints	0
Total dihedral-angle restraints	40
Number of unmapped restraints	0
Number of restraints per residue	14.0
Number of long range restraints per residue ¹	3.5

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.3	0.16
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	Max ($^{\circ}$)
1.0-10.0 (Small)	1.6	2.42
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Bestroints type	Count	$\%^1$	Vio	lated	3	Consis	tently	y $Violated^4$
Restraints type	Count	701	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	162	39.6	4	2.5	1.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	132	32.3	4	3.0	1.0	0	0.0	0.0
Sidechain-Sidechain	30	7.3	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	105	25.7	2	1.9	0.5	0	0.0	0.0
Backbone-Backbone	33	8.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	67	16.4	2	3.0	0.5	0	0.0	0.0
Sidechain-Sidechain	5	1.2	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	29	7.1	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	2	0.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	16	3.9	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	11	2.7	0	0.0	0.0	0	0.0	0.0
Long range $(i-j \ge 5)$	93	22.7	1	1.1	0.2	0	0.0	0.0
Backbone-Backbone	18	4.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	29	7.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	46	11.2	1	2.2	0.2	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	20	4.9	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	409	100.0	7	1.7	1.7	0	0.0	0.0
Backbone-Backbone	53	13.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	260	63.6	6	2.3	1.5	0	0.0	0.0
Sidechain-Sidechain	96	23.5	1	1.0	0.2	0	0.0	0.0

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å

are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	5	Mean (Å)	Max (Å)	SD^6 (Å)	Median (Å)
Model ID	IR^1	SQ^2	MR^3	LR ⁴	$ IC^5 $	Total	Mean (A)	Max (A)	$SD^{*}(A)$	Median (A)
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0	0	0.0	0.0	0.0	0.0
4	1	0	0	0	0	1	0.15	0.15	0.0	0.15
5	1	0	0	0	0	1	0.13	0.13	0.0	0.13
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	1	0	0	0	0	1	0.14	0.14	0.0	0.14
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0

Continued on next page...




continuea ji	Continueu from previous page										
Model ID			nber o	f viola	ations	5	Mean (Å)	Max (Å)	SD^{6} (Å)	Median (Å)	
Model ID	IR^{1}	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	SD(A)	Median (A)	
11	0	0	0	1	0	1	0.1	0.1	0.0	0.1	
12	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
13	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
14	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
15	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
16	1	0	0	0	0	1	0.11	0.11	0.0	0.11	
17	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0	
19	0	1	0	0	0	1	0.16	0.16	0.0	0.16	
20	1	1	0	0	0	2	0.13	0.15	0.02	0.13	

Continued from previous page...

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right




9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 382(IR:158, SQ:103, MR:29, LR:92, IC:0) restraints are not violated in the ensemble.

Nu		of vio		restra	Fraction of the ensemble		
IR^1	SQ^2	MR^3	LR ⁴	IC ⁵	Total	Count^6	%
3	2	0	1	0	6	1	5.0
1	0	0	0	0	1	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations






9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,29)	1:20:A:THR:H	1:20:A:THR:HG21	2	0.14	0.01	0.14
(1,29)	1:20:A:THR:H	1:20:A:THR:HG22	2	0.14	0.01	0.14
(1,29)	1:20:A:THR:H	1:20:A:THR:HG23	2	0.14	0.01	0.14


 $^1\mathrm{Number}$ of violated models, $^2\mathrm{Standard}$ deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table : All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,167)	1:26:A:LEU:HG	1:27:A:TYR:H	19	0.16
(1,180)	1:8:A:LYS:H	1:8:A:LYS:HB3	20	0.15
(1,26)	1:8:A:LYS:H	1:8:A:LYS:HE2	4	0.15
(1,26)	1:8:A:LYS:H	1:8:A:LYS:HE3	4	0.15
(1,29)	1:20:A:THR:H	1:20:A:THR:HG21	9	0.14
(1,29)	1:20:A:THR:H	1:20:A:THR:HG22	9	0.14
(1,29)	1:20:A:THR:H	1:20:A:THR:HG23	9	0.14
(1,29)	1:20:A:THR:H	1:20:A:THR:HG21	5	0.13
(1,29)	1:20:A:THR:H	1:20:A:THR:HG22	5	0.13
(1,29)	1:20:A:THR:H	1:20:A:THR:HG23	5	0.13
(1,332)	1:8:A:LYS:HG2	1:9:A:GLY:H	20	0.11
(1,332)	1:8:A:LYS:HG3	1:9:A:GLY:H	20	0.11
(1,15)	1:6:A:CYS:H	1:6:A:CYS:HB3	16	0.11
(1,88)	1:7:A:ARG:HE	1:12:A:LYS:HG2	11	0.1
(1,88)	1:7:A:ARG:HE	1:12:A:LYS:HG3	11	0.1



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count $\%^1$		Vie				Consistently Violated ⁴		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$	
PSI	19	47.5	7	36.8	17.5	0	0.0	0.0	
PHI	21	52.5	4	19.0	10.0	0	0.0	0.0	
Total	40	100.0	11	27.5	27.5	0	0.0	0.0	

 1 percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories





10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nun	nber o	of violations	\mathbf{M}_{oop} (°)	M_{OV} (°)	SD (°)	Median (°)
Model ID	PSI	PHI	Total	Mean $(^{\circ})$	$Max (^{\circ})$	$SD(^{\circ})$	Median ()
1	0	2	2	1.35	1.58	0.23	1.35
2	0	1	1	1.87	1.87	0.0	1.87
3	1	1	2	2.13	2.33	0.2	2.13
4	0	1	1	1.93	1.93	0.0	1.93
5	1	0	1	1.11	1.11	0.0	1.11
6	2	1	3	1.53	1.87	0.35	1.67
7	0	1	1	1.25	1.25	0.0	1.25
8	1	0	1	1.31	1.31	0.0	1.31
9	1	1	2	1.6	2.09	0.49	1.6
10	1	3	4	1.32	1.46	0.13	1.34
11	0	0	0	0.0	0.0	0.0	0.0
12	1	1	2	1.76	2.42	0.65	1.76
13	1	1	2	1.02	1.02	0.0	1.02
14	0	1	1	1.33	1.33	0.0	1.33
15	0	0	0	0.0	0.0	0.0	0.0
16	2	0	2	1.35	1.35	0.0	1.35
17	1	0	1	1.17	1.17	0.0	1.17
18	1	0	1	1.76	1.76	0.0	1.76
19	2	1	3	1.73	2.36	0.49	1.65
20	0	1	1	1.87	1.87	0.0	1.87







The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints	Fractio	n of the ensemble
PSI	PHI	Total	Count^1	%
4	1	5	1	5.0
0	0	0	2	10.0
2	2	4	3	15.0
0	0	0	4	20.0
1	0	1	5	25.0
0	0	0	6	30.0
0	0	0	7	35.0
0	0	0	8	40.0
0	1	1	9	45.0
0	0	0	10	50.0
0	0	0	11	55.0

Continued on next page...



PSI


PHI

Continued from pretious page										
Nun	nber o	of violated restraints	Fractio	n of the ensemble						
PSI	PHI	Total	Count^1	%						
0	0	0	12	60.0						
0	0	0	13	65.0						
0	0	0	14	70.0						
0	0	0	15	75.0						
0	0	0	16	80.0						
0	0	0	17	85.0						
0	0	0	18	90.0						
0	0	0	19	95.0						
0	0	0	20	100.0						

Continued from previous page...

 1 Number of models with violations






10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram : Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models



in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	9	1.79	0.36	1.87
(1,35)	1:27:A:TYR:N	1:27:A:TYR:CA	1:27:A:TYR:C	1:28:A:CYS:N	5	1.64	0.42	1.67
(1,28)	1:23:A:ILE:C	1:24:A:ARG:N	1:24:A:ARG:CA	1:24:A:ARG:C	3	1.54	0.29	1.58
(1,39)	1:29:A:CYS:N	1:29:A:CYS:CA	1:29:A:CYS:C	1:30:A:PRO:N	3	1.35	0.25	1.35
(1,25)	1:19:A:ALA:N	1:19:A:ALA:CA	1:19:A:ALA:C	1:20:A:THR:N	3	1.1	0.06	1.11
(1,11)	1:7:A:ARG:C	1:8:A:LYS:N	1:8:A:LYS:CA	1:8:A:LYS:C	3	1.09	0.05	1.12


 1 Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	12	2.42
(1,33)	1:26:A:LEU:N	1:26:A:LEU:CA	1:26:A:LEU:C	1:27:A:TYR:N	19	2.36
(1,35)	1:27:A:TYR:N	1:27:A:TYR:CA	1:27:A:TYR:C	1:28:A:CYS:N	3	2.33
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	9	2.09
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	3	1.93
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	4	1.93
(1,28)	1:23:A:ILE:C	1:24:A:ARG:N	1:24:A:ARG:CA	1:24:A:ARG:C	6	1.87
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	2	1.87
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	20	1.87
(1,35)	1:27:A:TYR:N	1:27:A:TYR:CA	1:27:A:TYR:C	1:28:A:CYS:N	18	1.76
(1,35)	1:27:A:TYR:N	1:27:A:TYR:CA	1:27:A:TYR:C	1:28:A:CYS:N	6	1.67
(1,39)	1:29:A:CYS:N	1:29:A:CYS:CA	1:29:A:CYS:C	1:30:A:PRO:N	19	1.65
(1,28)	1:23:A:ILE:C	1:24:A:ARG:N	1:24:A:ARG:CA	1:24:A:ARG:C	1	1.58
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	10	1.46
(1,30)	1:24:A:ARG:C	1:25:A:PHE:N	1:25:A:PHE:CA	1:25:A:PHE:C	10	1.4
(1,39)	1:29:A:CYS:N	1:29:A:CYS:CA	1:29:A:CYS:C	1:30:A:PRO:N	16	1.35
(1,13)	1:9:A:GLY:N	1:9:A:GLY:CA	1:9:A:GLY:C	1:10:A:HIS:N	16	1.35
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	14	1.33
(1,35)	1:27:A:TYR:N	1:27:A:TYR:CA	1:27:A:TYR:C	1:28:A:CYS:N	8	1.31
(1,40)	1:30:A:PRO:N	1:30:A:PRO:CA	1:30:A:PRO:C	1:31:A:ARG:N	10	1.29
(1,4)	1:3:A:LEU:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	7	1.25

Continued on next page...



Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation ($^{\circ}$)
(1,28)	1:23:A:ILE:C	1:24:A:ARG:N	1:24:A:ARG:CA	1:24:A:ARG:C	19	1.17
(1,25)	1:19:A:ALA:N	1:19:A:ALA:CA	1:19:A:ALA:C	1:20:A:THR:N	17	1.17
(1,11)	1:7:A:ARG:C	1:8:A:LYS:N	1:8:A:LYS:CA	1:8:A:LYS:C	10	1.13
(1,11)	1:7:A:ARG:C	1:8:A:LYS:N	1:8:A:LYS:CA	1:8:A:LYS:C	1	1.12
(1,35)	1:27:A:TYR:N	1:27:A:TYR:CA	1:27:A:TYR:C	1:28:A:CYS:N	5	1.11
(1,25)	1:19:A:ALA:N	1:19:A:ALA:CA	1:19:A:ALA:C	1:20:A:THR:N	12	1.11
(1,2)	1:2:A:LEU:N	1:2:A:LEU:CA	1:2:A:LEU:C	1:3:A:LEU:N	9	1.1
(1,39)	1:29:A:CYS:N	1:29:A:CYS:CA	1:29:A:CYS:C	1:30:A:PRO:N	6	1.04
(1,25)	1:19:A:ALA:N	1:19:A:ALA:CA	1:19:A:ALA:C	1:20:A:THR:N	13	1.02
(1,11)	1:7:A:ARG:C	1:8:A:LYS:N	1:8:A:LYS:CA	1:8:A:LYS:C	13	1.02

Continued from previous page...

