



Full wwPDB X-ray Structure Validation Report i

Aug 22, 2023 – 06:07 PM EDT

PDB ID : 7UVS
Title : Pfs230 domain 1 bound by LMIV230-02 Fab
Authors : Ivanochko, D.; Julien, J.P.
Deposited on : 2022-05-02
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	199	<div style="width: 39%;">39%</div> <div style="width: 84%; background-color: green;">84%</div> <div style="width: 14%; background-color: grey;">• 14%</div>

2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 10235 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LMIV230-02 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C 1662	N 1045	O 275	S 336	6	0	0
1	D	223	Total	C 1640	N 1034	O 271	S 330	5	0	0

- Molecule 2 is a protein called LMIV230-02 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	221	Total	C 1710	N 1071	O 289	S 344	6	0	0
2	E	220	Total	C 1704	N 1068	O 288	S 343	5	0	0

- Molecule 3 is a protein called Gametocyte surface protein P230.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	180	Total	C 1431	N 915	O 222	S 290	4	0	0
3	F	172	Total	C 1374	N 881	O 213	S 276	4	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	585	GLN	ASN	conflict	UNP P68874
C	732	GLY	-	expression tag	UNP P68874
C	733	SER	-	expression tag	UNP P68874
C	734	LEU	-	expression tag	UNP P68874
C	735	LYS	-	expression tag	UNP P68874
C	736	GLU	-	expression tag	UNP P68874
C	737	ASN	-	expression tag	UNP P68874
C	738	LEU	-	expression tag	UNP P68874

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	739	TYR	-	expression tag	UNP P68874
C	740	PHE	-	expression tag	UNP P68874
C	741	GLN	-	expression tag	UNP P68874
C	742	GLY	-	expression tag	UNP P68874
C	743	TRP	-	expression tag	UNP P68874
C	744	SER	-	expression tag	UNP P68874
C	745	HIS	-	expression tag	UNP P68874
C	746	PRO	-	expression tag	UNP P68874
C	747	GLN	-	expression tag	UNP P68874
C	748	PHE	-	expression tag	UNP P68874
C	749	GLU	-	expression tag	UNP P68874
C	750	LYS	-	expression tag	UNP P68874
F	585	GLN	ASN	conflict	UNP P68874
F	732	GLY	-	expression tag	UNP P68874
F	733	SER	-	expression tag	UNP P68874
F	734	LEU	-	expression tag	UNP P68874
F	735	LYS	-	expression tag	UNP P68874
F	736	GLU	-	expression tag	UNP P68874
F	737	ASN	-	expression tag	UNP P68874
F	738	LEU	-	expression tag	UNP P68874
F	739	TYR	-	expression tag	UNP P68874
F	740	PHE	-	expression tag	UNP P68874
F	741	GLN	-	expression tag	UNP P68874
F	742	GLY	-	expression tag	UNP P68874
F	743	TRP	-	expression tag	UNP P68874
F	744	SER	-	expression tag	UNP P68874
F	745	HIS	-	expression tag	UNP P68874
F	746	PRO	-	expression tag	UNP P68874
F	747	GLN	-	expression tag	UNP P68874
F	748	PHE	-	expression tag	UNP P68874
F	749	GLU	-	expression tag	UNP P68874
F	750	LYS	-	expression tag	UNP P68874

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	147	Total O 147 147	0	0
5	B	140	Total O 140 140	0	0
5	C	123	Total O 123 123	0	0
5	D	122	Total O 122 122	0	0
5	E	166	Total O 166 166	0	0
5	F	15	Total O 15 15	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LMIV230-02 Fab heavy chain




- Molecule 1: LMIV230-02 Fab heavy chain




- Molecule 2: LMIV230-02 Fab light chain



- Molecule 2: LMIV230-02 Fab light chain




- Molecule 3: Gametocyte surface protein P230





- Molecule 3: Gametocyte surface protein P230



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.36 Å 132.47 Å 125.07 Å 90.00° 102.35° 90.00°	Depositor
Resolution (Å)	29.85 – 2.06 29.85 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.85-2.06) 99.1 (29.85-2.06)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.01 (at 2.06 Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R , R_{free}	0.174 , 0.215 0.174 , 0.215	Depositor DCC
R_{free} test set	2019 reflections (2.25%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 54.2	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10235	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1703	0.64	0/2320
1	D	0.39	0/1681	0.57	0/2292
2	B	0.41	0/1748	0.60	0/2373
2	E	0.43	0/1742	0.61	0/2365
3	C	0.46	0/1457	0.68	0/1970
3	F	0.34	0/1400	0.64	0/1894
All	All	0.42	0/9731	0.62	0/13214

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1662	0	1616	6	0
1	D	1640	0	1596	4	0
2	B	1710	0	1660	2	0
2	E	1704	0	1656	2	0
3	C	1431	0	1437	3	0
3	F	1374	0	1384	3	0
4	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	147	0	0	0	0
5	B	140	0	0	0	0
5	C	123	0	0	0	0
5	D	122	0	0	0	0
5	E	166	0	0	0	0
5	F	15	0	0	0	0
All	All	10235	0	9349	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:67:VAL:HG22	1:D:82:VAL:HG22	1.85	0.59
2:B:78:LEU:HD11	2:B:104:LEU:HD21	1.90	0.53
1:A:30:ASN:O	1:A:53:SER:OG	2.28	0.52
2:E:54:ARG:NH1	2:E:62:PHE:O	2.44	0.49
1:A:67:VAL:HG22	1:A:82:VAL:HG22	1.94	0.48
3:C:712:GLU:HG2	3:C:717:LYS:HG2	1.96	0.47
3:C:647:PRO:HB3	3:C:688:ILE:HD12	1.96	0.47
3:F:707:ASP:OD2	3:F:720:ARG:NH1	2.49	0.46
1:A:83:ARG:NH1	1:A:85:GLU:OE1	2.49	0.45
3:C:653:LYS:HE2	3:C:653:LYS:HB3	1.75	0.44
3:F:656:THR:HB	3:F:657:LYS:HE2	1.99	0.44
1:D:60:ALA:HB3	1:D:63:PHE:HD2	1.84	0.43
1:A:12:LYS:HE2	1:A:18:VAL:HG23	2.01	0.42
1:D:12:LYS:HG3	1:D:18:VAL:HB	2.02	0.42
1:A:119:PRO:HB3	1:A:145:TYR:HB3	2.03	0.41
1:A:128:SER:N	1:A:216:CYS:OXT	2.54	0.41
2:B:40:PRO:HG2	2:B:166:GLU:HG3	2.02	0.41
2:E:27(D):TYR:CZ	2:E:27(F):SER:HB2	2.56	0.40
1:D:145:TYR:CE2	1:D:150:VAL:HG13	2.56	0.40
3:F:705:ILE:HD13	3:F:720:ARG:NH2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	224/226 (99%)	222 (99%)	2 (1%)	0	100 100
1	D	221/226 (98%)	219 (99%)	2 (1%)	0	100 100
2	B	219/221 (99%)	213 (97%)	6 (3%)	0	100 100
2	E	218/221 (99%)	212 (97%)	6 (3%)	0	100 100
3	C	178/199 (89%)	173 (97%)	5 (3%)	0	100 100
3	F	170/199 (85%)	163 (96%)	7 (4%)	0	100 100
All	All	1230/1292 (95%)	1202 (98%)	28 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	187/187 (100%)	186 (100%)	1 (0%)	88 89
1	D	184/187 (98%)	184 (100%)	0	100 100
2	B	194/194 (100%)	192 (99%)	2 (1%)	76 75
2	E	193/194 (100%)	193 (100%)	0	100 100
3	C	167/185 (90%)	167 (100%)	0	100 100
3	F	161/185 (87%)	161 (100%)	0	100 100
All	All	1086/1132 (96%)	1083 (100%)	3 (0%)	92 93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	PHE
2	B	60	ASP
2	B	215	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/226 (100%)	0.36	15 (6%) 18 18	24, 36, 81, 151	0
1	D	223/226 (98%)	0.53	30 (13%) 3 2	24, 43, 95, 130	0
2	B	221/221 (100%)	-0.08	3 (1%) 75 76	24, 36, 58, 125	0
2	E	220/221 (99%)	-0.05	3 (1%) 75 76	23, 37, 61, 82	0
3	C	180/199 (90%)	0.32	13 (7%) 15 16	23, 35, 71, 99	0
3	F	172/199 (86%)	2.13	77 (44%) 0 0	58, 88, 126, 160	0
All	All	1242/1292 (96%)	0.48	141 (11%) 5 5	23, 41, 102, 160	0

All (141) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	216	CYS	10.6
3	F	562	LEU	9.0
1	A	1	GLN	8.1
3	F	577	ASP	7.7
1	A	215	SER	7.6
1	D	132	SER	7.1
1	A	27	GLY	7.1
3	F	697	HIS	6.7
3	F	714	ASP	6.6
1	D	133	GLY	6.4
1	D	130	SER	6.2
3	F	695	VAL	6.1
3	F	730	TYR	6.0
3	F	704	PHE	5.9
1	D	27	GLY	5.9
1	D	26	GLY	5.9
3	F	705	ILE	5.8
3	F	578	SER	5.7
3	F	563	SER	5.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	728	GLU	5.5
3	F	656	THR	5.5
3	F	696	VAL	5.5
3	F	583	ALA	5.4
3	F	573	ALA	5.1
3	F	715	ASN	5.0
3	F	560	ILE	4.9
1	D	30	ASN	4.8
1	A	128	SER	4.8
2	B	215	CYS	4.8
3	F	604	SER	4.7
1	D	131	THR	4.7
1	D	28	THR	4.6
3	F	579	TYR	4.6
3	F	722	ILE	4.4
3	F	616	ASN	4.4
3	C	554	VAL	4.4
3	F	721	GLY	4.3
3	F	565	GLU	4.3
3	F	561	ASP	4.3
3	F	587	THR	4.3
3	F	698	LYS	4.3
1	D	25	SER	4.3
3	F	567	THR	4.2
3	F	564	TYR	4.2
1	D	24	ALA	4.2
1	D	75	THR	4.1
3	F	712	GLU	4.1
1	D	74	SER	4.0
3	F	605	GLY	4.0
3	F	668	TYR	3.7
3	F	602	THR	3.7
3	F	699	ALA	3.7
3	F	588	ASN	3.7
3	C	732	GLY	3.6
3	F	603	GLU	3.6
3	F	618	PRO	3.6
1	A	26	GLY	3.5
3	F	606	PRO	3.5
3	F	706	CYS	3.4
3	F	723	VAL	3.4
3	F	729	PRO	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	582	TYR	3.3
1	D	190	GLY	3.3
3	C	556	GLU	3.2
3	C	603	GLU	3.2
3	F	575	SER	3.2
1	A	131	THR	3.2
1	A	28	THR	3.2
3	F	580	ASP	3.2
3	F	694	PRO	3.2
3	F	727	VAL	3.2
3	F	713	ASP	3.2
1	D	64	GLN	3.1
3	C	559	LYS	3.1
1	D	191	THR	3.1
1	A	2	VAL	3.1
3	F	657	LYS	3.0
3	F	726	TYR	3.0
3	F	607	LYS	2.9
2	E	1	GLU	2.9
1	A	29	PHE	2.9
3	F	615	VAL	2.9
3	F	589	LYS	2.9
3	F	572	THR	2.9
1	A	127	SER	2.9
3	C	555	ASP	2.9
3	C	562	LEU	2.9
3	F	619	LEU	2.9
3	F	667	ILE	2.8
3	F	676	VAL	2.8
3	F	658	LEU	2.8
3	F	678	GLU	2.8
3	F	581	LYS	2.8
1	D	61	GLN	2.8
3	C	725	VAL	2.7
1	D	128	SER	2.7
3	F	703	TYR	2.7
2	E	134	VAL	2.7
1	D	65	GLY	2.7
2	B	213	GLY	2.7
3	C	557	LEU	2.7
1	D	186	SER	2.6
3	F	655	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	178	LEU	2.6
3	F	595	PHE	2.6
1	D	187	SER	2.6
3	F	599	LEU	2.6
1	A	157	GLY	2.6
1	A	132	SER	2.6
1	D	29	PHE	2.5
3	F	716	LYS	2.5
3	F	677	ASN	2.5
3	C	731	GLY	2.5
1	D	100(D)	GLY	2.5
3	F	617	GLU	2.4
1	A	214	LYS	2.4
3	F	597	ASP	2.4
3	F	700	THR	2.4
3	C	588	ASN	2.3
1	D	2	VAL	2.3
3	F	569	SER	2.3
1	D	57	ALA	2.3
3	F	711	THR	2.3
3	F	576	GLU	2.3
2	B	155	LEU	2.2
3	F	591	TYR	2.2
1	A	73	THR	2.2
2	E	70	ASP	2.2
3	F	659	LYS	2.2
1	D	142	VAL	2.1
1	D	54	PHE	2.1
1	D	3	GLN	2.1
1	D	129	LYS	2.1
3	F	654	GLU	2.1
3	C	723	VAL	2.1
3	F	725	VAL	2.1
1	D	72	ASP	2.0
3	C	672	ILE	2.0
3	F	683	PHE	2.0
3	F	566	THR	2.0
3	F	679	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NA	C	801	1/1	0.96	0.13	52,52,52,52	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.