



# Full wwPDB X-ray Structure Validation Report i

Nov 8, 2022 – 08:39 PM JST

PDB ID : 7VZS  
Title : FAD-dependent Glucose Dehydrogenase complexed with an inhibitor at pH7.56  
Authors : Nakajima, Y.  
Deposited on : 2021-11-16  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	572	2%	95% ..
1	B	572	%	95% 5% .

## 2 Entry composition [\(i\)](#)

There are 7 unique types of molecules in this entry. The entry contains 9682 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GMC oxidoreductase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	566	Total	C 4328	N 2738	O 747	S 835	8	0
1	B	568	Total	C 4359	N 2761	O 754	S 836	8	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A0A1S9DW10
B	1	MET	-	initiating methionine	UNP A0A1S9DW10

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C 53	N 27	O 9	P 15	2

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P			
2	B	1	53	27	9	15	2	0	0	

- Molecule 3 is D-glucal (three-letter code: DGO) (formula: C<sub>6</sub>H<sub>10</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	O					
3	A	1	10	6	4				0	0
3	A	1	10	6	4				0	0
3	A	1	10	6	4				0	0
3	B	1	10	6	4				0	0
3	B	1	10	6	4				0	0
3	B	1	10	6	4				0	0
3	B	1	10	6	4				0	0
3	B	1	10	6	4				0	0

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total C N O S					0	0
			15 8 2 4 1						
4	A	1	Total C N O S					0	0
			15 8 2 4 1						

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total O S			0	0
			5 4 1				
5	B	1	Total O S			0	0
			5 4 1				

- Molecule 6 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C<sub>10</sub>H<sub>22</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 16 10 6	0	0
6	B	1	Total C O 16 10 6	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	363	Total O 363 363	0	0
7	B	374	Total O 374 374	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GMC oxidoreductase



- Molecule 1: GMC oxidoreductase



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.93Å    67.08Å    150.68Å 90.00°    97.13°    90.00°	Depositor
Resolution (Å)	20.00 – 1.80 40.01 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.3 (20.00-1.80) 95.2 (40.01-1.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.28 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
$R$ , $R_{free}$	0.152 , 0.190 0.155 , 0.195	Depositor DCC
$R_{free}$ test set	4626 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.5	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 47.1	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9682	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, DGO, SO4, FAD, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.57	0/4436	0.63	0/6040
1	B	0.60	0/4472	0.63	0/6089
All	All	0.58	0/8908	0.63	0/12129

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4328	0	4276	15	0
1	B	4359	0	4305	22	0
2	A	53	0	31	2	0
2	B	53	0	31	3	0
3	A	30	0	0	0	0
3	B	50	0	0	0	0
4	A	30	0	34	3	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	16	0	22	0	0
6	B	16	0	22	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	363	0	0	0	0
7	B	374	0	0	1	0
All	All	9682	0	8721	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:202:SER:HB2	1:B:204:LEU:HD23	1.73	0.71
1:B:280:ARG:HE	6:B:608:1PE:H262	1.61	0.65
1:B:94:ASN:HB2	2:B:601:FAD:C5X	2.30	0.62
4:A:606:EPE:H62	1:B:58:PHE:O	2.00	0.61
1:A:94:ASN:HB2	2:A:601:FAD:C5X	2.34	0.58
1:B:284:ILE:HG13	6:B:608:1PE:H261	1.85	0.57
1:B:549:HIS:HB3	2:B:601:FAD:C2	2.35	0.56
1:A:506[A]:HIS:ND1	1:A:545:GLN:OE1	2.42	0.52
1:A:549:HIS:HB3	2:A:601:FAD:C2	2.40	0.52
1:A:350:ASP:OD1	1:B:478:LYS:NZ	2.43	0.51
1:B:551:VAL:HG22	2:B:601:FAD:O2	2.12	0.50
1:B:345:PHE:CD2	1:B:352:ILE:HD13	2.47	0.49
1:B:424:GLY:HA3	1:B:440:ASN:O	2.13	0.48
1:B:506[A]:HIS:CE1	1:B:550:LEU:HG	2.50	0.47
1:A:153:LYS:O	4:A:605:EPE:H32	2.16	0.46
1:B:506[B]:HIS:HE1	1:B:549:HIS:HA	1.79	0.46
1:B:280:ARG:HE	6:B:608:1PE:C15	2.30	0.45
1:B:283:LEU:HD13	6:B:608:1PE:H252	1.99	0.44
1:B:280:ARG:CZ	6:B:608:1PE:H242	2.48	0.44
1:A:281:SER:N	1:A:282:PRO:HD2	2.33	0.44
1:A:424:GLY:HA3	1:A:440:ASN:O	2.17	0.44
1:B:111:LYS:NZ	7:B:702:HOH:O	2.41	0.44
1:B:526[B]:LEU:HD22	1:B:537:VAL:HG22	1.99	0.43
1:A:99:THR:HA	1:A:190:ASN:OD1	2.18	0.43
1:B:282:PRO:HG3	1:B:510:THR:HB	2.01	0.43
1:A:402:LEU:O	1:A:413:SER:HA	2.19	0.43
1:B:283:LEU:HD12	6:B:608:1PE:H151	2.00	0.43
1:B:299:ASN:OD1	1:B:430:SER:HA	2.19	0.42
1:A:506[B]:HIS:HE1	1:A:549:HIS:HA	1.84	0.41
1:A:432:ASP:HA	1:A:433:PRO:HD2	1.86	0.41
4:A:606:EPE:H22	4:A:606:EPE:H101	1.79	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:479:PRO:HB3	1:A:496:TRP:CE2	2.56	0.41
1:A:526[B]:LEU:HD12	1:A:561:ALA:HB2	2.03	0.41
1:B:281:SER:N	1:B:282:PRO:HD2	2.35	0.41
1:B:10:ILE:HD13	1:B:266:VAL:HG23	2.04	0.40
1:A:199:ILE:N	1:A:199:ILE:HD12	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	569/572 (100%)	557 (98%)	11 (2%)	1 (0%)	47 33
1	B	572/572 (100%)	558 (98%)	14 (2%)	0	100 100
All	All	1141/1144 (100%)	1115 (98%)	25 (2%)	1 (0%)	51 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	347	ASN

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	458/458 (100%)	457 (100%)	1 (0%)	93 92
1	B	459/458 (100%)	458 (100%)	1 (0%)	93 92
All	All	917/916 (100%)	915 (100%)	2 (0%)	93 92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	475	LYS
1	B	395	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	237	ASN
1	B	243	ASN
1	B	347	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	DGO	B	604	-	10,10,10	2.14	1 (10%)	12,13,13	2.04	4 (33%)
5	SO4	A	607	-	4,4,4	0.18	0	6,6,6	0.23	0
3	DGO	A	604	-	10,10,10	2.16	1 (10%)	12,13,13	2.33	4 (33%)
3	DGO	B	605	-	10,10,10	2.33	1 (10%)	12,13,13	2.67	5 (41%)
3	DGO	B	602	-	10,10,10	2.04	1 (10%)	12,13,13	2.66	2 (16%)
3	DGO	B	606	-	10,10,10	2.26	1 (10%)	12,13,13	2.31	3 (25%)
4	EPE	A	605	-	15,15,15	0.84	1 (6%)	18,20,20	1.79	3 (16%)
4	EPE	A	606	-	15,15,15	0.99	1 (6%)	18,20,20	2.17	7 (38%)
6	1PE	B	608	-	15,15,15	0.49	0	14,14,14	0.46	0
3	DGO	A	602	-	10,10,10	2.22	1 (10%)	12,13,13	2.61	2 (16%)
2	FAD	B	601	-	53,58,58	1.08	3 (5%)	68,89,89	1.51	11 (16%)
2	FAD	A	601	-	53,58,58	1.18	4 (7%)	68,89,89	1.43	9 (13%)
6	1PE	A	608	-	15,15,15	0.49	0	14,14,14	0.35	0
3	DGO	A	603	-	10,10,10	2.16	1 (10%)	12,13,13	1.99	3 (25%)
5	SO4	B	607	-	4,4,4	0.31	0	6,6,6	0.25	0
3	DGO	B	603	-	10,10,10	2.28	1 (10%)	12,13,13	2.45	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DGO	B	604	-	-	0/2/16/16	0/1/1/1
3	DGO	A	604	-	-	0/2/16/16	0/1/1/1
3	DGO	B	605	-	-	0/2/16/16	0/1/1/1
3	DGO	B	602	-	-	0/2/16/16	0/1/1/1
3	DGO	B	606	-	-	2/2/16/16	0/1/1/1
4	EPE	A	605	-	-	1/9/19/19	0/1/1/1
4	EPE	A	606	-	-	8/9/19/19	0/1/1/1
6	1PE	B	608	-	-	6/13/13/13	-
3	DGO	A	602	-	-	0/2/16/16	0/1/1/1
2	FAD	B	601	-	-	4/30/50/50	0/6/6/6
2	FAD	A	601	-	-	5/30/50/50	0/6/6/6

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	1PE	A	608	-	-	8/13/13/13	-
3	DGO	A	603	-	-	0/2/16/16	0/1/1/1
3	DGO	B	603	-	-	0/2/16/16	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	605	DGO	O5-C1	6.96	1.47	1.36
3	B	606	DGO	O5-C1	6.83	1.46	1.36
3	B	603	DGO	O5-C1	6.81	1.46	1.36
3	A	602	DGO	O5-C1	6.57	1.46	1.36
3	A	603	DGO	O5-C1	6.43	1.46	1.36
3	B	604	DGO	O5-C1	6.36	1.46	1.36
3	A	604	DGO	O5-C1	6.26	1.45	1.36
3	B	602	DGO	O5-C1	6.16	1.45	1.36
2	A	601	FAD	C4X-N5	4.64	1.39	1.30
2	B	601	FAD	C4X-N5	4.30	1.39	1.30
2	A	601	FAD	C2A-N3A	3.97	1.38	1.32
4	A	606	EPE	C10-S	3.43	1.82	1.77
2	B	601	FAD	C2A-N3A	3.32	1.37	1.32
4	A	605	EPE	C10-S	2.88	1.81	1.77
2	A	601	FAD	C2A-N1A	2.81	1.39	1.33
2	A	601	FAD	C10-N1	2.23	1.37	1.33
2	B	601	FAD	C2A-N1A	2.22	1.38	1.33

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	DGO	O5-C1-C2	-7.25	118.44	125.26
2	A	601	FAD	N3A-C2A-N1A	-6.78	118.08	128.68
2	B	601	FAD	N3A-C2A-N1A	-6.25	118.91	128.68
3	B	605	DGO	O5-C1-C2	-6.10	119.52	125.26
3	B	602	DGO	O5-C1-C2	-6.08	119.53	125.26
3	B	602	DGO	C5-O5-C1	5.95	122.31	114.65
3	A	603	DGO	O5-C1-C2	-5.68	119.92	125.26
3	B	606	DGO	O5-C1-C2	-5.49	120.09	125.26
3	A	604	DGO	O5-C1-C2	-5.46	120.12	125.26
3	B	603	DGO	O5-C1-C2	-5.21	120.36	125.26
4	A	606	EPE	C5-N4-C3	5.20	120.54	108.83
3	A	602	DGO	C5-O5-C1	4.72	120.73	114.65
3	B	604	DGO	O5-C1-C2	-4.71	120.83	125.26
3	B	606	DGO	C5-O5-C1	4.65	120.64	114.65

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	605	DGO	C5-O5-C1	4.33	120.22	114.65
3	B	603	DGO	C5-O5-C1	4.19	120.05	114.65
3	B	604	DGO	C5-O5-C1	4.07	119.90	114.65
3	A	604	DGO	C5-O5-C1	4.07	119.89	114.65
3	B	603	DGO	O5-C5-C4	-4.05	105.41	110.83
2	B	601	FAD	C9A-C5X-N5	-3.94	118.15	122.43
4	A	605	EPE	O1S-S-C10	3.94	111.66	106.92
4	A	605	EPE	C5-N4-C3	3.93	117.69	108.83
3	B	605	DGO	O5-C5-C4	-3.28	106.44	110.83
4	A	605	EPE	C7-N4-C5	3.21	119.44	111.23
4	A	606	EPE	O1S-S-C10	3.09	110.63	106.92
3	A	604	DGO	O5-C5-C4	-3.08	106.70	110.83
4	A	606	EPE	C7-N4-C5	3.07	119.09	111.23
2	A	601	FAD	C4-C4X-N5	2.99	122.49	118.23
4	A	606	EPE	C7-N4-C3	2.97	118.84	111.23
3	B	605	DGO	C3-C2-C1	-2.95	118.38	122.23
2	A	601	FAD	C4X-C10-N10	2.89	120.71	116.48
2	A	601	FAD	C4X-C4-N3	2.80	120.31	113.19
2	B	601	FAD	C4-N3-C2	-2.79	120.48	125.64
2	B	601	FAD	C4-C4X-N5	2.73	122.12	118.23
4	A	606	EPE	O2S-S-C10	2.73	110.20	106.92
2	B	601	FAD	C4X-C4-N3	2.72	120.09	113.19
2	A	601	FAD	C4-N3-C2	-2.69	120.67	125.64
2	A	601	FAD	C10-C4X-N5	-2.69	119.15	124.86
3	A	603	DGO	C5-O5-C1	2.60	118.00	114.65
3	B	605	DGO	O5-C5-C6	2.56	110.48	106.53
2	A	601	FAD	C1B-N9A-C4A	-2.54	122.18	126.64
3	B	606	DGO	O5-C5-C6	2.51	110.41	106.53
2	B	601	FAD	C4X-C10-N10	2.45	120.06	116.48
2	B	601	FAD	C10-C4X-N5	-2.45	119.67	124.86
2	B	601	FAD	C10-N1-C2	2.33	121.56	116.90
3	A	604	DGO	O5-C5-C6	2.29	110.07	106.53
3	B	603	DGO	O5-C5-C6	2.23	109.97	106.53
2	A	601	FAD	C9A-C5X-N5	-2.22	120.02	122.43
4	A	606	EPE	O3S-S-C10	2.20	109.33	105.77
2	B	601	FAD	C5X-C9A-N10	2.20	120.22	117.95
2	B	601	FAD	O4-C4-C4X	-2.16	120.86	126.60
2	B	601	FAD	C2A-N1A-C6A	2.13	122.40	118.75
3	B	604	DGO	O5-C5-C6	2.11	109.78	106.53
3	A	603	DGO	O5-C5-C4	-2.10	108.01	110.83
3	B	604	DGO	O5-C5-C4	-2.08	108.04	110.83
4	A	606	EPE	C6-C5-N4	2.07	114.89	110.64

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	601	FAD	O4B-C1B-C2B	-2.03	103.96	106.93

There are no chirality outliers.

All (34) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	FAD	N10-C1'-C2'-O2'
2	A	601	FAD	PA-O3P-P-O5'
2	B	601	FAD	N10-C1'-C2'-O2'
2	B	601	FAD	PA-O3P-P-O5'
4	A	606	EPE	C10-C9-N1-C2
4	A	606	EPE	C10-C9-N1-C6
4	A	606	EPE	C9-C10-S-O2S
6	B	608	1PE	C12-C22-OH3-C23
6	B	608	1PE	OH2-C12-C22-OH3
6	B	608	1PE	OH6-C15-C25-OH5
4	A	606	EPE	N4-C7-C8-O8
4	A	606	EPE	C9-C10-S-O3S
6	B	608	1PE	C15-C25-OH5-C14
4	A	606	EPE	S-C10-C9-N1
6	B	608	1PE	C24-C14-OH5-C25
4	A	606	EPE	C9-C10-S-O1S
2	B	601	FAD	N10-C1'-C2'-C3'
6	A	608	1PE	C12-C22-OH3-C23
6	A	608	1PE	C14-C24-OH4-C13
3	B	606	DGO	C4-C5-C6-O6
4	A	606	EPE	C8-C7-N4-C5
2	A	601	FAD	P-O3P-PA-O1A
2	A	601	FAD	P-O3P-PA-O2A
3	B	606	DGO	O5-C5-C6-O6
6	A	608	1PE	C23-C13-OH4-C24
6	A	608	1PE	C15-C25-OH5-C14
2	B	601	FAD	O4B-C4B-C5B-O5B
6	A	608	1PE	OH4-C13-C23-OH3
6	A	608	1PE	C24-C14-OH5-C25
6	A	608	1PE	OH7-C16-C26-OH6
6	A	608	1PE	OH5-C14-C24-OH4
6	B	608	1PE	OH4-C13-C23-OH3
4	A	605	EPE	C8-C7-N4-C5
2	A	601	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.


5 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	605	EPE	1	0
4	A	606	EPE	2	0
6	B	608	1PE	6	0
2	B	601	FAD	3	0
2	A	601	FAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	566/572 (98%)	-0.27	10 (1%) 68 64	13, 20, 32, 45	0
1	B	568/572 (99%)	-0.35	8 (1%) 75 72	12, 18, 29, 41	0
All	All	1134/1144 (99%)	-0.31	18 (1%) 72 68	12, 19, 31, 45	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	245	SER	6.0
1	A	247	GLU	4.7
1	A	245	SER	4.4
1	A	246	ALA	4.0
1	B	246	ALA	4.0
1	B	207	ASP	3.7
1	A	329	VAL	3.4
1	A	259	ALA	3.1
1	B	185[A]	TRP	2.8
1	B	3	ASN	2.7
1	A	407	GLY	2.5
1	A	327	TYR	2.4
1	A	506[A]	HIS	2.3
1	B	247	GLU	2.2
1	A	408	GLY	2.2
1	B	167	SER	2.1
1	B	244	GLY	2.1
1	A	372	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	1PE	B	608	16/16	0.74	0.19	40,47,50,50	0
4	EPE	A	606	15/15	0.83	0.19	61,64,64,65	0
3	DGO	B	603	10/10	0.88	0.24	28,34,35,37	0
6	1PE	A	608	16/16	0.88	0.15	37,44,51,52	0
3	DGO	B	606	10/10	0.88	0.19	41,42,43,43	0
3	DGO	A	604	10/10	0.91	0.22	30,36,37,37	0
3	DGO	A	602	10/10	0.91	0.13	28,29,30,32	0
5	SO4	B	607	5/5	0.92	0.14	33,37,38,40	0
5	SO4	A	607	5/5	0.93	0.12	33,33,36,38	0
4	EPE	A	605	15/15	0.95	0.14	35,36,40,41	0
3	DGO	B	605	10/10	0.95	0.15	26,29,30,30	0
3	DGO	B	602	10/10	0.96	0.10	23,24,26,28	0
3	DGO	A	603	10/10	0.96	0.08	29,30,30,32	0
3	DGO	B	604	10/10	0.97	0.07	22,23,24,25	0
2	FAD	A	601	53/53	0.98	0.11	11,14,16,17	0
2	FAD	B	601	53/53	0.98	0.11	11,13,15,17	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.























## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.