



# Full wwPDB X-ray Structure Validation Report i

Oct 9, 2023 – 08:06 AM EDT

PDB ID : 8DAO  
Title : Crystal structure of SARS-CoV-2 spike stem fusion peptide in complex with neutralizing antibody COV44-79  
Authors : Lin, T.H.; Lee, C.C.D.; Yuan, M.; Wilson, I.A.  
Deposited on : 2022-06-13  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...



## 2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 6914 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called COV44-79 heavy chain variable domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	126	959	604	168	182	5	0	0	0
1	A	126	959	604	168	182	5	0	0	0

- Molecule 2 is a protein called COV44-79 light chain variable domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	109	838	526	143	165	4	0	0	0
2	B	109	838	526	143	165	4	0	0	0

- Molecule 3 is a protein called COV44-79 heavy chain constant domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	103	738	465	121	149	3	0	0	0
3	G	103	738	465	121	149	3	0	0	0

- Molecule 4 is a protein called COV44-79 light chain constant domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	F	106	820	510	139	169	2	0	0	0
4	H	106	820	510	139	169	2	0	0	0


- Molecule 5 is a protein called Spike protein S2 fusion peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	I	12	Total	C	N	O	0	0	0
			102	68	16	18			
5	J	12	Total	C	N	O	0	0	0
			102	68	16	18			


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: COV44-79 heavy chain variable domain




- Molecule 1: COV44-79 heavy chain variable domain




- Molecule 2: COV44-79 light chain variable domain




- Molecule 2: COV44-79 light chain variable domain




- Molecule 3: COV44-79 heavy chain constant domain




- Molecule 3: COV44-79 heavy chain constant domain



- Molecule 4: COV44-79 light chain constant domain



- Molecule 4: COV44-79 light chain constant domain



- Molecule 5: Spike protein S2 fusion peptide



- Molecule 5: Spike protein S2 fusion peptide



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.37 Å    100.37 Å    229.94 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	49.88 – 2.80 49.88 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.5 (49.88-2.80) 98.5 (49.88-2.80)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.13 (at 2.81 Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
$R$ , $R_{free}$	0.231 , 0.284 0.229 , 0.281	Depositor DCC
$R_{free}$ test set	1457 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.9	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 58.7	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.48$ , $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6914	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.25	0/978	0.53	0/1326
1	C	0.25	0/978	0.53	0/1326
2	B	0.28	0/858	0.56	0/1164
2	D	0.26	0/858	0.52	0/1164
3	E	0.27	0/756	0.50	0/1035
3	G	0.31	0/756	0.52	0/1035
4	F	0.25	0/836	0.47	0/1134
4	H	0.25	0/836	0.49	0/1134
5	I	0.25	0/104	0.43	0/137
5	J	0.26	0/104	0.47	0/137
All	All	0.26	0/7064	0.51	0/9592

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	959	0	956	19	0
1	C	959	0	956	13	0
2	B	838	0	815	8	0
2	D	838	0	813	16	0
3	E	738	0	732	7	0
3	G	738	0	732	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	820	0	793	8	0
4	H	820	0	793	17	0
5	I	102	0	104	6	0
5	J	102	0	104	7	0
All	All	6914	0	6798	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:34:ALA:HB3	2:B:89:GLN:HG2	1.60	0.82
2:D:34:ALA:HB3	2:D:89:GLN:HG2	1.67	0.76
3:G:116:THR:HG22	3:G:147:PRO:HD3	1.69	0.74
1:A:91:THR:HG23	1:A:123:SER:HA	1.70	0.74
5:I:818:ILE:HA	5:I:821:LEU:HD12	1.77	0.66
1:A:58:LYS:HB3	1:A:60:TYR:HE1	1.59	0.66
1:A:111:ASP:HB3	5:J:815:ARG:HB2	1.78	0.66
3:G:129:LYS:HD3	3:G:129:LYS:H	1.61	0.65
3:G:130:SER:HA	4:H:116:PHE:HB3	1.82	0.62
2:B:109:ARG:O	4:H:108:ARG:N	2.34	0.60
1:C:111:ASP:HB3	5:I:815:ARG:HB2	1.82	0.60
1:C:83:MET:HE3	1:C:86:LEU:HD21	1.83	0.60
2:D:105:ARG:NH2	4:F:173:TYR:OH	2.34	0.60
3:G:119:PRO:HB3	3:G:145:TYR:HB3	1.82	0.60
1:A:12:VAL:HG11	1:A:18:LEU:HB2	1.84	0.60
3:G:188:SER:HB2	3:G:192:GLN:HG2	1.83	0.59
4:H:163:VAL:HG22	4:H:175:LEU:HD12	1.83	0.59
5:J:814:LYS:O	5:J:818:ILE:HD12	2.03	0.58
2:D:1:ASP:O	4:H:189:HIS:NE2	2.33	0.57
1:A:13:GLN:HG3	1:A:14:PRO:HD2	1.88	0.56
4:F:170:ASP:OD1	4:F:170:ASP:N	2.32	0.56
4:H:201:LEU:HD13	4:H:205:VAL:HG13	1.86	0.56
2:D:18:ARG:HG3	2:D:76:SER:HA	1.88	0.55
1:C:51:ILE:HG12	1:C:72:ARG:HD2	1.89	0.55
1:A:102:PRO:HD3	2:B:91:ALA:HB1	1.88	0.55
2:D:11:MET:HE3	2:D:106:LEU:HD13	1.90	0.54
4:F:163:VAL:HG22	4:F:175:LEU:HD12	1.91	0.53
1:C:102:PRO:HD3	2:D:91:ALA:HB1	1.90	0.52
1:A:14:PRO:HD2	1:A:126:SER:HB3	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:193:THR:HG23	3:E:210:LYS:HE2	1.90	0.52
1:A:93:VAL:HG22	1:A:121:MET:HG2	1.92	0.52
1:A:58:LYS:HB3	1:A:60:TYR:CE1	2.42	0.51
2:D:30:SER:OG	2:D:31:LYS:N	2.41	0.51
3:E:200:HIS:CD2	3:E:202:PRO:HD2	2.46	0.50
3:E:129:LYS:HG3	3:E:130:SER:N	2.27	0.50
5:J:815:ARG:O	5:J:819:GLU:HG3	2.12	0.49
2:B:35:TRP:CE2	2:B:73:LEU:HB2	2.47	0.49
1:A:68:PHE:CZ	1:A:83:MET:HG2	2.47	0.49
3:E:188:SER:HB2	3:E:192:GLN:HE21	1.76	0.49
3:E:129:LYS:HG3	3:E:130:SER:H	1.77	0.49
4:H:133:VAL:HG22	4:H:178:THR:HG23	1.94	0.48
1:C:99:LEU:O	5:I:815:ARG:NH2	2.47	0.48
1:A:89:GLU:H	1:A:89:GLU:CD	2.15	0.48
1:A:18:LEU:HD12	1:A:19:ARG:N	2.28	0.48
3:G:134:GLY:HA2	3:G:186:SER:OG	2.14	0.48
1:C:51:ILE:HD12	1:C:58:LYS:HG3	1.95	0.47
5:J:818:ILE:HA	5:J:821:LEU:HD12	1.97	0.47
1:A:99:LEU:O	5:J:815:ARG:NH2	2.48	0.47
2:D:1:ASP:HB3	4:H:189:HIS:CD2	2.50	0.47
2:D:35:TRP:CZ3	2:D:88:CYS:HB3	2.50	0.46
4:F:136:LEU:HD22	4:F:175:LEU:HD22	1.97	0.46
3:G:126:PRO:HG3	3:G:138:LEU:HD23	1.97	0.46
4:F:145:LYS:HB2	4:F:145:LYS:HE2	1.77	0.46
1:C:107:PRO:O	2:D:49:TYR:OH	2.31	0.46
2:B:61:ARG:HB2	2:B:76:SER:HB2	1.98	0.45
4:F:108:ARG:HD3	4:F:109:THR:O	2.17	0.45
4:H:122:ASP:O	4:H:125:LEU:N	2.50	0.45
1:C:47:TRP:HZ2	1:C:50:VAL:HG12	1.82	0.45
2:B:20:THR:HG23	2:B:72:THR:HG23	1.99	0.44
1:A:31:GLY:O	5:J:816:SER:OG	2.32	0.44
2:D:20:THR:HG23	2:D:72:THR:HG23	1.99	0.44
5:I:814:LYS:O	5:I:818:ILE:HD12	2.17	0.44
4:F:147:GLN:NE2	4:F:154:LEU:HG	2.33	0.43
1:A:87:ARG:HB3	1:A:89:GLU:OE2	2.18	0.43
2:B:23:CYS:HG	2:B:88:CYS:HG	1.65	0.43
1:A:11:VAL:HG22	1:A:123:SER:HB2	2.01	0.43
1:C:107:PRO:HB3	2:D:54:LEU:H	1.82	0.43
1:C:111:ASP:O	5:I:815:ARG:HG3	2.19	0.43
1:A:18:LEU:HD12	1:A:19:ARG:H	1.84	0.43
2:D:33:LEU:HD13	2:D:71:PHE:CG	2.54	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:206:LYS:HA	3:E:206:LYS:HD2	1.64	0.43
2:D:1:ASP:HB2	4:H:185:ASP:OD2	2.20	0.42
4:H:108:ARG:NH1	4:H:109:THR:OG1	2.53	0.42
4:H:142:ARG:HH11	4:H:142:ARG:HG2	1.84	0.42
3:G:153:SER:HB2	3:G:197:ASN:HB2	2.02	0.42
3:G:189:LEU:HA	3:G:189:LEU:HD12	1.84	0.42
2:D:1:ASP:HB3	4:H:189:HIS:HE2	1.85	0.41
3:G:197:ASN:ND2	3:G:208:ASP:OD1	2.53	0.41
1:A:60:TYR:OH	1:A:70:ILE:HG22	2.20	0.41
1:C:53:ARG:NH1	5:I:820:ASP:OD1	2.53	0.41
4:F:140:TYR:CG	4:F:141:PRO:HA	2.56	0.41
3:E:189:LEU:HD23	3:E:189:LEU:HA	1.83	0.41
1:C:13:GLN:HB2	1:C:16:ARG:HG3	2.02	0.41
2:D:3:GLN:OE1	4:H:155:GLN:NE2	2.54	0.41
1:A:51:ILE:HG23	1:A:72:ARG:HH11	1.86	0.41
4:H:183:LYS:O	4:H:187:GLU:HG3	2.20	0.41
5:J:814:LYS:HG2	5:J:818:ILE:HD11	2.02	0.41
1:C:60:TYR:CE1	1:C:70:ILE:HG22	2.55	0.41
4:H:128:GLY:HA2	4:H:183:LYS:HB2	2.03	0.41
2:B:107:GLU:OE2	4:H:140:TYR:OH	2.26	0.40
4:H:142:ARG:HG2	4:H:142:ARG:NH1	2.36	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	124/126 (98%)	119 (96%)	5 (4%)	0	100 100
1	C	124/126 (98%)	120 (97%)	4 (3%)	0	100 100
2	B	107/109 (98%)	98 (92%)	9 (8%)	0	100 100
2	D	107/109 (98%)	100 (94%)	7 (6%)	0	100 100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	101/103 (98%)	94 (93%)	7 (7%)	0	100	100
3	G	101/103 (98%)	95 (94%)	6 (6%)	0	100	100
4	F	104/107 (97%)	103 (99%)	1 (1%)	0	100	100
4	H	104/107 (97%)	102 (98%)	2 (2%)	0	100	100
5	I	10/15 (67%)	10 (100%)	0	0	100	100
5	J	10/15 (67%)	10 (100%)	0	0	100	100
All	All	892/920 (97%)	851 (95%)	41 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/103 (100%)	101 (98%)	2 (2%)	57	85
1	C	103/103 (100%)	99 (96%)	4 (4%)	32	66
2	B	93/93 (100%)	91 (98%)	2 (2%)	52	83
2	D	93/93 (100%)	91 (98%)	2 (2%)	52	83
3	E	88/89 (99%)	86 (98%)	2 (2%)	50	82
3	G	88/89 (99%)	84 (96%)	4 (4%)	27	60
4	F	95/96 (99%)	90 (95%)	5 (5%)	22	54
4	H	95/96 (99%)	89 (94%)	6 (6%)	18	46
5	I	12/15 (80%)	11 (92%)	1 (8%)	11	32
5	J	12/15 (80%)	12 (100%)	0	100	100
All	All	782/792 (99%)	754 (96%)	28 (4%)	35	69

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	58	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	71	SER
1	C	82	GLU
1	C	96	CYS
2	D	18	ARG
2	D	60	SER
3	E	132	SER
3	E	197	ASN
4	F	108	ARG
4	F	114	SER
4	F	132	VAL
4	F	145	LYS
4	F	170	ASP
5	I	815	ARG
1	A	13	GLN
1	A	96	CYS
2	B	60	SER
2	B	89	GLN
3	G	129	LYS
3	G	200	HIS
3	G	206	LYS
3	G	208	ASP
4	H	108	ARG
4	H	121	SER
4	H	156	SER
4	H	159	SER
4	H	185	ASP
4	H	188	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	E	192	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	126/126 (100%)	0.36	7 (5%)	24	16	45, 79, 109, 125	0
1	C	126/126 (100%)	0.18	3 (2%)	59	49	36, 62, 88, 105	0
2	B	109/109 (100%)	0.07	1 (0%)	84	80	44, 62, 83, 98	0
2	D	109/109 (100%)	-0.09	1 (0%)	84	80	42, 63, 79, 86	0
3	E	103/103 (100%)	0.19	4 (3%)	39	29	35, 46, 91, 131	0
3	G	103/103 (100%)	0.56	5 (4%)	29	20	54, 80, 104, 138	0
4	F	106/107 (99%)	-0.15	0	100	100	37, 46, 65, 76	0
4	H	106/107 (99%)	-0.02	0	100	100	44, 57, 77, 93	0
5	I	12/15 (80%)	0.64	2 (16%)	1	1	81, 93, 98, 107	0
5	J	12/15 (80%)	1.11	3 (25%)	0	0	92, 105, 119, 122	0
All	All	912/920 (99%)	0.16	26 (2%)	51	41	35, 63, 102, 138	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	216	CYS	8.0
3	E	131	THR	5.5
3	E	130	SER	5.4
3	G	131	THR	4.7
3	G	130	SER	4.4
1	A	1	GLN	3.9
1	A	30	SER	3.8
3	E	132	SER	3.6
3	G	211	VAL	3.4
3	G	132	SER	3.3
1	A	29	PHE	3.3
1	C	118	GLN	2.9
3	E	114	ALA	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	24	ALA	2.7
5	J	815	ARG	2.6
1	C	1	GLN	2.4
2	B	1	ASP	2.4
1	A	82	GLU	2.3
1	A	19	ARG	2.3
2	D	54	LEU	2.3
5	I	822	LEU	2.2
5	J	816	SER	2.2
1	A	56	ARG	2.2
1	C	26	GLY	2.2
5	I	813	SER	2.2
5	J	823	PHE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.