

# Full wwPDB X-ray Structure Validation Report (i)

#### Jan 22, 2024 – 04:08 PM EST

PDB ID : 8FVV

Title: Rubrerythrin from B. pseudomallei: iron-bound

Authors: Monteiro, D.C.F.; Snell, M.E.; Budziszewski, G.R.; Bowman, S.E.J.

Deposited on : 2023-01-19

Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467 Xtriage (Phenix): 1.13

EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove) roteins) : Engh & Huber (2001)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	169	78%	5%	18%
1	В	169	73%	9%	18%
1	С	169	73%	8% •	18%
1	D	169	74%	8%	18%
1	Е	169	77%	5%	18%



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Length	Quality of chain		
	_		2%		
1	F	169	78%		18%



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 13347 atoms, of which 6068 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Rubrerythrin.

Mol	Chain	Residues			Atom	ıS			ZeroOcc	AltConf	Trace
1	A	139	Total	С	Н	N	О	S	35	0	0
1	Λ	139	2082	666	1008	188	218	2	30		
1	В	139	Total	С	Н	N	О	S	34	0	0
1	Ъ	139	2090	668	1012	189	219	2	34	U	U
1	С	139	Total	С	Н	N	О	S	34	0	0
1		109	2090	668	1012	189	219	2	94	U	
1	D	139	Total	С	Н	N	O	S	34	0	0
1	D	109	2090	668	1012	189	219	2	94	0	U
1	E	139	Total	С	Н	N	O	S	34	0	0
1	Ľ	109	2090	668	1012	189	219	2	54		U
1	F	139	Total	С	Н	N	О	S	34	0	0
1	I'	109	2090	668	1012	189	219	2	54	U	U

There are 174 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-28	MET	-	initiating methionine	UNP Q3JK21
A	-27	GLY	-	expression tag	UNP Q3JK21
A	-26	SER	-	expression tag	UNP Q3JK21
A	-25	SER	-	expression tag	UNP Q3JK21
A	-24	HIS	-	expression tag	UNP Q3JK21
A	-23	HIS	-	expression tag	UNP Q3JK21
A	-22	HIS	-	expression tag	UNP Q3JK21
A	-21	HIS	-	expression tag	UNP Q3JK21
A	-20	HIS	-	expression tag	UNP Q3JK21
A	-19	HIS	-	expression tag	UNP Q3JK21
A	-18	SER	-	expression tag	UNP Q3JK21
A	-17	SER	_	expression tag	UNP Q3JK21
A	-16	GLY	-	expression tag	UNP Q3JK21
A	-15	LEU		expression tag	UNP Q3JK21
A	-14	VAL	-	expression tag	UNP Q3JK21
A	-13	PRO	-	expression tag	UNP Q3JK21
A	-12	ARG	-	expression tag	UNP Q3JK21



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	GLY	-	expression tag	UNP Q3JK21
A	-10	SER	-	expression tag	UNP Q3JK21
A	-9	HIS	-	expression tag	UNP Q3JK21
A	-8	MET	-	expression tag	UNP Q3JK21
A	-7	GLU	-	expression tag	UNP Q3JK21
A	-6	ASN	-	expression tag	UNP Q3JK21
A	-5	LEU	-	expression tag	UNP Q3JK21
A	-4	TYR	-	expression tag	UNP Q3JK21
A	-3	PHE	-	expression tag	UNP Q3JK21
A	-2	GLN	-	expression tag	UNP Q3JK21
A	-1	GLY	-	expression tag	UNP Q3JK21
A	0	SER	-	expression tag	UNP Q3JK21
В	-28	MET	-	initiating methionine	UNP Q3JK21
В	-27	GLY	-	expression tag	UNP Q3JK21
В	-26	SER	-	expression tag	UNP Q3JK21
В	-25	SER	-	expression tag	UNP Q3JK21
В	-24	HIS	-	expression tag	UNP Q3JK21
В	-23	HIS	-	expression tag	UNP Q3JK21
В	-22	HIS	-	expression tag	UNP Q3JK21
В	-21	HIS	-	expression tag	UNP Q3JK21
В	-20	HIS	-	expression tag	UNP Q3JK21
В	-19	HIS	-	expression tag	UNP Q3JK21
В	-18	SER	-	expression tag	UNP Q3JK21
В	-17	SER	-	expression tag	UNP Q3JK21
В	-16	GLY	-	expression tag	UNP Q3JK21
В	-15	LEU	-	expression tag	UNP Q3JK21
В	-14	VAL	-	expression tag	UNP Q3JK21
В	-13	PRO	-	expression tag	UNP Q3JK21
В	-12	ARG	-	expression tag	UNP Q3JK21
В	-11	GLY	-	expression tag	UNP Q3JK21
В	-10	SER	-	expression tag	UNP Q3JK21
В	-9	HIS	-	expression tag	UNP Q3JK21
В	-8	MET	-	expression tag	UNP Q3JK21
В	-7	GLU		expression tag	UNP Q3JK21
В	-6	ASN	-	expression tag	UNP Q3JK21
В	-5	LEU	-	expression tag	UNP Q3JK21
В	-4	TYR	-	expression tag	UNP Q3JK21
В	-3	PHE	-	expression tag	UNP Q3JK21
В	-2	GLN		expression tag	UNP Q3JK21
В	-1	GLY	-	expression tag	UNP Q3JK21
В	0	SER	-	expression tag	UNP Q3JK21
С	-28	MET	-	initiating methionine	UNP Q3JK21



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
С	-27	GLY	-	expression tag	UNP Q3JK21
С	-26	SER	-	expression tag	UNP Q3JK21
С	-25	SER	-	expression tag	UNP Q3JK21
С	-24	HIS	-	expression tag	UNP Q3JK21
С	-23	HIS	-	expression tag	UNP Q3JK21
С	-22	HIS	-	expression tag	UNP Q3JK21
С	-21	HIS	-	expression tag	UNP Q3JK21
С	-20	HIS	-	expression tag	UNP Q3JK21
С	-19	HIS	-	expression tag	UNP Q3JK21
С	-18	SER	-	expression tag	UNP Q3JK21
С	-17	SER	-	expression tag	UNP Q3JK21
С	-16	GLY	-	expression tag	UNP Q3JK21
С	-15	LEU	-	expression tag	UNP Q3JK21
С	-14	VAL	-	expression tag	UNP Q3JK21
С	-13	PRO	-	expression tag	UNP Q3JK21
С	-12	ARG	-	expression tag	UNP Q3JK21
С	-11	GLY	-	expression tag	UNP Q3JK21
С	-10	SER	-	expression tag	UNP Q3JK21
С	-9	HIS	-	expression tag	UNP Q3JK21
С	-8	MET	-	expression tag	UNP Q3JK21
С	-7	GLU	-	expression tag	UNP Q3JK21
С	-6	ASN	-	expression tag	UNP Q3JK21
С	-5	LEU	-	expression tag	UNP Q3JK21
С	-4	TYR	-	expression tag	UNP Q3JK21
С	-3	PHE	-	expression tag	UNP Q3JK21
С	-2	GLN	_	expression tag	UNP Q3JK21
С	-1	GLY	-	expression tag	UNP Q3JK21
С	0	SER	-	expression tag	UNP Q3JK21
D	-28	MET	-	initiating methionine	UNP Q3JK21
D	-27	GLY	-	expression tag	UNP Q3JK21
D	-26	SER	-	expression tag	UNP Q3JK21
D	-25	SER	-	expression tag	UNP Q3JK21
D	-24	HIS	-	expression tag	UNP Q3JK21
D	-23	HIS	-	expression tag	UNP Q3JK21
D	-22	HIS	-	expression tag	UNP Q3JK21
D	-21	HIS	-	expression tag	UNP Q3JK21
D	-20	HIS	-	expression tag	UNP Q3JK21
D	-19	HIS	-	expression tag	UNP Q3JK21
D	-18	SER	-	expression tag	UNP Q3JK21
D	-17	SER	-	expression tag	UNP Q3JK21
D	-16	GLY	-	expression tag	UNP Q3JK21
D	-15	LEU	-	expression tag	UNP Q3JK21



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
D	-14	VAL	-	expression tag	UNP Q3JK21
D	-13	PRO	-	expression tag	UNP Q3JK21
D	-12	ARG	-	expression tag	UNP Q3JK21
D	-11	GLY	-	expression tag	UNP Q3JK21
D	-10	SER	-	expression tag	UNP Q3JK21
D	-9	HIS	-	expression tag	UNP Q3JK21
D	-8	MET	-	expression tag	UNP Q3JK21
D	-7	GLU	-	expression tag	UNP Q3JK21
D	-6	ASN	-	expression tag	UNP Q3JK21
D	-5	LEU	-	expression tag	UNP Q3JK21
D	-4	TYR	-	expression tag	UNP Q3JK21
D	-3	PHE	-	expression tag	UNP Q3JK21
D	-2	GLN	-	expression tag	UNP Q3JK21
D	-1	GLY	-	expression tag	UNP Q3JK21
D	0	SER	-	expression tag	UNP Q3JK21
Е	-28	MET	-	initiating methionine	UNP Q3JK21
Е	-27	GLY	-	expression tag	UNP Q3JK21
Е	-26	SER	-	expression tag	UNP Q3JK21
Е	-25	SER	-	expression tag	UNP Q3JK21
Е	-24	HIS	-	expression tag	UNP Q3JK21
Е	-23	HIS	-	expression tag	UNP Q3JK21
Е	-22	HIS	-	expression tag	UNP Q3JK21
Е	-21	HIS	-	expression tag	UNP Q3JK21
Е	-20	HIS	-	expression tag	UNP Q3JK21
Е	-19	HIS	-	expression tag	UNP Q3JK21
Е	-18	SER	-	expression tag	UNP Q3JK21
E	-17	SER	-	expression tag	UNP Q3JK21
E	-16	GLY	-	expression tag	UNP Q3JK21
Е	-15	LEU	_	expression tag	UNP Q3JK21
E	-14	VAL	-	expression tag	UNP Q3JK21
Е	-13	PRO	-	expression tag	UNP Q3JK21
E	-12	ARG	_	expression tag	UNP Q3JK21
E	-11	GLY	-	expression tag	UNP Q3JK21
Е	-10	SER	-	expression tag	UNP Q3JK21
Е	-9	HIS	-	expression tag	UNP Q3JK21
Е	-8	MET	-	expression tag	UNP Q3JK21
Е	-7	GLU	-	expression tag	UNP Q3JK21
Е	-6	ASN	-	expression tag	UNP Q3JK21
Е	-5	LEU	-	expression tag	UNP Q3JK21
Е	-4	TYR	-	expression tag	UNP Q3JK21
Е	-3	PHE	-	expression tag	UNP Q3JK21
Е	-2	GLN	-	expression tag	UNP Q3JK21



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-1	GLY	-	expression tag	UNP Q3JK21
E	0	SER	-	expression tag	UNP Q3JK21
F	-28	MET	_	initiating methionine	UNP Q3JK21
F	-27	GLY	_	expression tag	UNP Q3JK21
F	-26	SER	-	expression tag	UNP Q3JK21
F	-25	SER	-	expression tag	UNP Q3JK21
F	-24	HIS	_	expression tag	UNP Q3JK21
F	-23	HIS	-	expression tag	UNP Q3JK21
F	-22	HIS	-	expression tag	UNP Q3JK21
F	-21	HIS	-	expression tag	UNP Q3JK21
F	-20	HIS	-	expression tag	UNP Q3JK21
F	-19	HIS	-	expression tag	UNP Q3JK21
F	-18	SER	_	expression tag	UNP Q3JK21
F	-17	SER	-	expression tag	UNP Q3JK21
F	-16	GLY	-	expression tag	UNP Q3JK21
F	-15	LEU	-	expression tag	UNP Q3JK21
F	-14	VAL	-	expression tag	UNP Q3JK21
F	-13	PRO	_	expression tag	UNP Q3JK21
F	-12	ARG	-	expression tag	UNP Q3JK21
F	-11	GLY	_	expression tag	UNP Q3JK21
F	-10	SER	-	expression tag	UNP Q3JK21
F	-9	HIS	-	expression tag	UNP Q3JK21
F	-8	MET	-	expression tag	UNP Q3JK21
F	-7	GLU	-	expression tag	UNP Q3JK21
F	-6	ASN	-	expression tag	UNP Q3JK21
F	-5	LEU	-	expression tag	UNP Q3JK21
F	-4	TYR	-	expression tag	UNP Q3JK21
F	-3	PHE	-	expression tag	UNP Q3JK21
F	-2	GLN	-	expression tag	UNP Q3JK21
F	-1	GLY	-	expression tag	UNP Q3JK21
F	0	SER	-	expression tag	UNP Q3JK21

 $\bullet$  Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Fe 2 2	0	0
2	В	2	Total Fe 2 2	0	0
2	С	2	Total Fe 2 2	0	0
2	D	2	Total Fe 2 2	0	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	E	2	Total Fe 2 2	0	0
2	F	2	Total Fe 2 2	0	0

#### • Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	143	Total O 145 145	0	2
3	В	139	Total O 139 139	0	0
3	С	138	Total O 138 138	0	0
3	D	121	Total O 122 122	0	1
3	E	136	Total O 136 136	0	0
3	F	123	Total O 123 123	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rubrerythrin Chain A: 18% METY SER MET • Molecule 1: Rubrerythrin Chain B: 73% • Molecule 1: Rubrerythrin Chain C: 73% 18% • Molecule 1: Rubrerythrin Chain D: 74% 18% 







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants	202.17Å 202.17Å 69.83Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	29.90 - 1.93	Depositor
resolution (A)	29.90 - 1.93	EDS
% Data completeness	92.4 (29.90-1.93)	Depositor
(in resolution range)	76.5 (29.90 - 1.93)	EDS
$R_{merge}$	0.23	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.39  (at  1.93Å)	Xtriage
Refinement program	REFMAC 5.8.0403	Depositor
$R, R_{free}$	0.161 , $0.209$	Depositor
it, it <sub>free</sub>	0.171 , $0.214$	DCC
$R_{free}$ test set	2998 reflections $(4.91\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.9	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.40  ,  49.4	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.013 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	13347	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.50	0/1096	0.69	0/1478	
1	В	0.49	0/1100	0.72	0/1483	
1	С	0.48	0/1100	0.74	0/1483	
1	D	0.48	0/1100	0.69	0/1483	
1	Е	0.44	0/1100	0.67	0/1483	
1	F	0.42	0/1100	0.67	0/1483	
All	All	0.47	0/6596	0.70	0/8893	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1074	1008	1000	6	0
1	В	1078	1012	1006	9	1
1	С	1078	1012	1006	8	1
1	D	1078	1012	1006	6	1
1	Е	1078	1012	1006	7	0
1	F	1078	1012	1006	4	1
2	A	2	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	2	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
2	Е	2	0	0	0	0
2	F	2	0	0	0	0
3	A	145	0	0	2	0
3	В	139	0	0	0	0
3	С	138	0	0	2	0
3	D	122	0	0	1	0
3	Ε	136	0	0	2	0
3	F	123	0	0	1	0
All	All	7279	6068	6030	33	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:F:3:GLN:NE2	3:F:301:HOH:O	2.17	0.76
1:A:90:GLU:OE1	1:A:124:GLU:OE1	2.13	0.67
1:B:90:GLU:OE1	1:B:124:GLU:OE1	2.13	0.67
1:D:84:GLN:NE2	3:D:301:HOH:O	2.28	0.66
1:C:115:ASN:HB3	3:C:403:HOH:O	1.96	0.65
1:C:90:GLU:OE1	1:C:124:GLU:OE1	2.23	0.57
1:C:17:PHE:CE1	1:C:61:LEU:HG	2.39	0.57
1:A:120:LEU:HD22	1:B:56:HIS:CD2	2.45	0.51
1:B:94:TYR:CE1	1:B:125:ARG:HG3	2.46	0.51
1:A:56:HIS:CD2	1:B:120:LEU:HD22	2.46	0.50
1:A:107:ASP:OD2	3:A:301:HOH:O	2.18	0.50
1:B:3:GLN:HA	1:B:3:GLN:OE1	2.12	0.50
1:E:124:GLU:HA	1:E:124:GLU:OE1	2.12	0.49
1:B:124:GLU:OE1	1:B:124:GLU:HA	2.13	0.49
1:A:106:ARG:HD2	3:A:315:HOH:O	2.11	0.48
1:B:80:ARG:HH21	1:B:140:ASP:CG	2.16	0.48
1:C:133:LYS:HE2	3:C:417:HOH:O	2.13	0.48
1:C:120:LEU:HD22	1:D:56:HIS:CE1	2.48	0.48
1:D:90:GLU:OE1	1:D:124:GLU:OE1	2.32	0.47
1:E:86:ALA:O	1:E:90:GLU:HG2	2.14	0.47
1:E:139:VAL:HG13	3:E:362:HOH:O	2.14	0.47
1:E:106:ARG:HD2	3:E:301:HOH:O	2.16	0.44



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance} ({ m \AA})$	overlap (Å)
1:C:94:TYR:CD2	1:C:125:ARG:HA	2.53	0.43
1:E:42:ILE:HD12	1:F:83:LEU:HD11	1.99	0.43
1:B:8:LYS:HD2	1:B:8:LYS:HA	1.85	0.43
1:C:116:TRP:CZ2	1:D:59:GLY:HA3	2.54	0.42
1:E:59:GLY:HA3	1:F:116:TRP:CZ2	2.54	0.41
1:E:90:GLU:OE2	1:E:124:GLU:OE1	2.38	0.41
1:C:65:GLU:HG2	1:C:76:PHE:CE2	2.56	0.41
1:D:17:PHE:CE1	1:D:61:LEU:HG	2.55	0.41
1:D:65:GLU:HG2	1:D:76:PHE:CE2	2.55	0.41
1:A:30:PHE:CG	1:B:82:ASN:HB3	2.57	0.40
1:F:91:THR:HA	1:F:94:TYR:CE2	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:49:THR:HG1	1:F:40:ASN:HD21[5_555]	1.23	0.37
1:B:40:ASN:HD21	1:D:49:THR:HG1[8_544]	1.29	0.31

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	$137/169\ (81\%)$	134 (98%)	3 (2%)	0	100 100
1	В	137/169~(81%)	133 (97%)	4 (3%)	0	100 100
1	C	$137/169\ (81\%)$	133 (97%)	4 (3%)	0	100 100
1	D	137/169~(81%)	134 (98%)	3 (2%)	0	100 100
1	E	$137/169\ (81\%)$	134 (98%)	3 (2%)	0	100 100
1	F	137/169 (81%)	132 (96%)	5 (4%)	0	100 100
All	All	822/1014 (81%)	800 (97%)	22 (3%)	0	100 100



There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	104/131 (79%)	103 (99%)	1 (1%)	76 71
1	В	105/131 (80%)	102 (97%)	3 (3%)	42 28
1	С	105/131 (80%)	102 (97%)	3 (3%)	42 28
1	D	105/131 (80%)	101 (96%)	4 (4%)	33 18
1	E	105/131 (80%)	103 (98%)	2 (2%)	57 45
1	F	105/131 (80%)	103 (98%)	2 (2%)	57 45
All	All	629/786 (80%)	614 (98%)	15 (2%)	49 36

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	ARG
1	В	25	ARG
1	В	96	ASP
1	В	122	LYS
1	С	8	LYS
1	С	25	ARG
1	С	61	LEU
1	D	5	LYS
1	D	11	GLU
1	D	25	ARG
1	D	136	ASP
1	Е	25	ARG
1	Е	94	TYR
1	F	3	GLN
1	F	25	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	129	ASN
1	F	84	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	A	139/169 (82%)	-0.35	1 (0%) 87 91	14, 22, 39, 67	0
1	В	139/169 (82%)	-0.32	3 (2%) 62 69	14, 22, 43, 67	0
1	С	139/169 (82%)	-0.08	4 (2%) 51 59	16, 22, 43, 82	0
1	D	139/169 (82%)	-0.08	5 (3%) 42 50	15, 27, 47, 71	0
1	E	139/169 (82%)	-0.06	3 (2%) 62 69	17, 25, 46, 84	0
1	F	139/169 (82%)	-0.20	3 (2%) 62 69	17, 27, 47, 82	0
All	All	834/1014 (82%)	-0.18	19 (2%) 60 67	14, 24, 45, 84	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	2	ALA	8.7
1	Е	2	ALA	7.6
1	A	2	ALA	5.6
1	D	2	ALA	5.5
1	F	2	ALA	5.0
1	В	2	ALA	4.8
1	С	3	GLN	4.0
1	D	3	GLN	3.7
1	Е	3	GLN	3.6
1	В	3	GLN	2.7
1	F	8	LYS	2.4
1	Е	66	ALA	2.3
1	D	66	ALA	2.2
1	В	107	ASP	2.2
1	D	136	ASP	2.2
1	С	66	ALA	2.1
1	F	109	GLY	2.1
1	С	110	PHE	2.1
1	D	103	LYS	2.1



### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q < 0.9
2	FE	F	202	1/1	0.90	0.15	88,88,88,88	1
2	FE	В	202	1/1	0.94	0.08	59,59,59,59	1
2	FE	Е	202	1/1	0.96	0.24	66,66,66,66	1
2	FE	D	202	1/1	0.97	0.09	57,57,57,57	1
2	FE	С	202	1/1	0.98	0.16	36,36,36,36	1
2	FE	D	201	1/1	0.99	0.10	27,27,27,27	1
2	FE	В	201	1/1	0.99	0.08	34,34,34,34	0
2	FE	Е	201	1/1	0.99	0.07	23,23,23,23	1
2	FE	A	201	1/1	0.99	0.09	21,21,21,21	1
2	FE	F	201	1/1	0.99	0.10	31,31,31,31	0
2	FE	A	202	1/1	0.99	0.07	30,30,30,30	1
2	FE	С	201	1/1	1.00	0.06	21,21,21,21	1

## 6.5 Other polymers (i)

There are no such residues in this entry.

