



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 9, 2024 – 02:07 PM JST

PDB ID : 8IYA
Title : Complex of SETDB1-derived peptide bound to UBE2E1
Authors : Du, Y.X.; Liu, L.
Deposited on : 2023-04-04
Resolution : 2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	<div> <div>79%</div> <div>14%</div> <div>• 5%</div> </div>
1	B	170	<div> <div>%</div> <div>76%</div> <div>16%</div> <div>• 6%</div> </div>
1	C	170	<div> <div>2%</div> <div>74%</div> <div>16%</div> <div>11%</div> </div>
2	D	6	<div> <div>100%</div> </div>
2	E	6	<div> <div>83%</div> <div>17%</div> </div>
2	F	6	<div> <div>100%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin-conjugating enzyme E2 E1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	160	Total	C	N	O	S	0	0	0
			1259	807	213	236	3			
1	A	161	Total	C	N	O	S	0	0	0
			1268	813	214	238	3			
1	C	152	Total	C	N	O	S	0	0	0
			1197	766	203	225	3			

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	24	MET	-	initiating methionine	UNP P51965
B	25	HIS	-	expression tag	UNP P51965
B	26	HIS	-	expression tag	UNP P51965
B	27	HIS	-	expression tag	UNP P51965
B	28	HIS	-	expression tag	UNP P51965
B	29	HIS	-	expression tag	UNP P51965
B	30	HIS	-	expression tag	UNP P51965
B	31	HIS	-	expression tag	UNP P51965
B	32	HIS	-	expression tag	UNP P51965
B	33	LEU	-	expression tag	UNP P51965
B	34	GLU	-	expression tag	UNP P51965
B	35	VAL	-	expression tag	UNP P51965
B	36	LEU	-	expression tag	UNP P51965
B	37	PHE	-	expression tag	UNP P51965
B	38	GLN	-	expression tag	UNP P51965
B	39	GLY	-	expression tag	UNP P51965
B	40	PRO	-	expression tag	UNP P51965
B	67	SER	CYS	engineered mutation	UNP P51965
B	122	PRO	CYS	engineered mutation	UNP P51965
B	153	SER	CYS	engineered mutation	UNP P51965
B	159	PRO	CYS	engineered mutation	UNP P51965
A	24	MET	-	initiating methionine	UNP P51965
A	25	HIS	-	expression tag	UNP P51965

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	HIS	-	expression tag	UNP P51965
A	27	HIS	-	expression tag	UNP P51965
A	28	HIS	-	expression tag	UNP P51965
A	29	HIS	-	expression tag	UNP P51965
A	30	HIS	-	expression tag	UNP P51965
A	31	HIS	-	expression tag	UNP P51965
A	32	HIS	-	expression tag	UNP P51965
A	33	LEU	-	expression tag	UNP P51965
A	34	GLU	-	expression tag	UNP P51965
A	35	VAL	-	expression tag	UNP P51965
A	36	LEU	-	expression tag	UNP P51965
A	37	PHE	-	expression tag	UNP P51965
A	38	GLN	-	expression tag	UNP P51965
A	39	GLY	-	expression tag	UNP P51965
A	40	PRO	-	expression tag	UNP P51965
A	67	SER	CYS	engineered mutation	UNP P51965
A	122	PRO	CYS	engineered mutation	UNP P51965
A	153	SER	CYS	engineered mutation	UNP P51965
A	159	PRO	CYS	engineered mutation	UNP P51965
C	24	MET	-	initiating methionine	UNP P51965
C	25	HIS	-	expression tag	UNP P51965
C	26	HIS	-	expression tag	UNP P51965
C	27	HIS	-	expression tag	UNP P51965
C	28	HIS	-	expression tag	UNP P51965
C	29	HIS	-	expression tag	UNP P51965
C	30	HIS	-	expression tag	UNP P51965
C	31	HIS	-	expression tag	UNP P51965
C	32	HIS	-	expression tag	UNP P51965
C	33	LEU	-	expression tag	UNP P51965
C	34	GLU	-	expression tag	UNP P51965
C	35	VAL	-	expression tag	UNP P51965
C	36	LEU	-	expression tag	UNP P51965
C	37	PHE	-	expression tag	UNP P51965
C	38	GLN	-	expression tag	UNP P51965
C	39	GLY	-	expression tag	UNP P51965
C	40	PRO	-	expression tag	UNP P51965
C	67	SER	CYS	engineered mutation	UNP P51965
C	122	PRO	CYS	engineered mutation	UNP P51965
C	153	SER	CYS	engineered mutation	UNP P51965
C	159	PRO	CYS	engineered mutation	UNP P51965


- Molecule 2 is a protein called Histone-lysine N-methyltransferase SETDB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	6	Total	C	N	O	S	0	0	0
			46	27	6	12	1			
2	E	6	Total	C	N	O	S	0	0	0
			46	27	6	12	1			
2	F	6	Total	C	N	O	S	0	0	0
			46	27	6	12	1			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	CYS	LYS	conflict	UNP Q15047
E	1	CYS	LYS	conflict	UNP Q15047
F	1	CYS	LYS	conflict	UNP Q15047

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	14	Total 14	O 14	0	0
4	A	11	Total 11	O 11	0	0
4	C	9	Total 9	O 9	0	0
4	E	2	Total 2	O 2	0	0

- Molecule 1: Ubiquitin-conjugating enzyme E2 E1





- Molecule 2: Histone-lysine N-methyltransferase SETDB1

Chain F:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	102.75Å 50.08Å 109.25Å 90.00° 91.75° 90.00°	Depositor
Resolution (Å)	41.42 – 2.43 45.01 – 2.43	Depositor EDS
% Data completeness (in resolution range)	99.6 (41.42-2.43) 99.6 (45.01-2.43)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.205 , 0.247 0.204 , 0.246	Depositor DCC
R_{free} test set	1045 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.630	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3913	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1303	0.66	1/1778 (0.1%)
1	B	0.49	0/1294	0.66	0/1767
1	C	0.45	0/1230	0.64	0/1678
2	D	0.83	0/46	0.63	0/60
2	E	0.45	0/46	0.53	0/60
2	F	0.44	0/46	0.77	0/60
All	All	0.49	0/3965	0.65	1/5403 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	33	LEU	CA-CB-CG	6.49	130.22	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1268	0	1270	19	0
1	B	1259	0	1259	20	1
1	C	1197	0	1200	16	0
2	D	46	0	35	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	46	0	35	1	0
2	F	46	0	35	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
4	A	11	0	0	0	0
4	B	14	0	0	0	0
4	C	9	0	0	1	0
4	E	2	0	0	0	0
All	All	3913	0	3834	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:45:LEU:HD23	1:C:104:PRO:HB3	1.16	1.11
1:C:45:LEU:CD2	1:C:104:PRO:HB3	1.99	0.91
1:C:45:LEU:HD23	1:C:104:PRO:CB	2.09	0.71
1:B:179:GLU:OE2	1:B:182:ARG:NH1	2.22	0.71
1:A:172:GLN:NE2	1:A:179:GLU:HG3	2.14	0.63
1:C:176:ASN:ND2	1:C:179:GLU:HB2	2.13	0.63
1:B:136:LYS:HG3	1:B:137:ASP:H	1.61	0.63
1:C:166:VAL:HG11	1:C:169:ILE:HD12	1.81	0.61
1:B:118:ARG:HA	1:B:118:ARG:HE	1.65	0.60
1:A:35:VAL:HG23	1:A:36:LEU:HD12	1.82	0.60
1:A:120:TYR:OH	1:A:172:GLN:OE1	2.20	0.59
1:C:51:ARG:NH2	1:C:143:LEU:O	2.35	0.59
1:C:81:SER:HB3	1:C:98:LEU:HB2	1.86	0.57
1:A:33:LEU:HD13	1:A:34:GLU:OE1	2.05	0.57
1:B:81:SER:HB3	1:B:98:LEU:HB2	1.87	0.56
1:C:137:ASP:OD1	1:C:138:ASN:N	2.37	0.56
1:C:172:GLN:OE1	1:C:180:HIS:HA	2.06	0.54
1:B:37:PHE:CG	1:B:37:PHE:O	2.61	0.54
1:C:126:SER:OG	1:C:127:GLN:NE2	2.38	0.53
1:A:172:GLN:HE21	1:A:179:GLU:HG3	1.75	0.52
1:B:80:ARG:HH12	1:B:116:ARG:CZ	2.22	0.52
1:A:172:GLN:O	1:A:176:ASN:O	2.27	0.52
1:A:120:TYR:CZ	1:A:169:ILE:HG12	2.45	0.51
1:B:172:GLN:HG2	1:B:180:HIS:HB2	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:99:ASP:OD2	1:C:116:ARG:HD2	2.11	0.49
1:B:48:SER:O	1:B:52:ILE:HG13	2.13	0.49
1:A:81:SER:HB3	1:A:98:LEU:HB2	1.94	0.49
1:B:131:CYS:HB2	1:B:165:LEU:HG	1.95	0.48
1:A:66:ASN:HB3	1:A:84:LEU:HB2	1.96	0.48
1:B:136:LYS:CG	1:B:137:ASP:H	2.27	0.47
1:B:142:ALA:HB2	2:E:4:TYR:CG	2.49	0.47
1:A:112:LYS:HD2	1:A:112:LYS:HA	1.73	0.47
1:C:71:PRO:HA	1:C:79:TRP:HA	1.96	0.47
1:C:166:VAL:CG1	1:C:169:ILE:HD12	2.45	0.46
1:C:45:LEU:CD2	1:C:104:PRO:CB	2.81	0.46
1:B:99:ASP:OD2	1:B:116:ARG:HD2	2.16	0.46
1:B:136:LYS:HG3	1:B:137:ASP:N	2.31	0.45
1:B:190:ARG:HG2	1:B:191:TYR:CD2	2.52	0.45
1:B:118:ARG:CZ	1:B:126:SER:HA	2.47	0.45
1:B:118:ARG:NH1	1:B:126:SER:HA	2.32	0.45
1:A:33:LEU:O	1:A:35:VAL:HG22	2.16	0.44
1:B:118:ARG:HA	1:B:118:ARG:NE	2.30	0.44
1:A:133:ASP:OD1	1:A:134:ILE:N	2.51	0.44
1:C:167:GLY:O	1:C:171:THR:HG23	2.18	0.43
1:B:89:SER:HB3	1:B:157:THR:HG22	1.99	0.42
1:A:33:LEU:O	1:A:35:VAL:N	2.47	0.42
1:A:172:GLN:HE21	1:A:179:GLU:CG	2.31	0.42
1:B:48:SER:OG	1:B:77:TYR:OH	2.28	0.42
1:A:80:ARG:HH12	1:A:116:ARG:HE	1.68	0.41
1:A:131:CYS:HB2	1:A:165:LEU:HG	2.03	0.41
1:A:172:GLN:HG2	1:A:180:HIS:HB2	2.01	0.41
1:B:122:PRO:HG2	1:B:159:PRO:HB3	2.03	0.41
1:C:42:SER:OG	4:C:302:HOH:O	2.22	0.41
1:A:186:GLN:O	1:A:190:ARG:HG2	2.20	0.41
1:A:118:ARG:NH1	1:A:126:SER:HA	2.36	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:74:ASP:OD1	1:B:116:ARG:NH2[2_556]	2.04	0.16

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/170 (94%)	155 (98%)	4 (2%)	0	100	100
1	B	158/170 (93%)	155 (98%)	3 (2%)	0	100	100
1	C	150/170 (88%)	148 (99%)	2 (1%)	0	100	100
2	D	4/6 (67%)	4 (100%)	0	0	100	100
2	E	4/6 (67%)	4 (100%)	0	0	100	100
2	F	4/6 (67%)	4 (100%)	0	0	100	100
All	All	479/528 (91%)	470 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	143/152 (94%)	139 (97%)	4 (3%)	43	56
1	B	142/152 (93%)	136 (96%)	6 (4%)	30	39
1	C	135/152 (89%)	131 (97%)	4 (3%)	41	53
2	D	5/5 (100%)	5 (100%)	0	100	100
2	E	5/5 (100%)	5 (100%)	0	100	100
2	F	5/5 (100%)	5 (100%)	0	100	100
All	All	435/471 (92%)	421 (97%)	14 (3%)	39	50

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	41	ASN
1	B	43	LYS
1	B	54	LYS
1	B	118	ARG
1	B	182	ARG
1	B	190	ARG
1	A	74	ASP
1	A	80	ARG
1	A	118	ARG
1	A	136	LYS
1	C	54	LYS
1	C	80	ARG
1	C	136	LYS
1	C	182	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	201	-	4,4,4	0.28	0	6,6,6	0.13	0
3	SO4	A	201	-	4,4,4	0.27	0	6,6,6	0.22	0
3	SO4	C	201	-	4,4,4	0.26	0	6,6,6	0.22	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	161/170 (94%)	0.08	0	100	100	38, 52, 70, 84	0
1	B	160/170 (94%)	-0.03	1 (0%)	89	89	34, 46, 73, 99	0
1	C	152/170 (89%)	0.22	4 (2%)	56	52	39, 56, 74, 95	0
2	D	6/6 (100%)	0.16	0	100	100	43, 46, 61, 68	0
2	E	6/6 (100%)	0.37	0	100	100	49, 54, 66, 76	0
2	F	6/6 (100%)	0.22	0	100	100	54, 58, 79, 82	0
All	All	491/528 (92%)	0.09	5 (1%)	82	81	34, 52, 74, 99	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	45	LEU	4.3
1	C	178	ALA	3.4
1	B	42	SER	3.4
1	C	44	LEU	2.9
1	C	43	LYS	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	201	5/5	0.85	0.52	30,30,30,30	0
3	SO4	B	201	5/5	0.88	0.50	30,30,30,30	0
3	SO4	C	201	5/5	0.91	0.53	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.