

Full wwPDB X-ray Structure Validation Report (i)

Jun 23, 2025 - 04:16 pm BST

PDB ID	:	$8 \mathrm{RIC} / \mathrm{pdb} 0000 \mathrm{sric}$
Title	:	Crystallographic structure of oligosaccharide dehydrogenase from Pycnoporus
		cinnabarinus bound to Sinapic Acid, tetragonal crystal
Authors	:	Savino, C.; Sciara, G.; Gugole, E.; Vallone, B.; Fata, F.; Bulfaro, G.; Costanzo,
		A.; Montemiglio, L.C.
Deposited on	:	2023-12-18
Resolution	:	1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359(1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	А	591	% • 90%	9% •					
1	В	591	% 90%	9% •					
2	С	2	50%	50%					
2	F	2	50%	50%					



Mol	Chain	Length	Quality of chain					
2	G	2	100%					
3	Е	4	75%		25%			
3	Н	4	50%	25%	25%			

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SXX	А	602	-	-	Х	-
6	SO4	В	610	-	-	Х	-



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 11151 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glucose oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	591	Total 4727	C 3014	N 791	O 907	S 15	0	55	0
1	В	591	Total 4762	C 3039	N 792	O 915	S 16	0	64	0

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	С	2	Total C N O 28 16 2 10	0	0	0
2	F	2	Total C N O 28 16 2 10	0	0	0
2	G	2	Total C N O 28 16 2 10	0	0	0

• Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluco pyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	Е	4	Total 50	C 28	N 2	O 20	0	0	0



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	Н	4	Total 50	C 28	N 2	O 20	0	0	0

• Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	Λ	1	Total	С	Ν	0	Р	0	0
4 A	1	53	27	9	15	2	0	0	
4	Р	1	Total	С	Ν	0	Р	0	0
4 B	L	53	27	9	15	2	0	0	

• Molecule 5 is SINAPINATE (CCD ID: SXX) (formula: C₁₁H₁₂O₅) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total C O 16 11 5	0	0
5	А	1	Total C O 16 11 5	0	0
5	А	1	Total C O 16 11 5	0	0
5	В	1	Total C O 16 11 5	0	0
5	В	1	Total C O 16 11 5	0	0
5	В	1	Total C O 16 11 5	0	0
5	В	1	Total C O 16 11 5	0	0

• Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O_4S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	А	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	604	Total O 604 604	0	12
8	В	542	Total O 542 542	0	5



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Glucose oxidase

Chain C:	50%	50%
AG1 AG2		

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

$OI \cdot D$		
Chain F:	50%	50%
Unam 1.	20%	50%

NAG 1 NAG 2

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose



Chain G:	100%		
NAG 1 NAG 2			
• Molecule 3: alpha- eta-D-glucopyranose	D-mannopyranose-(1-6)-beta-I -(1-4)-2-acetamido-2-deoxy-bet	D-mannopyranose-(1-4)-2-acetamido-2-de a-D-glucopyranose	eoxy-b
Chain E:	75%	25%	
NAG1 NAG2 BMA3 MAN4			
• Molecule 3: alpha- eta-D-glucopyranose	D-mannopyranose-(1-6)-beta-J (1-4)-2-acetamido-2-deoxy-bet	D-mannopyranose-(1-4)-2-acetamido-2-de za-D-glucopyranose	eoxy-b

Chain H:	50%	25%	25%
NAG1 NAG2 MAA3 MAN4			



4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 42 21 2	Depositor	
Cell constants	147.53Å 147.53Å 142.79Å	Deneriten	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
$\mathbf{P}_{\text{oscolution}}(\mathbf{\hat{A}})$	49.04 - 1.86	Depositor	
Resolution (A)	49.04 - 1.86	EDS	
% Data completeness	100.0 (49.04-1.86)	Depositor	
(in resolution range)	$100.0 \ (49.04 - 1.86)$	EDS	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	1.27 (at 1.86Å)	Xtriage	
Refinement program	REFMAC 5.8.0405	Depositor	
D D	0.169 , 0.202	Depositor	
Λ, Λ_{free}	0.172 , 0.205	DCC	
R_{free} test set	6686 reflections $(5.08%)$	wwPDB-VP	
Wilson B-factor $(Å^2)$	22.3	Xtriage	
Anisotropy	0.066	Xtriage	
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.36, 38.8	EDS	
L-test for twinning ²	$< L > = 0.50, < L^2 > = 0.33$	Xtriage	
Estimated twinning fraction	0.000 for -h,-l,-k	Vtriago	
Estimated twinning fraction	0.000 for l,-k,h	Atriage	
F_o, F_c correlation	0.96	EDS	
Total number of atoms	11151	wwPDB-VP	
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, SXX, BMA, SO4, NAG, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bo	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	1.06	0/5002	1.21	1/6811~(0.0%)
1	В	1.10	5/5057~(0.1%)	1.23	3/6884~(0.0%)
All	All	1.08	5/10059~(0.0%)	1.22	4/13695~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	282[A]	LYS	C-O	13.14	1.34	1.23
1	В	282[B]	LYS	C-O	13.14	1.34	1.23
1	В	282[A]	LYS	CA-C	5.91	1.58	1.53
1	В	282[B]	LYS	CA-C	5.91	1.58	1.53
1	В	179	MET	N-CA	5.28	1.51	1.46

All (5) bond length outliers are listed below:

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	6	THR	CB-CA-C	-6.65	94.80	109.56
1	А	463	ASP	CA-CB-CG	5.50	118.10	112.60
1	В	494	GLY	CA-C-O	-5.50	117.84	122.29
1	В	371	THR	CA-CB-OG1	-5.38	101.53	109.60

There are no chirality outliers.



All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	590	ALA	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4727	0	4739	60	0
1	В	4762	0	4808	44	0
2	С	28	0	25	1	0
2	F	28	0	25	1	0
2	G	28	0	25	0	0
3	Е	50	0	43	1	0
3	Н	50	0	43	1	0
4	А	53	0	31	5	0
4	В	53	0	31	3	0
5	А	48	0	33	9	0
5	В	64	0	43	2	0
6	А	50	0	0	4	0
6	В	50	0	0	4	0
7	А	14	0	13	0	0
8	А	604	0	0	15	2
8	В	542	0	0	7	0
All	All	11151	0	9859	119	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:439:ASN:HD21	1:A:441[B]:THR:CG2	1.42	1.32
1:A:439:ASN:ND2	1:A:441[B]:THR:HG23	1.47	1.25
1:A:439:ASN:ND2	1:A:441[B]:THR:CG2	2.00	1.21
1:A:439:ASN:HD21	1:A:441[B]:THR:HG21	1.17	1.07
1:A:403[A]:ASP:OD1	8:A:701:HOH:O	1.74	1.05



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:42[B]:GLN:OE1	1:A:281[B]:ARG:NH2	1.95	0.99
1:A:439:ASN:ND2	1:A:441[A]:THR:OG1	1.96	0.97
1:A:148[B]:ARG:HD3	8:A:1086[B]:HOH:O	1.63	0.96
1:A:535[A]:MET:HE3	1:A:548:LEU:HD23	1.58	0.86
1:A:145[B]:ASP:OD1	1:A:148[B]:ARG:NH2	2.10	0.85
1:B:535[A]:MET:HE2	1:B:546:ALA:HA	1.60	0.81
1:B:535[A]:MET:HE3	1:B:548:LEU:HD23	1.64	0.79
5:A:602:SXX:H3	8:A:845:HOH:O	1.84	0.77
1:A:403[A]:ASP:OD2	8:A:702:HOH:O	2.03	0.75
1:A:535[A]:MET:HE3	1:A:548:LEU:CD2	2.16	0.75
5:A:602:SXX:O7	5:A:603:SXX:O14	2.05	0.74
1:B:454:ASN:OD1	8:B:701:HOH:O	2.06	0.72
6:A:614:SO4:O3	8:A:703:HOH:O	2.09	0.69
5:B:604:SXX:O13	8:B:702:HOH:O	2.09	0.68
1:A:439:ASN:ND2	1:A:441[A]:THR:HG1	1.85	0.67
6:B:609:SO4:O4	8:B:703:HOH:O	2.10	0.67
1:B:125:TRP:CE2	1:B:535[A]:MET:HE1	2.31	0.66
5:A:602:SXX:H10	8:A:845:HOH:O	1.94	0.66
8:A:1150:HOH:O	2:C:1:NAG:H81	1.96	0.65
1:A:397:ILE:HG12	1:A:536[A]:MET:HE1	1.79	0.65
1:B:441[B]:THR:HG21	3:H:1:NAG:H82	1.80	0.64
1:B:125:TRP:NE1	1:B:535[A]:MET:HE1	2.13	0.63
5:A:602:SXX:C14	5:A:603:SXX:H5	2.28	0.62
5:A:602:SXX:H143	5:A:603:SXX:H5	1.82	0.61
5:A:602:SXX:H151	8:A:893:HOH:O	2.00	0.61
1:B:115:LEU:HD13	1:B:536[A]:MET:CE	2.32	0.60
1:A:535[A]:MET:CE	1:A:548:LEU:HD23	2.30	0.60
6:A:605:SO4:O1	8:A:704:HOH:O	2.16	0.59
1:B:535[A]:MET:HE3	1:B:548:LEU:CD2	2.32	0.59
1:B:196[B]:LYS:NZ	8:B:707:HOH:O	2.32	0.59
1:A:6[B]:THR:HG22	1:A:8:ASP:H	1.69	0.58
5:A:602:SXX:C15	8:A:893:HOH:O	2.52	0.58
1:B:337[A]:LYS:HB3	1:B:488[A]:VAL:HG22	1.85	0.58
1:A:439:ASN:CG	1:A:441[B]:THR:HG23	2.24	0.57
1:B:237[B]:GLN:H	1:B:237[B]:GLN:CD	2.13	0.57
1:B:445:ASN:H	1:B:445:ASN:HD22	1.52	0.57
1:A:535[A]:MET:HE2	1:A:546:ALA:HA	1.87	0.57
1:A:441[B]:THR:HG22	8:A:763:HOH:O	2.04	0.57
5:B:602:SXX:C14	5:B:603:SXX:H5	2.34	0.57
1:A:427:ILE:HD12	1:A:478[A]:LEU:CD2	2.35	0.57
1:A:427:ILE:HD12	1:A:478[A]:LEU:HD22	1.87	0.56



Atom-1	Atom-2	Interatomic	Clash
		distance (A)	overlap (A)
I:B:97:ASN:HB2	4:B:601:FAD:C5X	2.36	0.56
1:A:403[B]:ASP:OD1	8:A:705:HOH:O	2.18	0.55
1:B:83[A]:VAL:HG12	1:B:519[A]:ASN:OD1	2.06	0.55
1:A:97:ASN:HB2	4:A:601:FAD:C5X	2.37	0.53
8:B:1079:HOH:O	2:F:1:NAG:H83	2.08	0.53
1:A:397:ILE:CG1	1:A:536[A]:MET:HE1	2.38	0.53
1:A:115:LEU:HD13	1:A:536[A]:MET:HE3	1.92	0.52
1:A:535[A]:MET:CE	1:A:548:LEU:CD2	2.86	0.52
1:B:83[A]:VAL:CG1	1:B:519[A]:ASN:OD1	2.57	0.52
1:A:73:ASP:OD2	1:A:85[B]:HIS:HD2	1.94	0.51
1:B:115:LEU:HD13	1:B:536[A]:MET:HE3	1.92	0.51
1:B:399[A]:GLN:OE1	1:B:403[A]:ASP:OD1	2.29	0.51
1:A:397:ILE:CD1	1:A:536[A]:MET:HE1	2.40	0.51
1:B:26:LEU:C	1:B:26:LEU:HD23	2.35	0.51
1:B:397:ILE:HG12	1:B:536[A]:MET:HE1	1.93	0.50
1:A:73:ASP:OD2	1:A:85[B]:HIS:CD2	2.64	0.50
5:A:602:SXX:O7	5:A:602:SXX:H152	2.12	0.49
1:B:403[B]:ASP:HB3	6:B:610:SO4:O3	2.13	0.49
1:B:450[A]:LYS:HG2	8:B:977:HOH:O	2.12	0.48
1:A:164:PRO:HG2	1:A:228:TYR:CE2	2.49	0.48
1:A:503[A]:ASN:ND2	8:A:720:HOH:O	2.46	0.48
1:B:13:ASN:HD22	1:B:277:LYS:H	1.62	0.48
1:A:9:PRO:HB3	1:A:245[B]:HIS:CE1	2.49	0.47
1:A:145[B]:ASP:OD1	1:A:148[B]:ARG:CZ	2.62	0.47
1:B:560:ASP:O	1:B:563:MET:HG3	2.14	0.47
1:A:331:GLN:HB2	1:A:526:VAL:HG22	1.96	0.47
1:B:445:ASN:HD22	1:B:445:ASN:N	2.13	0.47
1:B:283[B]:GLU:HG2	1:B:555:ASN:HB2	1.96	0.47
1:B:1:ALA:HB2	1:B:237[B]:GLN:HG3	1.98	0.46
1:B:275[A]:ARG:HD3	8:B:906:HOH:O	2.15	0.46
1:A:571:HIS:HB3	4:A:601:FAD:C2	2.46	0.46
1:A:441[B]:THR:HG21	3:E:1:NAG:H82	1.98	0.46
1:B:256:ASN:ND2	6:B:612:SO4:O1	2.42	0.46
1:A:337[B]:LYS:HD3	1:A:424:ASP:OD2	2.16	0.45
1:A:397:ILE·HD11	1:A:536[A]:MET:HE1	1.98	0.45
1:A:408[B]:HIS:HE1	6:A:613:SO4:O2	1.99	0.45
1:B:13:ASN·ND2	1:B:277:LYS·H	2.15	0.45
1.A.1.ALA.HB2	$1 \cdot A \cdot 237 [A] \cdot GLN \cdot HG2$	1.99	0.44
1.A.161[B]·THR·HG?3	1·A·162·THR·HG23	1 99	0.44
1.B.9.PRO.HR3	$1 \cdot B \cdot 245 [B] \cdot HIS \cdot CE1$	2.52	0.44
1.B.527.ALA.HB1	4:B:601:FAD:HM83	1.99	0.44



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:548:LEU:O	1:A:557:ARG:HA	2.19	0.43
1:B:571:HIS:HB3	4:B:601:FAD:C2	2.48	0.43
1:B:548:LEU:O	1:B:557:ARG:HA	2.17	0.43
1:A:85[B]:HIS:HE1	8:A:1052:HOH:O	2.01	0.43
1:A:560:ASP:O	1:A:563:MET:HG3	2.19	0.43
1:A:198:LEU:HD12	1:A:198:LEU:C	2.44	0.43
1:A:439:ASN:ND2	1:A:441[B]:THR:HG21	1.93	0.43
1:A:275[B]:ARG:HD3	1:A:275[B]:ARG:HA	1.66	0.42
1:A:97:ASN:HB2	4:A:601:FAD:C4X	2.48	0.42
1:A:367:SER:O	1:A:371[A]:THR:OG1	2.35	0.42
1:A:527:ALA:HB1	4:A:601:FAD:HM83	1.99	0.42
1:B:243:ILE:O	1:B:245[B]:HIS:HD2	2.01	0.42
1:B:73:ASP:OD2	1:B:85[A]:HIS:ND1	2.46	0.42
1:B:403[A]:ASP:HB3	6:B:610:SO4:O3	2.19	0.42
1:A:61:ILE:HD12	1:A:61:ILE:HA	1.87	0.41
1:A:161[B]:THR:CG2	1:A:162:THR:HG23	2.51	0.41
1:A:253:ASP:HB3	1:A:262:VAL:HG11	2.02	0.41
1:A:26:LEU:C	1:A:26:LEU:HD23	2.46	0.41
1:A:80:GLN:N	6:A:606:SO4:O3	2.49	0.41
1:B:411:PRO:HB2	1:B:569:SER:HB3	2.02	0.41
1:A:64:TYR:CD2	5:A:602:SXX:H153	2.56	0.41
1:A:97:ASN:HB2	4:A:601:FAD:N5	2.36	0.41
1:A:526:VAL:O	1:A:527:ALA:HB3	2.21	0.40
1:B:261:ALA:HB1	1:B:284[A]:VAL:CG1	2.51	0.40
8:A:824:HOH:O	1:B:337[A]:LYS:HE3	2.21	0.40
1:B:397:ILE:CD1	1:B:536[A]:MET:HE1	2.51	0.40
1:B:249[B]:LYS:HG2	1:B:265[B]:GLU:HB2	2.03	0.40
1:A:512[A]:LYS:NZ	1:A:512[A]:LYS:CB	2.84	0.40
1:B:112:TRP:CZ2	1:B:564[B]:MET:HG3	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:1160:HOH:O	8:A:1196:HOH:O[2_565]	2.04	0.16
8:A:1196:HOH:O	8:A:1196:HOH:O[2_565]	2.06	0.14



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	644/591~(109%)	626~(97%)	18 (3%)	0	100	100
1	В	652/591~(110%)	625~(96%)	27 (4%)	0	100	100
All	All	1296/1182~(110%)	1251 (96%)	45 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	520/465~(112%)	502~(96%)	18 (4%)	31 16
1	В	529/465~(114%)	510~(96%)	19 (4%)	30 15
All	All	1049/930~(113%)	1012 (96%)	37 (4%)	41 16

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	146[A]	GLN
1	А	146[B]	GLN
1	А	198	LEU
1	А	249[A]	LYS
1	А	249[B]	LYS
1	А	370	GLN
1	А	371[A]	THR
1	А	371[B]	THR



Mol	Chain	Res	Type
1	А	419	SER
1	А	439	ASN
1	А	444	SER
1	А	464[A]	LEU
1	А	464[B]	LEU
1	А	478[A]	LEU
1	А	478[B]	LEU
1	А	512[A]	LYS
1	А	512[B]	LYS
1	А	525	SER
1	В	6	THR
1	В	79	ASP
1	В	198	LEU
1	В	284[A]	VAL
1	В	284[B]	VAL
1	В	445	ASN
1	В	450[A]	LYS
1	В	450[B]	LYS
1	В	464[A]	LEU
1	В	464[B]	LEU
1	В	478[A]	LEU
1	В	478[B]	LEU
1	В	488[A]	VAL
1	В	488[B]	VAL
1	В	496[A]	LYS
1	В	496[B]	LYS
1	В	525	SER
1	В	554[A]	THR
1	В	554[B]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	13	ASN
1	А	128	ASN
1	А	235	ASN
1	А	439	ASN
1	В	13	ASN
1	В	373	GLN
1	В	445	ASN
1	В	454	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Bos	Link	Bo	ond leng	ths	Bond angles		
WIOI	туре	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	NAG	С	1	2,1	14,14,15	0.81	0	$17,\!19,\!21$	2.46	6 (35%)
2	NAG	С	2	2	14,14,15	1.21	2 (14%)	17,19,21	1.45	2 (11%)
3	NAG	Е	1	3	14,14,15	0.41	0	17,19,21	0.61	0
3	NAG	E	2	3	14,14,15	0.31	0	17,19,21	0.59	0
3	BMA	Е	3	3	11,11,12	0.29	0	$15,\!15,\!17$	0.67	0
3	MAN	Е	4	3	11,11,12	0.22	0	$15,\!15,\!17$	0.59	0
2	NAG	F	1	2,1	14,14,15	0.69	0	$17,\!19,\!21$	2.79	8 (47%)
2	NAG	F	2	2	$14,\!14,\!15$	0.65	0	$17,\!19,\!21$	1.67	3 (17%)
2	NAG	G	1	2,1	$14,\!14,\!15$	0.53	0	$17,\!19,\!21$	1.23	2 (11%)
2	NAG	G	2	2	$14,\!14,\!15$	0.36	0	$17,\!19,\!21$	1.16	2 (11%)
3	NAG	Н	1	3	14,14,15	0.63	0	$17,\!19,\!21$	1.25	2 (11%)
3	NAG	Н	2	3	14,14,15	0.34	0	17,19,21	0.64	0
3	BMA	Н	3	3	11,11,12	0.27	0	$15,\!15,\!17$	0.69	1 (6%)
3	MAN	Н	4	3	11,11,12	0.23	0	$15,\!15,\!17$	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	С	1	2,1	-	4/6/23/26	0/1/1/1
2	NAG	С	2	2	-	2/6/23/26	0/1/1/1
3	NAG	Е	1	3	-	0/6/23/26	0/1/1/1
3	NAG	Е	2	3	-	0/6/23/26	0/1/1/1
3	BMA	Е	3	3	-	1/2/19/22	0/1/1/1
3	MAN	Е	4	3	-	1/2/19/22	0/1/1/1
2	NAG	F	1	2,1	-	4/6/23/26	0/1/1/1
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1
2	NAG	G	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
3	NAG	Н	1	3	-	2/6/23/26	0/1/1/1
3	NAG	Н	2	3	-	0/6/23/26	0/1/1/1
3	BMA	Н	3	3	-	1/2/19/22	0/1/1/1
3	MAN	Н	4	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	С	2	NAG	C1-C2	2.55	1.56	1.52
2	С	2	NAG	C2-N2	2.07	1.49	1.46

All	(26)	bond	angle	outliers	are	listed	below:
-----	------	------	-------	----------	-----	--------	--------

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	F	1	NAG	C1-O5-C5	8.60	123.85	112.19
2	С	1	NAG	C1-O5-C5	7.46	122.30	112.19
2	F	2	NAG	C4-C3-C2	4.50	117.61	111.02
2	С	2	NAG	C1-O5-C5	4.30	118.02	112.19
2	F	1	NAG	O3-C3-C2	-3.67	101.87	109.47
2	F	2	NAG	C1-O5-C5	3.17	116.49	112.19
2	G	1	NAG	C1-C2-N2	-3.09	105.21	110.49
2	F	1	NAG	O5-C5-C4	3.00	118.13	110.83
2	С	1	NAG	C8-C7-N2	2.89	120.98	116.10
2	С	1	NAG	O7-C7-C8	-2.86	116.75	122.06
2	G	2	NAG	C1-C2-N2	2.85	115.36	110.49
2	F	1	NAG	C6-C5-C4	-2.80	106.45	113.00
2	С	1	NAG	O5-C5-C4	2.68	117.35	110.83
2	G	2	NAG	C1-O5-C5	2.56	115.66	112.19
2	F	2	NAG	C3-C4-C5	2.54	114.77	110.24
2	F	1	NAG	C3-C4-C5	2.49	114.69	110.24
3	Н	1	NAG	C4-C3-C2	-2.49	107.37	111.02
2	С	1	NAG	C6-C5-C4	-2.45	107.27	113.00



Conti	nued fron	ı previ	ous page				
Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$ $ Ideal(o)
2	F	1	NAG	C8-C7-N2	2.29	119.97	116.10
2	С	2	NAG	C2-N2-C7	2.24	126.10	122.90
2	С	1	NAG	C2-N2-C7	2.19	126.02	122.90
2	F	1	NAG	O7-C7-C8	-2.15	118.07	122.06
3	Н	1	NAG	C2-N2-C7	-2.10	119.91	122.90
2	G	1	NAG	C4-C3-C2	2.05	114.02	111.02
3	Н	3	BMA	C1-O5-C5	2.02	114.93	112.19
2	F	1	NAG	C4-C3-C2	2.00	113.95	111.02

There are no chirality outliers.

Mol	Chain	\mathbf{Res}	Type	Atoms
2	С	1	NAG	O7-C7-N2-C2
2	С	1	NAG	C8-C7-N2-C2
2	F	1	NAG	O5-C5-C6-O6
2	С	1	NAG	C4-C5-C6-O6
2	F	2	NAG	C4-C5-C6-O6
2	F	1	NAG	C4-C5-C6-O6
2	F	1	NAG	C8-C7-N2-C2
2	F	1	NAG	O7-C7-N2-C2
3	Н	1	NAG	C8-C7-N2-C2
2	С	2	NAG	O5-C5-C6-O6
2	F	2	NAG	O5-C5-C6-O6
2	С	1	NAG	O5-C5-C6-O6
3	Н	1	NAG	O7-C7-N2-C2
3	Е	4	MAN	O5-C5-C6-O6
3	Е	3	BMA	O5-C5-C6-O6
3	Н	3	BMA	O5-C5-C6-O6
2	С	2	NAG	C4-C5-C6-O6

All (17) torsion outliers are listed below:

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	1	NAG	1	0
3	Е	1	NAG	1	0
3	Н	1	NAG	1	0
2	F	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,





bond angles, torsion angles, and ring geometry for oligosaccharide.

















5.6 Ligand geometry (i)

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Cha		n Bos	Tink	Bo	ond leng	Bond angles			
WIOI	туре	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SXX	В	602	-	16,16,16	0.77	1 (6%)	21,21,21	0.48	0
6	SO4	А	611	-	4,4,4	0.32	0	6,6,6	0.08	0
5	SXX	А	602	-	16,16,16	0.90	1 (6%)	21,21,21	0.58	0
6	SO4	А	605	-	4,4,4	0.35	0	6,6,6	0.09	0
6	SO4	В	614	-	4,4,4	0.38	0	6,6,6	0.07	0
6	SO4	А	612	-	4,4,4	0.35	0	6,6,6	0.24	0
6	SO4	А	614	-	4,4,4	0.39	0	6,6,6	0.21	0



Mol	Mol Type Chain		Dog	Tink	Bo	ond leng	$_{\rm sths}$	Bond angles		
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
6	SO4	В	612	-	4,4,4	0.44	0	$6,\!6,\!6$	0.11	0
6	SO4	В	615	-	4,4,4	0.37	0	$6,\!6,\!6$	0.09	0
5	SXX	В	604	-	16,16,16	0.90	1 (6%)	21,21,21	0.80	0
6	SO4	В	609	-	4,4,4	0.35	0	$6,\!6,\!6$	0.11	0
4	FAD	В	601	-	53,58,58	0.74	0	68,89,89	0.83	2 (2%)
6	SO4	А	609	-	4,4,4	0.35	0	$6,\!6,\!6$	0.09	0
7	NAG	А	615	1	14,14,15	0.62	0	$17,\!19,\!21$	1.65	6 (35%)
5	SXX	В	603	-	16,16,16	0.64	1 (6%)	21,21,21	0.59	0
6	SO4	А	610	-	4,4,4	0.50	0	$6,\!6,\!6$	0.37	0
6	SO4	В	606	-	4,4,4	0.32	0	$6,\!6,\!6$	0.11	0
5	SXX	А	603	-	16,16,16	0.68	1 (6%)	$21,\!21,\!21$	0.68	0
6	SO4	А	608	-	4,4,4	0.25	0	$6,\!6,\!6$	0.18	0
5	SXX	В	605	-	16, 16, 16	0.60	0	21,21,21	0.64	0
5	SXX	А	604	-	16,16,16	0.74	1 (6%)	$21,\!21,\!21$	0.47	0
6	SO4	В	610	-	4,4,4	0.28	0	$6,\!6,\!6$	0.15	0
6	SO4	В	607	-	4,4,4	0.38	0	$6,\!6,\!6$	0.19	0
6	SO4	В	613	-	4,4,4	0.40	0	$6,\!6,\!6$	0.10	0
6	SO4	В	608	-	4,4,4	0.35	0	$6,\!6,\!6$	0.40	0
4	FAD	А	601	-	53,58,58	0.78	2(3%)	$68,\!89,\!89$	0.80	0
6	SO4	А	613	-	4,4,4	0.27	0	$6,\!6,\!6$	0.20	0
6	SO4	В	611	-	4,4,4	0.47	0	$6,\!6,\!6$	0.10	0
6	SO4	A	607	-	4,4,4	0.35	0	6,6,6	0.14	0
6	SO4	А	606	-	4,4,4	0.41	0	$6,\!6,\!6$	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SXX	В	602	-	-	5/9/9/9	0/1/1/1
5	SXX	В	604	-	-	4/9/9/9	0/1/1/1
7	NAG	А	615	1	-	0/6/23/26	0/1/1/1
5	SXX	А	602	-	-	7/9/9/9	0/1/1/1
5	SXX	В	603	-	-	4/9/9/9	0/1/1/1
5	SXX	А	603	-	-	2/9/9/9	0/1/1/1
5	SXX	В	605	-	-	3/9/9/9	0/1/1/1
4	FAD	В	601	-	-	5/30/50/50	0/6/6/6
5	SXX	А	604	-	-	3/9/9/9	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	А	601	-	-	3/30/50/50	0/6/6/6

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
5	А	602	SXX	O13-C12	-3.07	1.22	1.30
5	В	602	SXX	O13-C12	-2.90	1.22	1.30
5	В	604	SXX	O13-C12	-2.86	1.22	1.30
5	А	603	SXX	O13-C12	-2.52	1.23	1.30
5	А	604	SXX	O13-C12	-2.28	1.24	1.30
5	В	603	SXX	O13-C12	-2.14	1.24	1.30
4	А	601	FAD	C1'-C2'	-2.13	1.49	1.52
4	А	601	FAD	C2-N3	-2.04	1.34	1.39

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
7	А	615	NAG	C1-O5-C5	2.64	115.77	112.19
7	А	615	NAG	C1-C2-N2	-2.62	106.02	110.49
4	В	601	FAD	O2P-P-O1P	2.59	125.07	112.24
4	В	601	FAD	C5A-C6A-N6A	2.59	124.28	120.35
7	А	615	NAG	C2-N2-C7	2.55	126.53	122.90
7	А	615	NAG	C3-C4-C5	2.30	114.34	110.24
7	А	615	NAG	C4-C3-C2	2.25	114.31	111.02
7	А	615	NAG	O5-C1-C2	2.04	114.50	111.29

There are no chirality outliers.

All (36) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	601	FAD	N10-C1'-C2'-O2'
4	В	601	FAD	N10-C1'-C2'-O2'
4	В	601	FAD	PA-O3P-P-O5'
5	В	602	SXX	C4-C10-C11-C12
5	В	604	SXX	C4-C10-C11-C12
5	В	602	SXX	C1-C2-O9-C15
5	В	602	SXX	C3-C2-O9-C15
5	А	602	SXX	C3-C2-O9-C15
5	В	603	SXX	C1-C6-O8-C14
5	В	605	SXX	C4-C10-C11-C12
5	A	602	SXX	C4-C10-C11-C12



Mol	Chain	Res	Type	Atoms
5	А	602	SXX	C1-C2-O9-C15
5	В	603	SXX	C5-C6-O8-C14
5	А	602	SXX	C1-C6-O8-C14
5	В	602	SXX	C10-C11-C12-O13
5	А	604	SXX	C11-C10-C4-C3
5	В	602	SXX	C10-C11-C12-O14
5	А	604	SXX	C11-C10-C4-C5
5	В	603	SXX	C10-C11-C12-O13
5	В	603	SXX	C10-C11-C12-O14
5	А	602	SXX	C11-C10-C4-C5
5	А	603	SXX	C10-C11-C12-O13
5	А	603	SXX	C10-C11-C12-O14
5	А	602	SXX	C11-C10-C4-C3
4	А	601	FAD	PA-O3P-P-O5'
5	В	605	SXX	C1-C6-O8-C14
5	А	602	SXX	C5-C6-O8-C14
5	В	605	SXX	C5-C6-O8-C14
5	В	604	SXX	C1-C2-O9-C15
4	А	601	FAD	O4B-C4B-C5B-O5B
5	А	604	SXX	C4-C10-C11-C12
5	В	604	SXX	C3-C2-O9-C15
4	В	601	FAD	P-O3P-PA-O2A
4	В	601	FAD	O4B-C4B-C5B-O5B
5	В	604	SXX	C10-C11-C12-O13
4	В	601	FAD	N10-C1'-C2'-C3'

Continued from previous page...

There are no ring outliers.

14 monomers are involved in 27 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	В	602	SXX	1	0
5	А	602	SXX	9	0
6	А	605	SO4	1	0
6	А	614	SO4	1	0
6	В	612	SO4	1	0
5	В	604	SXX	1	0
6	В	609	SO4	1	0
4	В	601	FAD	3	0
5	В	603	SXX	1	0
5	А	603	SXX	3	0
6	В	610	SO4	2	0
4	А	601	FAD	5	0



Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	А	613	SO4	1	0
6	А	606	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2		$OWAB(Å^2)$	Q<0.9	
1	А	591/591~(100%)	-0.44	4 (0%)	84	86	10, 21, 34, 68	55~(9%)
1	В	591/591~(100%)	-0.42	3 (0%)	87	89	10, 21, 33, 52	64 (10%)
All	All	1182/1182 (100%)	-0.43	7 (0%)	85	88	10, 21, 34, 68	119 (10%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	591	GLN	5.1
1	А	502	GLY	3.5
1	В	591[A]	GLN	3.2
1	В	1	ALA	3.1
1	А	439	ASN	2.8
1	А	403[A]	ASP	2.4
1	В	503[A]	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	NAG	F	2	14/15	0.49	0.17	72,98,107,109	0
3	NAG	Е	2	14/15	0.53	0.17	35,44,54,71	0



8F	2 T	\mathbf{C}
\mathbf{OI}	υı	\mathbf{U}

	The second		Dec		DCCC	DCD	\mathbf{D} for the rest (λ^2)	$\mathbf{O} < \mathbf{O} \mathbf{O}$
IVIOI	Type	Unain	Res	Atoms	RSUU	RSR	$B-factors(A^{-})$	Q<0.9
2	NAG	С	2	14/15	0.56	0.18	$68,\!90,\!99,\!106$	0
2	NAG	G	1	14/15	0.65	0.17	53,63,74,95	0
2	NAG	F	1	14/15	0.76	0.14	$55,\!67,\!75,\!89$	0
3	NAG	Е	1	14/15	0.78	0.14	24,26,27,32	0
2	NAG	С	1	14/15	0.79	0.15	48,62,74,75	0
2	NAG	G	2	14/15	0.79	0.16	$95,\!110,\!121,\!135$	0
3	BMA	Е	3	11/12	-	-	91,99,105,108	0
3	MAN	Е	4	11/12	-	-	30,30,30,30	0
3	NAG	Н	1	14/15	-	-	$20,\!22,\!25,\!29$	0
3	NAG	H	2	14/15	-	-	35,41,55,69	0
3	BMA	Н	3	11/12	-	-	76,97,106,109	0
3	MAN	H	4	11/12	-	-	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





















6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SXX	В	603	16/16	0.71	0.24	49,60,69,71	16
5	SXX	А	603	16/16	0.73	0.24	54,56,67,67	16
5	SXX	В	605	16/16	0.73	0.21	$59,\!66,\!77,\!82$	0
6	SO4	А	605	5/5	0.73	0.15	75,75,80,81	5
5	SXX	В	604	16/16	0.74	0.21	31,43,48,51	16
6	SO4	В	606	5/5	0.75	0.13	80,84,89,104	0
5	SXX	В	602	16/16	0.76	0.23	59,70,76,78	0
6	SO4	А	614	5/5	0.78	0.17	37,39,45,49	5
6	SO4	А	611	5/5	0.79	0.10	81,83,93,95	0
5	SXX	А	602	16/16	0.82	0.20	56,69,73,77	0
6	SO4	B	615	5/5	0.82	0.11	52,54,56,59	5



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors (A^2)	$Q{<}0.9$			
7	NAG	А	615	14/15	0.82	0.12	48,55,71,93	0			
5	SXX	А	604	16/16	0.83	0.17	38,47,58,62	0			
6	SO4	В	607	5/5	0.84	0.12	$54,\!55,\!65,\!71$	5			
6	SO4	В	610	5/5	0.84	0.11	$57,\!61,\!67,\!77$	0			
6	SO4	В	608	5/5	0.85	0.14	$63,\!64,\!69,\!75$	0			
6	SO4	В	614	5/5	0.85	0.14	45,46,49,51	5			
6	SO4	А	613	5/5	0.86	0.11	46,49,52,53	5			
6	SO4	А	612	5/5	0.86	0.11	57,59,62,74	5			
6	SO4	В	613	5/5	0.86	0.14	48,49,51,52	5			
6	SO4	В	612	5/5	0.88	0.11	44,44,51,52	5			
6	SO4	А	606	5/5	0.89	0.11	$50,\!51,\!56,\!58$	5			
6	SO4	В	609	5/5	0.89	0.10	58,62,67,70	5			
6	SO4	В	611	5/5	0.90	0.13	37,38,39,42	5			
6	SO4	А	609	5/5	0.91	0.08	$60,\!65,\!70,\!80$	0			
6	SO4	А	608	5/5	0.92	0.09	47,54,61,65	0			
6	SO4	А	607	5/5	0.93	0.08	49,52,52,56	5			
6	SO4	A	610	5/5	0.96	0.08	41,43,48,58	0			
4	FAD	A	601	53/53	0.98	0.04	13,17,20,21	0			
4	FAD	В	601	53/53	0.98	0.04	13,17,20,21	0			

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

































6.5 Other polymers (i)

There are no such residues in this entry.

