



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2025 – 04:49 PM EST

PDB ID : 8U04
Title : Reductasporine biosynthetic pathway imine reductase RedE, apo
Authors : Daniel-Ivad, P.; Ryan, K.S.
Deposited on : 2023-08-28
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6077 (1.64-1.60)
Clashscore	180529	6617 (1.64-1.60)
Ramachandran outliers	177936	6498 (1.64-1.60)
Sidechain outliers	177891	6497 (1.64-1.60)
RSRZ outliers	164620	6075 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	
1	C	315	
2	D	315	

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 9463 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RedE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	291	2095	1319	366	397	13	0	0	0
1	B	295	2102	1322	367	401	12	0	0	0
1	C	288	2076	1308	364	392	12	0	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A0F7G0Y4
A	-18	GLY	-	expression tag	UNP A0A0F7G0Y4
A	-17	SER	-	expression tag	UNP A0A0F7G0Y4
A	-16	SER	-	expression tag	UNP A0A0F7G0Y4
A	-15	HIS	-	expression tag	UNP A0A0F7G0Y4
A	-14	HIS	-	expression tag	UNP A0A0F7G0Y4
A	-13	HIS	-	expression tag	UNP A0A0F7G0Y4
A	-12	HIS	-	expression tag	UNP A0A0F7G0Y4
A	-11	HIS	-	expression tag	UNP A0A0F7G0Y4
A	-10	HIS	-	expression tag	UNP A0A0F7G0Y4
A	-9	SER	-	expression tag	UNP A0A0F7G0Y4
A	-8	SER	-	expression tag	UNP A0A0F7G0Y4
A	-7	GLY	-	expression tag	UNP A0A0F7G0Y4
A	-6	LEU	-	expression tag	UNP A0A0F7G0Y4
A	-5	VAL	-	expression tag	UNP A0A0F7G0Y4
A	-4	PRO	-	expression tag	UNP A0A0F7G0Y4
A	-3	ARG	-	expression tag	UNP A0A0F7G0Y4
A	-2	GLY	-	expression tag	UNP A0A0F7G0Y4
A	-1	SER	-	expression tag	UNP A0A0F7G0Y4
A	0	HIS	-	expression tag	UNP A0A0F7G0Y4
B	-19	MET	-	initiating methionine	UNP A0A0F7G0Y4
B	-18	GLY	-	expression tag	UNP A0A0F7G0Y4
B	-17	SER	-	expression tag	UNP A0A0F7G0Y4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	expression tag	UNP A0A0F7G0Y4
B	-15	HIS	-	expression tag	UNP A0A0F7G0Y4
B	-14	HIS	-	expression tag	UNP A0A0F7G0Y4
B	-13	HIS	-	expression tag	UNP A0A0F7G0Y4
B	-12	HIS	-	expression tag	UNP A0A0F7G0Y4
B	-11	HIS	-	expression tag	UNP A0A0F7G0Y4
B	-10	HIS	-	expression tag	UNP A0A0F7G0Y4
B	-9	SER	-	expression tag	UNP A0A0F7G0Y4
B	-8	SER	-	expression tag	UNP A0A0F7G0Y4
B	-7	GLY	-	expression tag	UNP A0A0F7G0Y4
B	-6	LEU	-	expression tag	UNP A0A0F7G0Y4
B	-5	VAL	-	expression tag	UNP A0A0F7G0Y4
B	-4	PRO	-	expression tag	UNP A0A0F7G0Y4
B	-3	ARG	-	expression tag	UNP A0A0F7G0Y4
B	-2	GLY	-	expression tag	UNP A0A0F7G0Y4
B	-1	SER	-	expression tag	UNP A0A0F7G0Y4
B	0	HIS	-	expression tag	UNP A0A0F7G0Y4
C	-19	MET	-	initiating methionine	UNP A0A0F7G0Y4
C	-18	GLY	-	expression tag	UNP A0A0F7G0Y4
C	-17	SER	-	expression tag	UNP A0A0F7G0Y4
C	-16	SER	-	expression tag	UNP A0A0F7G0Y4
C	-15	HIS	-	expression tag	UNP A0A0F7G0Y4
C	-14	HIS	-	expression tag	UNP A0A0F7G0Y4
C	-13	HIS	-	expression tag	UNP A0A0F7G0Y4
C	-12	HIS	-	expression tag	UNP A0A0F7G0Y4
C	-11	HIS	-	expression tag	UNP A0A0F7G0Y4
C	-10	HIS	-	expression tag	UNP A0A0F7G0Y4
C	-9	SER	-	expression tag	UNP A0A0F7G0Y4
C	-8	SER	-	expression tag	UNP A0A0F7G0Y4
C	-7	GLY	-	expression tag	UNP A0A0F7G0Y4
C	-6	LEU	-	expression tag	UNP A0A0F7G0Y4
C	-5	VAL	-	expression tag	UNP A0A0F7G0Y4
C	-4	PRO	-	expression tag	UNP A0A0F7G0Y4
C	-3	ARG	-	expression tag	UNP A0A0F7G0Y4
C	-2	GLY	-	expression tag	UNP A0A0F7G0Y4
C	-1	SER	-	expression tag	UNP A0A0F7G0Y4
C	0	HIS	-	expression tag	UNP A0A0F7G0Y4

- Molecule 2 is a protein called RedE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	278	2019	1273	353	382	11	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	MET	-	initiating methionine	UNP A0A0F7G0Y4
D	-18	GLY	-	expression tag	UNP A0A0F7G0Y4
D	-17	SER	-	expression tag	UNP A0A0F7G0Y4
D	-16	SER	-	expression tag	UNP A0A0F7G0Y4
D	-15	HIS	-	expression tag	UNP A0A0F7G0Y4
D	-14	HIS	-	expression tag	UNP A0A0F7G0Y4
D	-13	HIS	-	expression tag	UNP A0A0F7G0Y4
D	-12	HIS	-	expression tag	UNP A0A0F7G0Y4
D	-11	HIS	-	expression tag	UNP A0A0F7G0Y4
D	-10	HIS	-	expression tag	UNP A0A0F7G0Y4
D	-9	SER	-	expression tag	UNP A0A0F7G0Y4
D	-8	SER	-	expression tag	UNP A0A0F7G0Y4
D	-7	GLY	-	expression tag	UNP A0A0F7G0Y4
D	-6	LEU	-	expression tag	UNP A0A0F7G0Y4
D	-5	VAL	-	expression tag	UNP A0A0F7G0Y4
D	-4	PRO	-	expression tag	UNP A0A0F7G0Y4
D	-3	ARG	-	expression tag	UNP A0A0F7G0Y4
D	-2	GLY	-	expression tag	UNP A0A0F7G0Y4
D	-1	SER	-	expression tag	UNP A0A0F7G0Y4
D	0	HIS	-	expression tag	UNP A0A0F7G0Y4

- Molecule 3 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total I 3 3	0	0
3	B	2	Total I 2 2	0	0
3	C	2	Total I 2 2	0	0
3	D	1	Total I 1 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	C	1	Total Cl 1 1	0	0
4	D	1	Total Cl 1 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	303	Total 303	O 303	0	0
7	B	342	Total 342	O 342	0	0
7	C	267	Total 267	O 267	0	0
7	D	238	Total 238	O 238	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.78Å 186.72Å 71.69Å 90.00° 111.12° 90.00°	Depositor
Resolution (Å)	93.36 – 1.62 93.36 – 1.62	Depositor EDS
% Data completeness (in resolution range)	96.2 (93.36-1.62) 95.9 (93.36-1.62)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.62Å)	Xtrriage
Refinement program	PHENIX 1.19_4092	Depositor
R, R_{free}	0.164 , 0.208 0.164 , 0.205	Depositor DCC
R_{free} test set	7670 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtrriage
Anisotropy	0.359	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.466 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	9463	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GOL, CL, IOD, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2131	0.48	0/2911
1	B	0.25	0/2137	0.48	0/2919
1	C	0.25	0/2112	0.48	0/2886
2	D	0.25	0/2063	0.48	0/2820
All	All	0.25	0/8443	0.48	0/11536

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2095	0	2026	15	0
1	B	2102	0	2016	15	0
1	C	2076	0	2018	14	0
2	D	2019	0	1955	14	0
3	A	3	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	1	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	B	6	0	8	1	0
6	D	4	0	6	0	0
7	A	303	0	0	2	0
7	B	342	0	0	2	0
7	C	267	0	0	3	0
7	D	238	0	0	2	0
All	All	9463	0	8029	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:149:CYS:SG	7:D:571:HOH:O	2.53	0.67
1:B:289:ARG:NH1	7:B:403:HOH:O	2.29	0.65
1:A:270:ARG:HH12	1:A:287:PHE:HB3	1.63	0.62
1:A:138:GLN:OE1	1:A:156:ASN:ND2	2.30	0.62
1:C:22:LEU:HD11	1:C:46:ALA:HB2	1.80	0.62
2:D:22:LEU:HD11	2:D:46:ALA:HB2	1.80	0.61
1:C:191:ALA:HB3	1:C:254:ARG:HD3	1.81	0.61
1:B:22:LEU:HD11	1:B:46:ALA:HB2	1.82	0.60
1:C:80:ALA:HB1	1:C:109:HIS:CD2	2.38	0.58
1:C:208:ASN:ND2	7:C:411:HOH:O	2.33	0.56
2:D:33:ARG:HH22	2:D:67:ALA:HB2	1.72	0.54
1:B:285:VAL:O	1:B:289:ARG:HG2	2.07	0.54
1:C:97:VAL:HG13	3:C:301:IOD:I	2.79	0.53
1:A:34:THR:HG22	1:B:60:PRO:HB3	1.91	0.53
1:B:0:HIS:O	1:B:27:ARG:NE	2.43	0.52
1:C:98:HIS:ND1	7:C:404:HOH:O	2.27	0.50
1:C:267:LEU:HG	1:C:284:LEU:HD22	1.94	0.50
1:A:22:LEU:HD11	1:A:46:ALA:HB2	1.93	0.50
1:A:254:ARG:NH2	7:A:413:HOH:O	2.39	0.49
1:B:69:TYR:OH	1:B:115:ASP:OD2	2.25	0.49
1:A:215:GLY:HA2	1:A:218:MET:HE2	1.94	0.49
1:C:82:GLU:N	1:C:82:GLU:OE1	2.45	0.49
1:B:76:LEU:HD22	1:B:83:LEU:HD11	1.95	0.48
1:B:133:LEU:HB3	1:B:157:LEU:HD11	1.95	0.48
2:D:17:LEU:HD22	2:D:151:LEU:HD13	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:10:LEU:HD11	2:D:30:VAL:HB	1.96	0.48
1:C:160:ASP:HB3	1:C:163:MET:HG3	1.96	0.47
1:A:160:ASP:HB3	1:A:163:MET:HG3	1.97	0.47
1:A:225:VAL:HG13	1:A:285:VAL:HG11	1.97	0.47
1:C:10:LEU:HD21	1:C:30:VAL:HG21	1.96	0.46
1:A:221:TYR:HB3	1:A:282:ALA:HB2	1.95	0.46
2:D:207:ALA:O	2:D:211:MET:HG2	2.15	0.46
2:D:192:ASP:OD1	2:D:197:ASN:ND2	2.34	0.46
1:B:157:LEU:HD13	1:B:167:TYR:CD2	2.52	0.45
2:D:239:LEU:HD13	2:D:242:ARG:HH21	1.82	0.45
1:A:17:LEU:HD22	1:A:151:LEU:HD13	1.98	0.44
2:D:157:LEU:HD13	2:D:167:TYR:CD2	2.52	0.44
1:A:238:HIS:O	1:A:242:ARG:HG3	2.18	0.44
1:C:38:GLY:N	7:C:415:HOH:O	2.36	0.44
1:B:225:VAL:HG13	1:B:285:VAL:HG11	1.99	0.44
2:D:237:LEU:HD21	2:D:281:TYR:HA	2.00	0.44
1:B:77:ASP:OD2	5:B:301:GOL:H31	2.17	0.43
2:D:82:GLU:OE1	2:D:82:GLU:N	2.48	0.43
1:B:56:VAL:HG21	1:B:79:LEU:HD13	2.00	0.43
1:A:157:LEU:HD13	1:A:167:TYR:CD2	2.54	0.42
1:B:27:ARG:NH1	7:B:416:HOH:O	2.44	0.42
2:D:289:ARG:HD2	7:D:488:HOH:O	2.18	0.42
1:C:30:VAL:HG12	1:C:48:GLU:HA	2.01	0.42
1:C:10:LEU:HB2	1:C:32:ASN:HB2	2.02	0.42
1:C:17:LEU:HD23	1:C:125:ILE:HG21	2.02	0.42
1:B:77:ASP:HB3	1:B:78:PRO:HD3	2.02	0.41
2:D:238:HIS:CE1	2:D:277:GLY:HA3	2.55	0.41
1:A:77:ASP:HB3	1:A:78:PRO:HD3	2.03	0.41
1:A:208:ASN:ND2	7:A:417:HOH:O	2.42	0.41
1:B:-1:SER:N	1:B:25:GLY:HA2	2.36	0.41
1:A:222:ALA:HB3	1:A:223:PRO:HD3	2.04	0.40
2:D:66:LEU:HD12	2:D:72:VAL:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/315 (91%)	282 (98%)	5 (2%)	1 (0%)	37	20
1	B	290/315 (92%)	284 (98%)	6 (2%)	0	100	100
1	C	285/315 (90%)	274 (96%)	11 (4%)	0	100	100
2	D	274/315 (87%)	270 (98%)	4 (2%)	0	100	100
All	All	1137/1260 (90%)	1110 (98%)	26 (2%)	1 (0%)	48	28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	127	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/230 (88%)	199 (98%)	3 (2%)	60	39
1	B	200/230 (87%)	198 (99%)	2 (1%)	73	57
1	C	201/230 (87%)	198 (98%)	3 (2%)	60	39
2	D	198/231 (86%)	196 (99%)	2 (1%)	73	57
All	All	801/921 (87%)	791 (99%)	10 (1%)	67	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	VAL
1	A	235	PHE
1	A	254	ARG
1	B	98	HIS
1	B	235	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	107	GLN
1	C	235	PHE
1	C	289	ARG
2	D	34	THR
2	D	235	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	138	GLN
1	B	156	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	C	149	1	8,9,10	0.48	0	6,9,11	1.49	1 (16%)
1	CME	B	149	1	8,9,10	0.49	0	6,9,11	2.06	2 (33%)
1	CME	A	149	1	8,9,10	0.50	0	6,9,11	2.10	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	C	149	1	-	0/5/8/10	-
1	CME	B	149	1	-	1/5/8/10	-
1	CME	A	149	1	-	1/5/8/10	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	149	CME	CE-SD-SG	4.26	122.17	103.46
1	A	149	CME	CB-SG-SD	-4.05	93.38	103.86
1	C	149	CME	CE-SD-SG	3.42	118.46	103.46
1	A	149	CME	CE-SD-SG	2.88	116.11	103.46
1	B	149	CME	CB-SG-SD	-2.40	97.66	103.86

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	149	CME	SD-CE-CZ-OH
1	B	149	CME	CE-SD-SG-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 11 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	301	-	5,5,5	0.97	0	5,5,5	0.94	0
6	EDO	D	302	-	3,3,3	0.49	0	2,2,2	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	301	-	-	2/4/4/4	-
6	EDO	D	302	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	301	GOL	O1-C1-C2-C3
5	B	301	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	301	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/315 (92%)	-1.17	0 100 100	12, 19, 30, 46	0
1	B	294/315 (93%)	-1.16	0 100 100	12, 19, 31, 52	0
1	C	287/315 (91%)	-1.12	0 100 100	14, 21, 34, 91	0
2	D	278/315 (88%)	-1.09	0 100 100	13, 20, 35, 73	0
All	All	1149/1260 (91%)	-1.14	0 100 100	12, 20, 32, 91	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	B	149	10/11	0.99	0.04	9,15,39,45	0
1	CME	A	149	10/11	1.00	0.03	11,16,31,38	0
1	CME	C	149	10/11	1.00	0.03	14,22,34,42	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	301	6/6	0.98	0.05	26,26,26,27	0
6	EDO	D	302	4/4	0.98	0.07	25,25,25,25	0
4	CL	D	303	1/1	0.99	0.04	36,36,36,36	0
3	IOD	B	302	1/1	1.00	0.06	42,42,42,42	0
3	IOD	B	303	1/1	1.00	0.02	31,31,31,31	1
3	IOD	C	301	1/1	1.00	0.06	41,41,41,41	0
3	IOD	C	302	1/1	1.00	0.07	93,93,93,93	0
3	IOD	D	301	1/1	1.00	0.07	42,42,42,42	0
4	CL	A	303	1/1	1.00	0.01	30,30,30,30	0
4	CL	C	303	1/1	1.00	0.05	34,34,34,34	0
3	IOD	A	301	1/1	1.00	0.03	41,41,41,41	0
3	IOD	A	302	1/1	1.00	0.08	62,62,62,62	0
3	IOD	A	304	1/1	1.00	0.09	71,71,71,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.