

#### Full wwPDB X-ray Structure Validation Report (i)

Nov 12, 2024 – 06:52 AM JST

PDB ID : 8X3X

Title: ThDP-dependent HKA synthase

Authors : Yu, J.H.; Liu, T.

Deposited on : 2023-11-14

Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

#### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	Similar resolution $(\#\text{Entries, resolution range}(\mathring{A}))$		
$R_{free}$	164625	1881 (2.16-2.16)		
Clashscore	180529	2047 (2.16-2.16)		
Ramachandran outliers	177936	2027 (2.16-2.16)		
Sidechain outliers	177891	2026 (2.16-2.16)		
RSRZ outliers	164620	1882 (2.16-2.16)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	584	81%	9% •	9%
1	В	584	81%	10%	• 8%



#### 2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 8534 atoms, of which 70 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called BbmA.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	533	Total 3969	C 2532	11	O 737	S 26	0	0	0
1	В	537	Total 4008	C 2559	N 680	O 742	S 27	0	0	0


• Molecule 2 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula:  $C_{12}H_{19}N_4O_7P_2S$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf			
9	2 A	1	Total	С	Н	N	О	Р	S	0	0	
			42	12	16	4	7	2	1	0	0	
2	D	1	Total	С	Н	N	О	Р	S	0	0	
2	Б	D 1	42	12	16	4	7	2	1			


• Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	A 1	Total	С	Н	N	О	Р	0	0
3			39	10	12	5	10	2		0
9	D	B 1	Total	С	Н	N	О	Р	0	0
3	Б		39	10	12	5	10	2	U	0

• Molecule 4 is 3-(4-HYDROXY-PHENYL)PYRUVIC ACID (three-letter code: ENO) (formula: C<sub>9</sub>H<sub>8</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total 20	C 9	H 7	O 4	0	0



 $Continued\ from\ previous\ page...$ 

			$\mathbf{Mol}$
Total C H O	1	В	4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	В	4

• Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	В	1	Total Mg 1 1	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	203	Total O 203 203	0	0
6	В	150	Total O 150 150	0	0



#### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





#### 4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 2 21 21	Depositor	
Cell constants	97.34Å 110.36Å 112.51Å	Donositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	48.67 - 2.15	Depositor	
rtesolution (A)	48.67 - 2.15	EDS	
% Data completeness	99.9 (48.67-2.15)	Depositor	
(in resolution range)	99.9 (48.67-2.15)	EDS	
$R_{merge}$	0.08	Depositor	
$R_{sym}$	(Not available)	Depositor	
$< I/\sigma(I) > 1$	4.14 (at 2.16Å)	Xtriage	
Refinement program	PHENIX 1.17.1_3660	Depositor	
$R, R_{free}$	0.155 , $0.196$	Depositor	
it, it free	0.157 , $0.196$	DCC	
$R_{free}$ test set	64594 reflections $(2.98%)$	wwPDB-VP	
Wilson B-factor (Å <sup>2</sup> )	35.5	Xtriage	
Anisotropy	0.362	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, 44.9	EDS	
L-test for twinning <sup>2</sup>	$< L > = 0.48, < L^2> = 0.31$	Xtriage	
Estimated twinning fraction	0.017 for -h,l,k	Xtriage	
$F_o, F_c$ correlation	0.96	EDS	
Total number of atoms	8534	wwPDB-VP	
Average B, all atoms $(\mathring{A}^2)$	42.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

#### 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TPP, ENO, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.53	0/4056	0.61	1/5514~(0.0%)	
1	В	0.45	0/4096	0.56	0/5567	
All	All	0.49	0/8152	0.59	1/11081 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
1	A	245	SER	N-CA-CB	5.37	118.56	110.50

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3969	0	4007	55	0
1	В	4008	0	4050	45	0
2	A	26	16	16	0	0
2	В	26	16	16	0	0
3	A	27	12	12	0	0
3	В	27	12	12	1	0
4	A	13	7	7	1	0
4	В	13	7	7	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
5	A	1	0	0	0	0
5	В	1	0	0	0	0
6	A	203	0	0	4	0
6	В	150	0	0	2	0
All	All	8464	70	8127	94	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 6.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A	A. 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:514:LYS:HE3	1:A:534:MET:HE3	1.41	1.03
1:B:204:PHE:HB3	1:B:343:LYS:HE2	1.53	0.91
1:A:514:LYS:HE3	1:A:534:MET:CE	2.01	0.91
1:A:549:LEU:HD21	1:A:554:LYS:HB2	1.63	0.80
1:B:16:GLU:HG3	1:B:150:LEU:HD13	1.63	0.80
1:B:549:LEU:HD21	1:B:554:LYS:HB2	1.63	0.79
1:B:104:THR:HG22	1:B:168:ASP:OD2	1.87	0.75
1:B:136:ILE:HD11	1:B:163:LEU:HD22	1.69	0.75
1:B:534:MET:O	1:B:535:ASN:HB3	1.88	0.72
1:A:338:ILE:O	1:A:341:SER:HB3	1.92	0.70
1:B:268:ASP:O	1:B:272:SER:HB3	1.92	0.69
1:A:85:ILE:HG22	1:A:129:ILE:HD13	1.75	0.68
1:B:188:LYS:HE2	1:B:326:ASP:OD2	1.93	0.68
1:B:488:GLN:HE21	1:B:561:ASN:ND2	1.91	0.67
1:A:514:LYS:NZ	1:A:536:THR:HB	2.10	0.66
1:B:198:ARG:HG2	1:B:198:ARG:HH11	1.59	0.66
1:A:16:GLU:HG3	1:A:150:LEU:HD13	1.77	0.66
1:A:129:ILE:C	1:A:129:ILE:HD12	2.15	0.65
1:A:268:ASP:O	1:A:272:SER:HB3	1.96	0.65
1:A:204:PHE:HB3	1:A:343:LYS:NZ	2.12	0.65
1:B:204:PHE:CB	1:B:343:LYS:HE2	2.27	0.65
1:B:204:PHE:HB3	1:B:343:LYS:CE	2.25	0.64
1:A:79:THR:HB	1:B:82:LEU:HD23	1.80	0.63
1:B:204:PHE:CG	1:B:343:LYS:HD3	2.33	0.63
1:A:514:LYS:CE	1:A:534:MET:CE	2.75	0.63
1:B:534:MET:SD	1:B:538:CYS:HB2	2.40	0.62
1:A:148:ARG:HD2	6:A:831:HOH:O	2.01	0.60
1:B:147:ARG:NH1	6:B:701:HOH:O	2.20	0.60
1:A:514:LYS:CE	1:A:534:MET:HE3	2.24	0.57



Continued from previous page...

Continuea from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap(Å)
1:A:136:ILE:HD11	1:A:163:LEU:HD22	1.86	0.56
1:B:180:PRO:HG2	1:B:183:ASN:OD1	2.05	0.56
1:A:514:LYS:HB2	1:A:534:MET:HE1	1.86	0.55
1:A:255:LEU:O	1:A:265:LYS:HG3	2.07	0.55
1:A:149:ALA:HB1	1:A:161:ILE:HG21	1.89	0.54
1:A:549:LEU:CD2	1:A:554:LYS:HB2	2.37	0.54
1:A:129:ILE:O	1:A:129:ILE:CD1	2.56	0.54
1:A:514:LYS:HZ1	1:A:536:THR:HB	1.72	0.54
1:B:16:GLU:CG	1:B:150:LEU:HD13	2.37	0.53
1:A:536:THR:HB	1:A:537:PRO:HD2	1.90	0.53
1:B:276:ASP:HA	1:B:300:SER:HB3	1.90	0.53
1:A:28:PRO:HB2	1:A:100:ALA:HB1	1.90	0.53
1:A:374:PRO:HB2	1:A:401:TRP:CE2	2.44	0.52
1:B:556:ARG:O	1:B:556:ARG:HG2	2.10	0.52
1:A:82:LEU:HD23	1:B:79:THR:HB	1.92	0.52
1:A:129:ILE:HD12	1:A:129:ILE:O	2.09	0.51
1:A:115:SER:HB3	1:A:122:LEU:HG	1.93	0.51
1:B:115:SER:HA	1:B:120:VAL:O	2.11	0.50
1:A:190:ALA:HB1	1:A:191:PRO:HD2	1.92	0.50
1:B:338:ILE:O	1:B:341:SER:HB3	2.11	0.50
1:A:193:ILE:HG12	6:A:782:HOH:O	2.11	0.50
1:B:495:ILE:HG13	1:B:495:ILE:O	2.11	0.49
1:A:101:GLN:HB3	1:A:166:PRO:HA	1.93	0.49
1:A:514:LYS:HZ2	1:A:536:THR:HB	1.77	0.49
1:A:209:PRO:HD2	6:A:723:HOH:O	2.13	0.49
1:A:168:ASP:HB3	1:B:563:SER:O	2.13	0.48
1:A:420:LEU:HG	1:B:76:PRO:HD3	1.96	0.48
1:B:28:PRO:HB2	1:B:100:ALA:HB1	1.95	0.48
1:A:514:LYS:HB2	1:A:534:MET:CE	2.43	0.48
1:B:37:PHE:O	1:B:40:LYS:HE3	2.12	0.48
1:B:21:VAL:HG23	1:B:42:ILE:HG21	1.95	0.48
1:A:129:ILE:HD12	1:A:130:THR:HG23	1.95	0.47
1:A:453:MET:HG2	1:B:49:HIS:CD2	2.48	0.47
1:A:481:VAL:HG22	4:A:603:ENO:H5	1.95	0.47
1:A:129:ILE:C	1:A:129:ILE:CD1	2.81	0.47
1:B:531:ALA:O	1:B:534:MET:HG2	2.14	0.46
1:A:255:LEU:HA	1:A:265:LYS:HE2	1.96	0.46
1:A:151:ARG:HD3	1:A:184:TYR:O	2.16	0.46
1:B:169:ILE:HG13	1:B:170:ALA:N	2.31	0.46
1:A:302:ALA:HA	1:A:320:ASP:OD2	2.15	0.46
1:A:514:LYS:HZ2	1:A:536:THR:CB	2.30	0.44



Continued from previous page...

A + 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:148:ARG:NH2	6:A:705:HOH:O	2.50	0.44
1:A:514:LYS:HB2	1:A:514:LYS:HE2	1.72	0.44
1:A:514:LYS:HE2	1:A:514:LYS:H	1.83	0.44
1:B:198:ARG:HG2	1:B:198:ARG:NH1	2.31	0.44
1:B:22:PHE:O	1:B:70:CYS:HA	2.17	0.43
1:B:555:GLN:NE2	1:B:555:GLN:HA	2.32	0.43
1:A:374:PRO:HB2	1:A:401:TRP:CD2	2.53	0.43
1:A:276:ASP:HA	1:A:300:SER:HB2	1.99	0.43
1:B:168:ASP:OD1	1:B:168:ASP:N	2.48	0.43
1:A:255:LEU:C	1:A:265:LYS:HE2	2.39	0.42
1:B:562:LYS:O	1:B:562:LYS:HG2	2.16	0.42
1:A:106:ALA:HB1	1:A:111:ALA:HB2	2.01	0.42
1:B:190:ALA:HB1	1:B:191:PRO:HD2	2.01	0.42
1:B:307:ASP:OD1	3:B:602:ADP:O2'	2.37	0.42
1:B:101:GLN:HB3	1:B:166:PRO:HA	2.02	0.42
1:A:115:SER:HA	1:A:120:VAL:O	2.21	0.41
1:A:204:PHE:HB3	1:A:343:LYS:HZ1	1.80	0.41
1:A:181:PRO:HA	1:A:184:TYR:CZ	2.55	0.41
1:B:292:SER:OG	1:B:559:MET:HG3	2.20	0.41
1:B:336:LEU:HD23	1:B:336:LEU:HA	1.82	0.41
1:A:82:LEU:CD2	1:B:79:THR:HB	2.51	0.41
1:B:204:PHE:CD2	1:B:343:LYS:HD3	2.56	0.40
1:B:224:GLU:HG2	6:B:748:HOH:O	2.21	0.40
1:A:48:LYS:HA	1:A:48:LYS:HD3	1.90	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	529/584 (91%)	519 (98%)	9 (2%)	1 (0%)	44 44



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	В	533/584 (91%)	520 (98%)	13 (2%)	0	100	100
All	All	1062/1168 (91%)	1039 (98%)	22 (2%)	1 (0%)	48	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	245	SER

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	420/462 (91%)	411 (98%)	9 (2%)	48 53
1	В	$424/462 \ (92\%)$	416 (98%)	8 (2%)	52 57
All	All	844/924 (91%)	827 (98%)	17 (2%)	50 55

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	82	LEU
1	A	194	MET
1	A	341	SER
1	A	514	LYS
1	A	534	MET
1	A	535	ASN
1	A	554	LYS
1	A	556	ARG
1	В	208	ARG
1	В	328	SER
1	В	341	SER
1	В	343	LYS
1	В	394	ASP
1	В	556	ARG



Continued from previous page...

Mol	Chain	Res	Type
1	В	558	ASP
1	В	562	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	555	GLN
1	В	555	GLN
1	В	561	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Res	Link	Bond lengths			Bond angles		
WIOI IY	Type				Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ENO	В	603	-	13,13,13	1.04	0	17,17,17	1.41	1 (5%)
2	TPP	A	601	5	22,27,27	0.50	0	29,40,40	0.78	1 (3%)
3	ADP	В	602	-	24,29,29	0.91	1 (4%)	29,45,45	1.56	7 (24%)



Mol Type	Tuno	Chain	Res	Link	Bond lengths			Bond angles		
	туре		ites	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TPP	В	601	5	22,27,27	0.43	0	29,40,40	0.74	1 (3%)
3	ADP	A	602	-	24,29,29	0.82	0	29,45,45	1.52	5 (17%)
4	ENO	A	603	-	13,13,13	1.03	1 (7%)	17,17,17	1.77	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ENO	В	603	-	-	4/8/8/8	0/1/1/1
2	TPP	A	601	5	-	2/16/17/17	0/2/2/2
3	ADP	В	602	-	-	1/12/32/32	0/3/3/3
2	TPP	В	601	5	-	2/16/17/17	0/2/2/2
3	ADP	A	602	-	-	1/12/32/32	0/3/3/3
4	ENO	A	603	-	-	4/8/8/8	0/1/1/1

All (2) bond length outliers are listed below:

I	Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\rm Observed(\mathring{A})$	Ideal(Å)
	3	В	602	ADP	O4'-C1'	2.50	1.44	1.41
	4	A	603	ENO	O4-C2	-2.06	1.18	1.23

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	A	603	ENO	C4-C3-C2	-6.01	105.84	113.64
3	A	602	ADP	N3-C2-N1	-4.57	121.54	128.68
4	В	603	ENO	C4-C3-C2	-4.31	108.04	113.64
3	В	602	ADP	N3-C2-N1	-3.99	122.45	128.68
3	A	602	ADP	C1'-N9-C4	-3.51	120.47	126.64
3	В	602	ADP	C1'-N9-C4	-3.20	121.01	126.64
3	В	602	ADP	C2'-C3'-C4'	2.82	108.13	102.64
4	A	603	ENO	O1-C1-C2	2.31	120.28	113.97
3	В	602	ADP	C5-C6-N6	-2.23	116.97	120.35
3	В	602	ADP	N6-C6-N1	2.21	123.17	118.57
3	В	602	ADP	PA-O3A-PB	-2.15	125.44	132.83
3	A	602	ADP	O3'-C3'-C4'	-2.15	104.83	111.05
3	A	602	ADP	C2-N1-C6	2.14	122.41	118.75
3	В	602	ADP	O2A-PA-O1A	2.07	122.48	112.24



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	В	601	TPP	C5-C4-N3	2.05	111.68	107.57
3	A	602	ADP	N6-C6-N1	2.05	122.82	118.57
2	A	601	TPP	C5-C4-N3	2.02	111.61	107.57

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	TPP	PA-O3A-PB-O2B
2	В	601	TPP	PA-O3A-PB-O2B
4	В	603	ENO	O1-C1-C2-C3
2	A	601	TPP	PA-O3A-PB-O1B
4	A	603	ENO	O2-C1-C2-O4
4	A	603	ENO	O2-C1-C2-C3
4	В	603	ENO	O2-C1-C2-C3
4	A	603	ENO	O1-C1-C2-O4
4	В	603	ENO	O1-C1-C2-O4
4	A	603	ENO	O1-C1-C2-C3
2	В	601	TPP	PA-O3A-PB-O1B
3	A	602	ADP	O4'-C4'-C5'-O5'
3	В	602	ADP	O4'-C4'-C5'-O5'
4	В	603	ENO	O2-C1-C2-O4


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	602	ADP	1	0
4	A	603	ENO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















#### 5.7 Other polymers (i)

There are no such residues in this entry.



#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



#### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	533/584 (91%)	-0.40	8 (1%) 71 75	23, 36, 58, 99	0
1	В	537/584 (91%)	-0.23	16 (2%) 52 58	27, 41, 66, 89	0
All	All	$1070/1168 \; (91\%)$	-0.31	24 (2%) 62 67	23, 39, 63, 99	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	559	MET	4.2
1	A	557	PHE	4.0
1	A	343	LYS	3.6
1	A	514	LYS	3.6
1	В	417	GLY	3.5
1	A	204	PHE	3.4
1	В	204	PHE	3.3
1	A	558	ASP	3.3
1	A	417	GLY	3.1
1	В	371	GLY	3.0
1	В	194	MET	2.9
1	В	560	LEU	2.9
1	В	555	GLN	2.9
1	В	1	LYS	2.8
1	В	343	LYS	2.8
1	В	182	MET	2.7
1	В	563	SER	2.7
1	A	555	GLN	2.6
1	A	556	ARG	2.6
1	В	534	MET	2.3
1	В	564	TYR	2.3
1	В	561	ASN	2.2
1	В	557	PHE	2.2
1	В	336	LEU	2.1



#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
4	ENO	В	603	13/13	0.85	0.12	55,61,73,73	0
4	ENO	A	603	13/13	0.88	0.12	48,59,68,73	0
2	TPP	В	601	26/26	0.97	0.06	33,43,56,57	0
2	TPP	A	601	26/26	0.98	0.05	33,42,56,61	0
3	ADP	A	602	27/27	0.98	0.05	23,29,42,45	0
5	MG	A	604	1/1	0.98	0.03	46,46,46,46	0
3	ADP	В	602	27/27	0.99	0.04	28,34,43,46	0
5	MG	В	604	1/1	0.99	0.04	45,45,45,45	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 

### Electron density around ENO A 603:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












#### Electron density around TPP B 601:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around TPP A 601:

 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










## 




# 









#### 6.5 Other polymers (i)

There are no such residues in this entry.

