

Full wwPDB X-ray Structure Validation Report (i)

Mar 31, 2025 – 10:06 PM JST

PDB ID : 8ZWW / pdb 00008zww

Title : Structure-Based Mechanism and Specificity of Human Galactosyltransferase

B3GalT5

Authors : Lo, J.M.; Ma, C. Deposited on : 2024-06-13

Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 2.0rc1

EDS: FAILED buster-report: 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.42

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS failed to run properly.

Mol	Chain	Length	(Quality of chain					
1	A	278		89%	7% •				
1	В	278	79% 16%						
2	С	5	20%	80%					
2	D	5	60%		40%				
3	Е	2	50%	50	%				
3	G	2	50%	50	%				
4	F	3	33%	67%					
5	Н	6	33%	50%	17%				



2 Entry composition (i)


There are 10 unique types of molecules in this entry. The entry contains 5301 atoms, of which 323 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Beta-1,3-galactosyltransferase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total 2206	C 1432	N 371	O 390	S 13	0	3	0
1	В	266	Total 2189	C 1418	N 370	O 388	S 13	0	0	0

• Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-b eta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-b eta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	С	5	Total	С	Н	N	О	0	0	0	
2		9	103	32	44	1	26	U	0	U	
9	D	5	Total	С	Н	N	О	0	0	0	
2	Ъ	5	103	32	44	1	26	U	U		

• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	9	Total	С	Н	N	О	0	0	0
9	15	2	55	16	27	2	10	0	0	0
2	С	9	Total	С	Н	N	О	0	0	0
3	G	2	55	16	27	2	10	U	0	



• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[al pha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	3	Total	С	Н	N	0	0	0	0
			64	22	26	2	14			

• Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran ose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	Н	6	Total 138	C 40	H 67	N 2	O 29	0	0	0

• Molecule 6 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Mn 1 1	0	0
6	В	1	Total Mn 1 1	0	0


• Molecule 7 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		A	Aton	ns		ZeroOcc	AltConf		
7	Λ	1	Total	С	Н	N	О	Р	0	0	
'	A	1	36	9	11	2	12	2	0		
7	D	1	Total	С	Н	N	О	Р	0	0	
'	1 D	1	36	9	11	2	12	2	0	U	

• Molecule 8 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PRO PYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
0	Λ	1	Total	С	Н	N	О	0	0
0	A	1	45	11	26	2	6	U	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues		Ato	oms		ZeroOcc	AltConf	
0	D	1	Total	С	Н	N	О	0	0
0	Б	1	45	11	26	2	6	U	

 \bullet Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $\rm C_8H_{15}NO_6).$



Mol	Chain	Residues	${f Atoms}$					ZeroOcc	AltConf
0	B	1	Total	С	Н	N	О	0	0
9	Ъ	1	28	8	14	1	5	0	0

• Molecule 10 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	131	Total O 131 131	0	0
10	В	65	Total O 65 65	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.





• Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-al pha-D-galactopyranose-(1-4)-beta-D-galactopyranose

Chain D: 60% 40%

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 50% 50%





 \bullet Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 50% 50%



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 33% 67%



 $\bullet \ \, \text{Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]} \\ \text{beta-D-mannopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]} \\ 2\text{-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]} \\ 2\text{-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)} \\$

Chain H: 33% 50% 17%





4 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	47.71Å 86.13Å 87.61Å	Depositor
a, b, c, α , β , γ	90.00° 95.68° 90.00°	Depositor
Resolution (Å)	29.06 - 1.87	Depositor
% Data completeness	70.3 (29.06-1.87)	Depositor
(in resolution range)	70.9 (23.00 1.01)	_
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.52 (at 1.87Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.195 , 0.234	Depositor
Wilson B-factor (A^2)	21.7	Xtriage
Anisotropy	0.228	Xtriage
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5301	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.13% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, BMA, GLA, GAL, B3P, NGA, MN, MAN, FUC, NAG, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.43	0/2274	0.61	0/3075
1	В	0.32	0/2248	0.56	0/3040
All	All	0.38	0/4522	0.59	0/6115

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2206	0	2205	14	1
1	В	2189	0	2174	37	0
2	С	59	44	51	0	0
2	D	59	44	51	4	0
3	Е	28	27	25	0	0
3	G	28	27	25	0	0
4	F	38	26	34	2	0
5	Н	71	67	61	0	1
6	A	1	0	0	0	0
6	В	1	0	0	0	0
7	A	25	11	11	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	В	25	11	11	1	0
8	A	19	26	26	0	0
8	В	19	26	26	0	0
9	В	14	14	13	0	0
10	A	131	0	0	0	0
10	В	65	0	0	1	0
All	All	4978	323	4713	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash
Atom-1			overlap (Å)
1:B:144:ARG:HG2	1:B:144:ARG:HH11	1.52	0.73
1:B:113:SER:HA	1:B:119:ILE:HD13	1.73	0.69
1:B:241:LEU:HD13	2:D:4:NGA:H62	1.76	0.67
1:B:65:SER:HA	7:B:402:UDP:O2'	1.95	0.67
1:A:209:ARG:HG3	1:A:209:ARG:HH11	1.59	0.66
1:A:42:GLY:O	1:A:43:ASN:HB2	1.95	0.65
1:A:80:TRP:CZ3	1:A:161[B]:ILE:HD12	2.32	0.65
1:B:89:LYS:NZ	1:B:171:LEU:HD21	2.11	0.64
1:B:209:ARG:HH11	1:B:209:ARG:HG3	1.61	0.64
1:B:193:GLN:HB3	1:B:195:PHE:CE2	2.35	0.61
1:B:128:TYR:CE1	1:B:197:LYS:HE2	2.39	0.57
2:D:1:BGC:O1	2:D:1:BGC:H6C1	2.06	0.56
1:B:232:VAL:HG21	1:B:252:ARG:HG2	1.87	0.55
1:B:89:LYS:HZ2	1:B:171:LEU:HD11	1.72	0.55
1:B:58:PHE:CB	1:B:89:LYS:HB3	2.36	0.55
1:B:84:ARG:NH2	10:B:502:HOH:O	2.40	0.53
1:B:154:LYS:NZ	1:B:216:GLY:O	2.38	0.53
1:B:128:TYR:CZ	1:B:197:LYS:HE2	2.43	0.53
1:B:56:PRO:HB3	1:B:148:GLN:HB3	1.92	0.52
1:B:57:PRO:HA	1:B:90:GLN:HB3	1.92	0.52
1:B:144:ARG:HH11	1:B:144:ARG:CG	2.22	0.52
4:F:1:NAG:O5	4:F:3:FUC:H5	2.10	0.51
1:B:300:GLU:O	1:B:303:ARG:HG2	2.11	0.51
1:B:209:ARG:HG3	1:B:209:ARG:NH1	2.25	0.51
1:B:242:GLU:HG3	2:D:5:GAL:O4	2.10	0.50
1:A:128:TYR:CZ	1:A:197:LYS:HE2	2.47	0.50
1:B:113:SER:O	1:B:117:GLY:N	2.37	0.50

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({ m \AA})$	$ ext{overlap }(ext{Å})$
1:B:278:PHE:HA	1:B:281:ILE:HG12	1.94	0.48
4:F:1:NAG:H61	4:F:2:NAG:O7	2.14	0.48
1:A:107:LYS:O	1:A:111:GLN:HG3	2.14	0.48
1:B:63:VAL:HG22	1:B:155:THR:CG2	2.44	0.47
1:A:232:VAL:HG21	1:A:252:ARG:HG2	1.96	0.47
1:A:42:GLY:O	1:A:43:ASN:CB	2.62	0.47
1:B:198:TRP:HH2	2:D:1:BGC:H6C2	1.80	0.46
1:A:58:PHE:CB	1:A:89:LYS:HB3	2.46	0.46
1:A:155:THR:HG21	1:A:161[B]:ILE:HD11	1.97	0.45
1:B:141:TRP:CD1	1:B:145:PHE:HD2	2.35	0.44
1:A:155:THR:HG21	1:A:161[B]:ILE:CD1	2.47	0.44
1:B:89:LYS:HZ3	1:B:171:LEU:HD21	1.83	0.44
1:B:128:TYR:CD1	1:B:197:LYS:HE2	2.53	0.44
1:B:73:ARG:HG2	1:B:95:PHE:CD2	2.54	0.43
1:B:66:SER:O	1:B:73:ARG:NH1	2.48	0.43
1:B:70:LEU:O	1:B:74:MET:HG2	2.19	0.43
1:B:154:LYS:HD2	1:B:219:TYR:CZ	2.54	0.43
1:B:58:PHE:CG	1:B:89:LYS:HB3	2.53	0.42
1:A:194:PRO:HA	1:A:199:PHE:CD1	2.54	0.42
1:B:180:PHE:CD2	1:B:221:PHE:HB3	2.54	0.42
1:A:56:PRO:HB3	1:A:148:GLN:HB3	2.02	0.42
1:A:80:TRP:CH2	1:A:161[B]:ILE:HD12	2.55	0.41
1:A:180:PHE:CD2	1:A:221:PHE:HB3	2.55	0.41
1:B:232:VAL:CG2	1:B:252:ARG:HG2	2.50	0.41
1:B:45:LEU:HG	1:B:122:LYS:HA	2.03	0.40
1:B:154:LYS:O	1:B:218:GLY:HA2	2.21	0.40
1:B:175:ARG:HB3	1:B:175:ARG:CZ	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1 Atom-2		$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:209:ARG:NH1	5:H:1:NAG:O3[1_656]	2.10	0.10

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles
1	A	$268/278 \; (96\%)$	259 (97%)	8 (3%)	1 (0%)	30 18
1	В	$264/278 \; (95\%)$	256 (97%)	8 (3%)	0	100 100
All	All	532/556 (96%)	515 (97%)	16 (3%)	1 (0%)	44 34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	A	$247/255 \ (97\%)$	247 (100%)	0	100	100
1	В	244/255 (96%)	244 (100%)	0	100	100
All	All	491/510 (96%)	491 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

23 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Т	Clasia.	Das	T :1-	Вс	nd leng	ths	В	ond ang	cles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	С	1	2	12,12,12	0.22	0	17,17,17	0.54	0
2	GAL	С	2	2	11,11,12	1.04	1 (9%)	15,15,17	1.87	4 (26%)
2	GLA	С	3	2	11,11,12	1.57	2 (18%)	15,15,17	1.13	1 (6%)
2	NGA	С	4	2	14,14,15	0.56	0	17,19,21	1.17	2 (11%)
2	GAL	С	5	2	11,11,12	0.75	0	15,15,17	1.40	2 (13%)
2	BGC	D	1	2	12,12,12	0.47	0	17,17,17	1.35	3 (17%)
2	GAL	D	2	2	11,11,12	1.14	1 (9%)	15,15,17	1.41	1 (6%)
2	GLA	D	3	2	11,11,12	1.72	3 (27%)	15,15,17	0.97	1 (6%)
2	NGA	D	4	2	14,14,15	0.60	0	17,19,21	1.02	0
2	GAL	D	5	2	11,11,12	1.03	1 (9%)	15,15,17	1.53	4 (26%)
3	NAG	Е	1	3,1	14,14,15	0.34	0	17,19,21	0.43	0
3	NAG	Е	2	3	14,14,15	0.45	0	17,19,21	0.62	1 (5%)
4	NAG	F	1	1,4	14,14,15	1.29	1 (7%)	17,19,21	1.01	1 (5%)
4	NAG	F	2	4	14,14,15	1.40	2 (14%)	17,19,21	0.81	0
4	FUC	F	3	4	10,10,11	0.84	0	14,14,16	0.82	0
3	NAG	G	1	3,1	14,14,15	0.47	0	17,19,21	0.53	0
3	NAG	G	2	3	14,14,15	0.61	1 (7%)	17,19,21	0.79	1 (5%)
5	NAG	Н	1	5,1	14,14,15	0.28	0	17,19,21	0.98	1 (5%)
5	NAG	Н	2	5	14,14,15	0.54	0	17,19,21	0.54	0
5	BMA	Н	3	5	11,11,12	0.61	0	15,15,17	1.00	1 (6%)
5	MAN	Н	4	5	11,11,12	0.87	0	15,15,17	1.10	1 (6%)
5	MAN	Н	5	5	11,11,12	0.79	0	15,15,17	0.80	0
5	FUC	Н	6	5	10,10,11	1.08	1 (10%)	14,14,16	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	С	1	2	-	2/2/22/22	0/1/1/1
2	GAL	С	2	2	-	2/2/19/22	0/1/1/1
2	GLA	С	3	2	-	2/2/19/22	0/1/1/1
2	NGA	С	4	2	-	0/6/23/26	0/1/1/1
2	GAL	С	5	2	-	1/2/19/22	0/1/1/1
2	BGC	D	1	2	-	0/2/22/22	0/1/1/1
2	GAL	D	2	2	-	2/2/19/22	0/1/1/1
2	GLA	D	3	2	-	2/2/19/22	0/1/1/1
2	NGA	D	4	2	-	2/6/23/26	0/1/1/1
2	GAL	D	5	2	-	1/2/19/22	0/1/1/1
3	NAG	Е	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	Е	2	3	-	0/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	2/6/23/26	0/1/1/1
4	FUC	F	3	4	-	-	0/1/1/1
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
5	NAG	Н	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	Н	2	5	-	0/6/23/26	0/1/1/1
5	BMA	Н	3	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	4	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	5	5	-	0/2/19/22	0/1/1/1
5	FUC	Н	6	5	-	-	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	D	3	GLA	O5-C1	4.65	1.51	1.43
4	F	2	NAG	C1-C2	4.60	1.59	1.52
4	F	1	NAG	O5-C1	-4.50	1.36	1.43
2	С	3	GLA	O5-C1	4.14	1.50	1.43
2	D	5	GAL	O5-C1	-2.80	1.39	1.43
2	D	2	GAL	C1-C2	2.73	1.58	1.52
5	Н	6	FUC	O5-C1	-2.71	1.39	1.43
2	С	3	GLA	O5-C5	2.45	1.48	1.43
4	F	2	NAG	O5-C1	2.17	1.47	1.43
3	G	2	NAG	O5-C1	2.15	1.47	1.43
2	D	3	GLA	O5-C5	2.14	1.47	1.43
2	С	2	GAL	C2-C3	2.09	1.55	1.52
2	D	3	GLA	C2-C3	-2.01	1.49	1.52



All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
2	С	2	GAL	C1-O5-C5	4.66	118.51	112.19
2	D	1	BGC	C3-C4-C5	3.93	117.25	110.24
2	D	2	GAL	C1-O5-C5	3.76	117.28	112.19
2	С	5	GAL	C1-O5-C5	3.31	116.68	112.19
5	Н	1	NAG	C1-O5-C5	3.21	116.53	112.19
2	С	2	GAL	C2-C3-C4	-2.81	106.04	110.89
2	D	5	GAL	C1-O5-C5	2.73	115.90	112.19
4	F	1	NAG	O4-C4-C3	-2.71	104.09	110.35
5	Н	4	MAN	C1-C2-C3	-2.68	106.37	109.67
2	С	4	NGA	O5-C5-C6	2.66	111.37	107.20
2	С	5	GAL	O2-C2-C1	2.59	114.45	109.15
2	D	5	GAL	C2-C3-C4	-2.58	106.43	110.89
3	G	2	NAG	C1-O5-C5	2.45	115.51	112.19
2	D	1	BGC	O5-C5-C4	2.43	114.10	109.69
2	D	5	GAL	O5-C1-C2	2.36	114.42	110.77
2	С	3	GLA	C1-C2-C3	2.28	112.47	109.67
2	D	1	BGC	C4-C3-C2	2.27	114.78	110.82
2	С	4	NGA	O3-C3-C2	-2.19	104.92	109.47
2	С	2	GAL	O5-C5-C6	-2.15	103.84	107.20
2	D	3	GLA	O5-C5-C6	2.13	110.54	107.20
2	С	2	GAL	C1-C2-C3	2.12	112.28	109.67
2	D	5	GAL	C1-C2-C3	2.10	112.24	109.67
3	Е	2	NAG	C1-O5-C5	2.09	115.02	112.19
5	Н	3	BMA	C1-O5-C5	2.05	114.96	112.19

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	4	NGA	C8-C7-N2-C2
2	D	4	NGA	O7-C7-N2-C2
4	F	2	NAG	O5-C5-C6-O6
5	Н	1	NAG	C8-C7-N2-C2
5	Н	1	NAG	O7-C7-N2-C2
4	F	2	NAG	C4-C5-C6-O6
3	G	2	NAG	O5-C5-C6-O6
2	С	3	GLA	O5-C5-C6-O6
2	С	5	GAL	O5-C5-C6-O6
2	С	2	GAL	C4-C5-C6-O6
3	G	2	NAG	C4-C5-C6-O6
2	D	2	GAL	C4-C5-C6-O6

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	5	GAL	O5-C5-C6-O6
2	D	3	GLA	C4-C5-C6-O6
2	С	2	GAL	O5-C5-C6-O6
2	D	3	GLA	O5-C5-C6-O6
2	D	2	GAL	O5-C5-C6-O6
2	С	3	GLA	C4-C5-C6-O6
2	С	1	BGC	O5-C5-C6-O6
2	С	1	BGC	C4-C5-C6-O6

There are no ring outliers.


7 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	4	NGA	1	0
5	Н	1	NAG	0	1
2	D	1	BGC	2	0
4	F	1	NAG	2	0
2	D	5	GAL	1	0
4	F	2	NAG	1	0
4	F	3	FUC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.
























5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	cles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	UDP	A	402	6	24,26,26	0.41	0	37,40,40	0.77	2 (5%)
8	ВЗР	A	403	-	18,18,18	0.80	0	21,23,23	1.06	2 (9%)
9	NAG	В	404	1	14,14,15	0.18	0	17,19,21	0.67	0
7	UDP	В	402	6	24,26,26	0.37	0	37,40,40	0.69	1 (2%)
8	ВЗР	В	403	-	18,18,18	0.89	1 (5%)	21,23,23	0.91	1 (4%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	UDP	A	402	6	-	1/16/32/32	0/2/2/2
8	ВЗР	A	403	_	-	0/28/28/28	-
9	NAG	В	404	1	-	2/6/23/26	0/1/1/1
7	UDP	В	402	6	-	1/16/32/32	0/2/2/2
8	ВЗР	В	403	-	-	0/28/28/28	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
8	В	403	ВЗР	C3-N1	-2.14	1.44	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
8	В	403	ВЗР	C11-C8-C10	-2.45	104.85	110.04
7	A	402	UDP	O3B-PB-O2B	2.40	116.81	107.64
7	В	402	UDP	O3B-PB-O3A	2.28	112.27	104.64
8	A	403	ВЗР	C7-C4-C6	-2.23	105.34	110.04
7	A	402	UDP	O3B-PB-O3A	2.17	111.90	104.64
8	A	403	ВЗР	C7-C4-C5	-2.03	105.75	110.04

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	В	404	NAG	O5-C5-C6-O6
9	В	404	NAG	C4-C5-C6-O6
7	A	402	UDP	C5'-O5'-PA-O3A
7	В	402	UDP	PB-O3A-PA-O2A

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	В	402	UDP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,



bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.


6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.


6.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.












Electron density around Chain E: $\,$

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around Chain G:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












6.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around MN B 401: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)



















6.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

