



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 27, 2024 – 08:31 AM EDT

PDB ID : 9C7K  
Title : Crystal structure of pentalenene synthase variant F76A with PEG molecule in the active site  
Authors : Prem Kumar, R.; Ellenburg, W.H.; Oprrian, D.D.  
Deposited on : 2024-06-10  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	
1	B	337	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4908 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pentalenene synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	292	2268	1418	415	426	9	0	0	0
1	B	312	2458	1534	453	459	12	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	ALA	PHE	engineered mutation	UNP Q55012
B	76	ALA	PHE	engineered mutation	UNP Q55012

- Molecule 2 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	13	8	5	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	B	1	13	8	5	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0
3	A	1	5	4	1	0	0
3	A	1	5	4	1	0	0
3	A	1	5	4	1	0	0
3	B	1	5	4	1	0	0
3	B	1	5	4	1	0	0
3	B	1	5	4	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	84	84	84	0	0

Continued on next page...

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	B	37	Total	O	0	0
			37	37		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	182.42Å 182.42Å 56.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.98 – 2.50 19.98 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.98-2.50) 95.8 (19.98-2.50)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 2.50Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.211 , 0.239 0.211 , 0.239	Depositor DCC
$R_{free}$ test set	1862 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.3	Xtrriage
Anisotropy	0.257	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.030 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4908	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/2320	0.46	0/3152
1	B	0.23	0/2519	0.47	0/3423
All	All	0.23	0/4839	0.47	0/6575

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2268	0	2135	13	0
1	B	2458	0	2325	18	0
2	A	13	0	18	1	0
2	B	13	0	18	1	0
3	A	20	0	0	0	0
3	B	15	0	0	0	0
4	A	84	0	0	1	0
4	B	37	0	0	2	0
All	All	4908	0	4496	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:126:ARG:NH1	4:B:502:HOH:O	2.32	0.62
1:A:6:ASP:O	1:A:256:GLN:NE2	2.35	0.60
1:B:44:ARG:NH1	4:B:505:HOH:O	2.35	0.59
1:B:122:ASP:OD1	1:B:125:ARG:NH1	2.36	0.58
1:A:177:ILE:HG21	1:A:212:VAL:HG13	1.86	0.57
1:B:9:ILE:HG22	1:B:263:LEU:HG	1.87	0.56
1:A:224:LEU:HD13	1:A:237:VAL:HG11	1.90	0.54
1:B:163:CYS:O	1:B:242:ARG:NH2	2.42	0.53
1:B:149:GLY:HA3	1:B:176:THR:HG22	1.91	0.51
1:A:248:LYS:O	1:A:250:ARG:N	2.43	0.50
1:A:291:GLU:OE2	1:A:294:ARG:NH2	2.36	0.47
2:A:401:PG4:H42	2:A:401:PG4:H22	1.62	0.47
1:A:273:LEU:HD12	1:A:294:ARG:HD3	1.95	0.47
1:A:198:VAL:HG21	1:A:282:LEU:HD11	1.96	0.46
1:A:210:ILE:HG23	1:A:262:ARG:HB3	1.98	0.46
2:B:401:PG4:H41	2:B:401:PG4:H61	1.73	0.46
1:B:179:VAL:HB	1:B:212:VAL:HG22	1.98	0.45
1:B:8:HIS:CD2	1:B:260:ARG:HG3	2.52	0.45
1:B:110:PRO:HB2	1:B:112:THR:HG22	2.00	0.44
1:A:275:LYS:NZ	4:A:517:HOH:O	2.47	0.43
1:B:6:ASP:OD1	1:B:306:TYR:OH	2.24	0.43
1:B:227:GLU:HB3	1:B:232:GLU:HB3	2.00	0.43
1:B:61:ALA:HB2	1:B:189:ALA:HB1	2.01	0.43
1:B:238:MET:H	1:B:238:MET:HG2	1.36	0.42
1:A:258:GLU:OE1	1:A:262:ARG:NH1	2.49	0.42
1:A:22:ALA:HB1	1:A:70:VAL:HG21	2.01	0.42
1:A:35:LEU:HB3	1:A:78:LEU:HD13	2.00	0.42
1:A:179:VAL:HB	1:A:212:VAL:HG22	2.02	0.41
1:B:9:ILE:HG13	1:B:9:ILE:O	2.20	0.41
1:B:180:GLN:HB3	1:B:181:PRO:HD3	2.02	0.41
1:B:124:TRP:CZ2	1:B:140:ALA:HB2	2.55	0.41
1:B:224:LEU:HD21	1:B:248:LYS:HE3	2.03	0.41
1:B:16:SER:HB2	1:B:59:PRO:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/337 (84%)	280 (99%)	4 (1%)	0	100	100
1	B	310/337 (92%)	305 (98%)	5 (2%)	0	100	100
All	All	594/674 (88%)	585 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/272 (83%)	225 (100%)	0	100	100
1	B	247/272 (91%)	246 (100%)	1 (0%)	89	96
All	All	472/544 (87%)	471 (100%)	1 (0%)	92	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	238	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	402	-	4,4,4	0.24	0	6,6,6	0.11	0
3	SO4	B	404	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	A	405	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	A	404	-	4,4,4	0.24	0	6,6,6	0.08	0
2	PG4	B	401	-	12,12,12	0.12	0	11,11,11	0.61	0
3	SO4	B	402	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	B	403	-	4,4,4	0.23	0	6,6,6	0.08	0
2	PG4	A	401	-	12,12,12	0.13	0	11,11,11	0.58	0
3	SO4	A	403	-	4,4,4	0.23	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	B	401	-	-	5/10/10/10	-
2	PG4	A	401	-	-	7/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	PG4	O3-C5-C6-O4
2	A	401	PG4	O2-C3-C4-O3
2	A	401	PG4	C4-C3-O2-C2
2	B	401	PG4	O3-C5-C6-O4
2	A	401	PG4	O4-C7-C8-O5
2	A	401	PG4	C3-C4-O3-C5
2	B	401	PG4	O1-C1-C2-O2
2	B	401	PG4	C8-C7-O4-C6
2	B	401	PG4	O2-C3-C4-O3
2	A	401	PG4	C6-C5-O3-C4
2	A	401	PG4	C1-C2-O2-C3
2	B	401	PG4	C6-C5-O3-C4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	PG4	1	0
2	A	401	PG4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/337 (86%)	0.13	21 (7%) 23 21	42, 55, 111, 121	0
1	B	312/337 (92%)	0.21	7 (2%) 62 59	49, 73, 90, 99	0
All	All	604/674 (89%)	0.17	28 (4%) 38 35	42, 66, 101, 121	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	315	TYR	5.7
1	A	245	GLY	4.8
1	A	246	TRP	4.2
1	A	161	ALA	4.2
1	A	160	ASN	3.4
1	A	166	ALA	3.4
1	A	230	ARG	3.3
1	B	316	ASP	3.2
1	A	168	GLN	3.1
1	A	159	TRP	3.0
1	A	172	MET	3.0
1	B	314	ARG	2.9
1	A	90	ASN	2.8
1	A	238	MET	2.7
1	A	237	VAL	2.6
1	A	229	ALA	2.4
1	B	112	THR	2.4
1	A	175	HIS	2.3
1	A	177	ILE	2.3
1	A	249	SER	2.2
1	A	169	TYR	2.2
1	B	310	ARG	2.2
1	B	5	VAL	2.2
1	B	9	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	158	PHE	2.1
1	A	236	MET	2.0
1	A	235	ASN	2.0
1	A	250	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	B	402	5/5	0.64	0.19	105,110,112,126	0
3	SO4	B	403	5/5	0.74	0.18	105,108,121,121	0
3	SO4	A	402	5/5	0.78	0.17	79,81,99,105	0
3	SO4	A	405	5/5	0.79	0.18	102,103,110,117	0
3	SO4	A	404	5/5	0.82	0.15	79,79,100,109	0
2	PG4	B	401	13/13	0.82	0.15	70,79,82,83	0
3	SO4	B	404	5/5	0.83	0.23	105,106,119,121	0
2	PG4	A	401	13/13	0.85	0.14	56,64,75,76	0
3	SO4	A	403	5/5	0.93	0.15	65,76,87,88	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.