



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 15, 2025 – 02:09 PM EDT

PDB ID : 9CU3 / pdb_00009cu3
Title : Allosteric mechanism of DriD transcription activator
Authors : Cannistraci, E.G.; Schumacher, M.A.
Deposited on : 2024-07-25
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	
1	B	271	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3719 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WYL domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	196	Total	C	N	O	S	0	1	0
			1545	978	284	279	4			
1	A	203	Total	C	N	O	S	0	0	0
			1590	1007	289	290	4			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	57	MET	-	initiating methionine	UNP Q9A999
B	58	GLY	-	expression tag	UNP Q9A999
B	59	SER	-	expression tag	UNP Q9A999
B	60	SER	-	expression tag	UNP Q9A999
B	61	HIS	-	expression tag	UNP Q9A999
B	62	HIS	-	expression tag	UNP Q9A999
B	63	HIS	-	expression tag	UNP Q9A999
B	64	HIS	-	expression tag	UNP Q9A999
B	65	HIS	-	expression tag	UNP Q9A999
B	66	HIS	-	expression tag	UNP Q9A999
B	67	SER	-	expression tag	UNP Q9A999
B	68	SER	-	expression tag	UNP Q9A999
B	69	GLY	-	expression tag	UNP Q9A999
B	70	LEU	-	expression tag	UNP Q9A999
B	71	VAL	-	expression tag	UNP Q9A999
B	72	PRO	-	expression tag	UNP Q9A999
B	73	ARG	-	expression tag	UNP Q9A999
B	74	GLY	-	expression tag	UNP Q9A999
B	75	SER	-	expression tag	UNP Q9A999
B	76	HIS	-	expression tag	UNP Q9A999
B	77	MET	-	expression tag	UNP Q9A999
A	57	MET	-	initiating methionine	UNP Q9A999
A	58	GLY	-	expression tag	UNP Q9A999
A	59	SER	-	expression tag	UNP Q9A999
A	60	SER	-	expression tag	UNP Q9A999

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	HIS	-	expression tag	UNP Q9A999
A	62	HIS	-	expression tag	UNP Q9A999
A	63	HIS	-	expression tag	UNP Q9A999
A	64	HIS	-	expression tag	UNP Q9A999
A	65	HIS	-	expression tag	UNP Q9A999
A	66	HIS	-	expression tag	UNP Q9A999
A	67	SER	-	expression tag	UNP Q9A999
A	68	SER	-	expression tag	UNP Q9A999
A	69	GLY	-	expression tag	UNP Q9A999
A	70	LEU	-	expression tag	UNP Q9A999
A	71	VAL	-	expression tag	UNP Q9A999
A	72	PRO	-	expression tag	UNP Q9A999
A	73	ARG	-	expression tag	UNP Q9A999
A	74	GLY	-	expression tag	UNP Q9A999
A	75	SER	-	expression tag	UNP Q9A999
A	76	HIS	-	expression tag	UNP Q9A999
A	77	MET	-	expression tag	UNP Q9A999

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	292	Total	O	0	0
			292	292		
4	A	260	Total	O	0	0
			260	260		

- Molecule 1: WYL domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.80Å 88.80Å 141.52Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.40 – 1.90 44.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	90.3 (44.40-1.90) 90.3 (44.40-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.21-5207	Depositor
R, R_{free}	0.202 , 0.231 0.202 , 0.232	Depositor DCC
R_{free} test set	49520 reflections (4.22%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3719	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1629	0.60	0/2209
1	B	0.39	0/1584	0.62	0/2147
All	All	0.38	0/3213	0.61	0/4356

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1590	0	1568	13	0
1	B	1545	0	1516	13	0
2	A	6	0	8	2	0
2	B	6	0	8	1	0
3	A	5	0	0	0	0
3	B	15	0	0	0	0
4	A	260	0	0	6	0
4	B	292	0	0	5	0
All	All	3719	0	3100	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:255:ARG:HH22	2:B:401:GOL:H12	1.36	0.90
1:A:268:GLN:NE2	4:A:505:HOH:O	2.28	0.66
1:A:252:HIS:HE1	4:A:506:HOH:O	1.85	0.59
1:A:166:PHE:HB3	1:A:214:LEU:HD13	1.90	0.52
1:B:303:ILE:O	1:B:310:LYS:HE2	2.13	0.49
1:A:318:ARG:HA	1:A:327:TRP:HZ2	1.78	0.48
1:B:166:PHE:HB3	1:B:214:LEU:HD22	1.96	0.47
1:A:262:TRP:CH2	2:A:401:GOL:H12	2.50	0.46
1:B:200:GLY:HA2	4:B:740:HOH:O	2.16	0.45
1:B:252[A]:HIS:HE1	4:B:501:HOH:O	1.98	0.44
1:A:160:GLY:HA3	1:A:162:GLN:HE21	1.84	0.43
1:A:178:ARG:NH2	1:A:197:GLU:OE1	2.49	0.43
1:B:326:ALA:HB3	4:B:588:HOH:O	2.17	0.43
1:B:302:GLU:OE1	4:B:501:HOH:O	2.22	0.42
1:B:163:ALA:N	4:B:503:HOH:O	2.30	0.41
1:B:230:GLN:NE2	4:A:504:HOH:O	2.47	0.41
1:A:260:LEU:HD23	1:A:270:VAL:HG11	2.02	0.41
1:B:181:THR:HB	1:B:196:LEU:HG	2.01	0.41
1:B:268:GLN:HA	1:B:281:THR:O	2.21	0.41
1:A:322:ARG:HD2	4:A:610:HOH:O	2.19	0.41
1:A:288:ARG:NH1	4:A:507:HOH:O	2.41	0.41
1:B:260:LEU:HD23	1:B:270:VAL:HG21	2.03	0.41
1:A:183:LEU:HD11	1:A:196:LEU:HG	2.03	0.40
1:A:255:ARG:HH22	2:A:401:GOL:H32	1.84	0.40
1:A:269:GLN:HB2	4:A:660:HOH:O	2.20	0.40
1:B:165:SER:HA	1:B:178:ARG:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/271 (74%)	199 (99%)	2 (1%)	0	100	100
1	B	195/271 (72%)	192 (98%)	3 (2%)	0	100	100
All	All	396/542 (73%)	391 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/211 (77%)	159 (98%)	3 (2%)	52	49
1	B	156/211 (74%)	155 (99%)	1 (1%)	84	86
All	All	318/422 (75%)	314 (99%)	4 (1%)	65	65

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	214	LEU
1	A	130	LEU
1	A	167	ARG
1	A	225	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	148	GLN
1	A	162	GLN
1	A	225	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	402	-	4,4,4	0.66	0	6,6,6	0.31	0
3	SO4	B	403	-	4,4,4	0.75	0	6,6,6	0.11	0
2	GOL	A	401	-	5,5,5	0.44	0	5,5,5	0.86	0
2	GOL	B	401	-	5,5,5	0.31	0	5,5,5	0.59	0
3	SO4	A	402	-	4,4,4	0.70	0	6,6,6	0.15	0
3	SO4	B	404	-	4,4,4	1.04	0	6,6,6	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	401	-	-	4/4/4/4	-
2	GOL	A	401	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	GOL	O1-C1-C2-C3
2	B	401	GOL	C1-C2-C3-O3
2	A	401	GOL	C1-C2-C3-O3
2	A	401	GOL	O2-C2-C3-O3
2	A	401	GOL	O1-C1-C2-C3
2	B	401	GOL	O2-C2-C3-O3
2	A	401	GOL	O1-C1-C2-O2
2	B	401	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	GOL	2	0
2	B	401	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/271 (74%)	0.37	5 (2%) 58 60	26, 46, 84, 115	0
1	B	196/271 (72%)	-0.09	6 (3%) 51 53	20, 36, 62, 123	1 (0%)
All	All	399/542 (73%)	0.14	11 (2%) 55 57	20, 40, 81, 123	1 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	133	ALA	5.3
1	B	145	TYR	4.4
1	A	242	ASP	3.2
1	A	170	GLY	3.0
1	B	132	GLN	3.0
1	B	327	TRP	2.9
1	A	125	PRO	2.5
1	B	252[A]	HIS	2.4
1	A	214	LEU	2.3
1	A	327	TRP	2.2
1	B	134	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	404	5/5	0.71	0.17	37,37,49,68	0
3	SO4	A	402	5/5	0.76	0.11	89,93,101,103	0
3	SO4	B	403	5/5	0.88	0.09	66,66,72,79	0
2	GOL	B	401	6/6	0.91	0.11	43,46,47,51	0
2	GOL	A	401	6/6	0.93	0.10	32,35,39,40	0
3	SO4	B	402	5/5	0.97	0.06	48,50,52,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.