

Jul 1, 2025 – 12:18 PM EDT

PDB ID	:	$9 \mathrm{DDO} \ / \ \mathrm{pdb} \ 00009 \mathrm{ddo}$
EMDB ID	:	EMD-46778
Title	:	E. coli TonB-ExbBD TonB bound to ExbB chain C
Authors	:	Celia, H.; Botos, I.
Deposited on		
Resolution	:	2.80 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


EMDB validation analysis Mogul		0.0.1.dev118 2022.3.0, CSD as543be (2022)
ő		4-5-2 with Phenix2.0rc1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM} {f structures} \ (\#{f Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	244	89%	• 8%
1	В	244	92%	• 7%
1	С	244	90%	• 8%
1	D	244	91%	• 8%
1	Е	244	92%	7%
2	F	261	8% • 91%	
3	Y	163	19% • 80%	
3	Z	163	15% • 83%	



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 9252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Ate	oms			AltConf	Trace
1	А	224	Total	С	Ν	0	S	0	0
	A	224	1695	1079	296	314	6	0	0
1	В	227	Total	С	Ν	0	S	0	0
	D	221	1719	1093	300	319	7	0	0
1	С	224	Total	С	Ν	0	S	0	0
	C	224	1695	1079	296	314	6	0	0
1	D	225	Total	С	Ν	0	S	0	0
	D	220	1705	1085	299	315	6	0	0
1	E	226	Total	С	Ν	0	S	0	0
	Ľ	220	1712	1089	300	317	6		U

• Molecule 1 is a protein called Biopolymer transport protein ExbB.

• Molecule 2 is a protein called Protein TonB.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	F	23	Total 168	C 113	N 26	O 28	S 1	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	240	GLY	-	expression tag	UNP P02929
F	241	GLY	-	expression tag	UNP P02929
F	242	GLY	-	expression tag	UNP P02929
F	243	SER	-	expression tag	UNP P02929
F	244	GLU	-	expression tag	UNP P02929
F	245	ASN	-	expression tag	UNP P02929
F	246	LEU	-	expression tag	UNP P02929
F	247	TYR	-	expression tag	UNP P02929
F	248	PHE	-	expression tag	UNP P02929
F	249	GLN	-	expression tag	UNP P02929
F	250	GLY	-	expression tag	UNP P02929
F	251	GLY	-	expression tag	UNP P02929
F	252	SER	-	expression tag	UNP P02929

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
F	253	ALA	-	expression tag	UNP P02929
F	254	TRP	-	expression tag	UNP P02929
F	255	SER	-	expression tag	UNP P02929
F	256	HIS	-	expression tag	UNP P02929
F	257	PRO	-	expression tag	UNP P02929
F	258	GLN	-	expression tag	UNP P02929
F	259	PHE	-	expression tag	UNP P02929
F	260	GLU	-	expression tag	UNP P02929
F	261	LYS	-	expression tag	UNP P02929

Continued from previous page...

• Molecule 3 is a protein called Biopolymer transport protein ExbD.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	Y	32	Total 244	C 161		-	${ m S} { m 3}$	0	0
3	Z	28	Total 216	C 144			${ m S} { m 3}$	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	142	GLU	-	expression tag	UNP P0ABV2
Y	143	ASN	-	expression tag	UNP P0ABV2
Y	144	LEU	-	expression tag	UNP P0ABV2
Y	145	TYR	-	expression tag	UNP P0ABV2
Y	146	PHE	-	expression tag	UNP P0ABV2
Y	147	GLN	-	expression tag	UNP P0ABV2
Y	148	GLY	-	expression tag	UNP P0ABV2
Y	149	ASN	-	expression tag	UNP P0ABV2
Y	150	ALA	-	expression tag	UNP P0ABV2
Y	151	GLY	-	expression tag	UNP P0ABV2
Y	152	SER	-	expression tag	UNP P0ABV2
Y	153	GLY	-	expression tag	UNP P0ABV2
Y	154	HIS	-	expression tag	UNP P0ABV2
Y	155	HIS	-	expression tag	UNP P0ABV2
Y	156	HIS	-	expression tag	UNP P0ABV2
Y	157	HIS	-	expression tag	UNP P0ABV2
Y	158	HIS	-	expression tag	UNP P0ABV2
Y	159	HIS	-	expression tag	UNP P0ABV2
Y	160	HIS	-	expression tag	UNP P0ABV2
Y	161	HIS	-	expression tag	UNP P0ABV2
Y	162	HIS	-	expression tag \tilde{a}	UNP P0ABV2


Continued on next page...



	-	vious page		~	
Chain	Residue	Modelled	Actual	Comment	Reference
Y	163	HIS	-	expression tag	UNP P0ABV2
Z	142	GLU	-	expression tag	UNP P0ABV2
Z	143	ASN	-	expression tag	UNP P0ABV2
Z	144	LEU	-	expression tag	UNP P0ABV2
Z	145	TYR	-	expression tag	UNP P0ABV2
Z	146	PHE	-	expression tag	UNP P0ABV2
Z	147	GLN	-	expression tag	UNP P0ABV2
Z	148	GLY	-	expression tag	UNP P0ABV2
Z	149	ASN	-	expression tag	UNP P0ABV2
Z	150	ALA	-	expression tag	UNP P0ABV2
Z	151	GLY	-	expression tag	UNP P0ABV2
Z	152	SER	-	expression tag	UNP P0ABV2
Z	153	GLY	-	expression tag	UNP P0ABV2
Z	154	HIS	-	expression tag	UNP P0ABV2
Z	155	HIS	-	expression tag	UNP P0ABV2
Ζ	156	HIS	-	expression tag	UNP P0ABV2
Z	157	HIS	-	expression tag	UNP P0ABV2
Ζ	158	HIS	-	expression tag	UNP P0ABV2
Ζ	159	HIS	-	expression tag	UNP P0ABV2
Z	160	HIS	-	expression tag	UNP P0ABV2
Z	161	HIS	-	expression tag	UNP P0ABV2
Z	162	HIS	-	expression tag	UNP P0ABV2
Ζ	163	HIS	-	expression tag	UNP P0ABV2

Continued from previous page...

• Molecule 4 is (1S)-2-{[(2-AMINOETHOXY)(HYDROXY)PHOSPHORYL]OXY}-1-[(PA LMITOYLOXY)METHYL]ETHYL STEARATE (CCD ID: PEV) (formula: C₃₉H₇₈NO₈P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
4	D	1	Total	С	Ν	Ο	Р	0
4	D	1	49	39	1	8	1	0
4	Л	1	Total	С	Ν	0	Р	0
4	D	1	49	39	1	8	1	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Biopolymer transport protein ExbB





Chain F	8% •			91%				
MET THR LEU ASP LEU PRO	ARG ARG PHE V11 V11 V11 V12	A22 V32 HIS GLN VAL ILE	LEU PRO ALA ALA GLN PRO FRO	SER VAL THR MET VAL THR THR	ALA ASP ASP LEU GLU PRO PRO	GLN ALA VAL GLN PRO PRO	GLU PRO VAL VAL	GLU PRO GLU FRO GLU
PRO GLU PRO PRO PRO GLU	PRO PRO LYS GLU ALA PRO VAL VAL TLE	GLU LYS PRO LYS PRO PRO LYS LYS	LYS PRO LYS PRO PRO VAL LYS LYS	VAL GLN GLN GLN PRO LYS ARG	ASP VAL LYS PRO VAL GLU SER	ARG PRO ALA SER PRO PHE	ASN THR ALA PRO	ALA ARG LEU THR
SER SER THR ALA ALA	ALA THR SER LYS PRO VAL SER VAL	ALA SER GLY PRO ARG ALA LEU LEU	ANG ASN GLN FRO GLN FYR PRO ALA	ALA GLN ALA LEU ARG ILE	GLV GLN VAL LYS VAL LYS VAL	PHE ASP VAL THR PRO ASP	ALT ARG ASP ASN	VAL GLN ILE LEU
SER ALA LYS PRO ALA ASN	MET PHE GLU GLU VAL LYS ASN	MET ARG ARG ARG ARG CLU FLU	LYS LYS PRO GLY SER GLY ILE VAL	ASN ILE LEU LEU LYS ILE	ASN GLY THR THR GLU GLN GLN	GLY GLY GLY SER GLU ASN	TYR PHE GLY GLY	GLY SER ALA TRP
SER HIS PRO GLN PHE GLU	LYS							
• Molec	ule 3: Biopo	lymer trans	port protei	n ExbD				
Chain Y	: 19%	·		80%				
MET ALA MET HIS LEU ASN	GLU ASN LEU ASP D11 N12 V20	T 42 VAL VAL VAL VAL VAL ASP VAL ASN LEU	PRO ALA SER THR FRO PRO GLN	PRO ARG PRO GLU LYS PRO VAL	TYR LEU SER VAL LYS ALA	ASN ASN MET PHE ILE GLY	ASN ASP PRO VAL THR	ASP ASP GLU THR MET
ILE THR ALA LEU ASN ALA	LEU THR GLU GLU GLV GLY LYS LYS ASP ASP THR	ILE PHE ARG ALA ASP LYS THR	ASP TYR GLU THR LEU MET LYS	VAL ASP THR LEU HIS GLN	ALA GLY TYR LEU LYS ILE GLY	LEU VAL GLY GLU GLU THR	LYS ALA LYS GLU	ASN LEU TYR PHE
GLN GLY ASN ALA GLY SER	GLY HIS HIS HIS HIS HIS HIS HIS	SIH						
• Molec	ule 3: Biopo	lymer trans	port protei	n ExbD				
Chain Z	: 15% •			83%			_	
MET ALA MET HIS LEU ASN	GLU ASN LEU ASP ASP N12 V20 T21	L30 L33 LEU LEU	VAL ASP VAL LYS VAL ASN VAL ASN	ALA SER THR SER SER PRO	GLN PRO PRO GLU LYS PRO	VAL TYR LEU SER VAL LYS	ASP ASP SER MET	PHE ILE GLY ASN
ASP PRO VAL THR ASP GLU	THR MET ILE THR ALA ASN ASN LEU	THR GLU GLY LYS LYS LYS ASP THR THR	LLE PHE ARC ALA ASP LYS THR	ASP TYR GLU THR LEU MET	LIS VAL ASP THR LEU HIS	GLN ALA GLY TYR LEU LYS	GLY GLY VAL GLY	GLU GLU THR ALA
LYS ALA LYS GLU ASN LEU	TYR PHE GLN GLN GLY ASN ALA CLY SER GLY	HIS HIS HIS HIS HIS HIS SIH SIH SIH SIH	SIH					



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	227584	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	69.9	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.470	Depositor
Minimum map value	-0.164	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.018	Depositor
Recommended contour level	0.08	Depositor
Map size (Å)	212.48, 212.48, 212.48	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^{\circ}$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.83, 0.83, 0.83	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.66	0/1719	1.15	0/2326
1	В	0.67	0/1743	1.17	0/2358
1	С	0.70	0/1719	1.20	0/2326
1	D	0.66	0/1730	1.18	0/2341
1	Ε	0.67	0/1737	1.17	0/2351
2	F	0.73	0/173	1.18	0/239
3	Y	0.77	0/248	1.23	0/339
3	Ζ	0.66	0/220	1.10	0/300
All	All	0.68	0/9289	1.17	0/12580

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1695	0	1727	6	0
1	В	1719	0	1751	6	0
1	С	1695	0	1727	3	0
1	D	1705	0	1734	2	0
1	Е	1712	0	1741	0	0

Continued on next page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	168	0	176	1	0
3	Y	244	0	251	1	0
3	Ζ	216	0	224	2	0
4	В	49	0	77	0	0
4	D	49	0	77	0	0
All	All	9252	0	9485	17	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ARG:HA	1:A:117:ARG:HE	1.53	0.71
1:A:145:LEU:HD11	1:B:185:LEU:HD21	1.76	0.68
1:B:117:ARG:HA	1:B:117:ARG:NH2	2.10	0.65
1:D:114:ARG:HB3	1:D:114:ARG:HH11	1.65	0.61
1:A:142:PHE:HA	1:A:145:LEU:HG	1.83	0.61
1:D:114:ARG:HH11	1:D:114:ARG:CB	2.15	0.59
1:A:145:LEU:HD11	1:B:185:LEU:CD2	2.31	0.59
1:B:117:ARG:HA	1:B:117:ARG:CZ	2.37	0.54
1:C:156:PHE:CZ	3:Z:33:ILE:HG21	2.44	0.52
1:C:148:THR:HG22	1:C:177:ALA:O	2.10	0.52
1:B:117:ARG:HH21	1:B:117:ARG:HG2	1.75	0.52
1:C:148:THR:HG21	1:C:181:THR:OG1	2.13	0.49
1:A:94:GLU:OE1	1:A:114:ARG:NH2	2.47	0.47
1:B:118:ARG:HH21	1:B:118:ARG:HG3	1.80	0.46
3:Y:20:VAL:HG13	3:Z:21:THR:HG23	2.00	0.43
2:F:11:TRP:N	2:F:12:PRO:HD3	2.35	0.41
1:A:142:PHE:HA	1:A:145:LEU:CG	2.49	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	222/244~(91%)	222~(100%)	0	0	100	100
1	В	225/244~(92%)	225~(100%)	0	0	100	100
1	\mathbf{C}	222/244~(91%)	220~(99%)	2(1%)	0	100	100
1	D	223/244~(91%)	222 (100%)	1 (0%)	0	100	100
1	Ε	224/244~(92%)	224 (100%)	0	0	100	100
2	F	21/261~(8%)	21 (100%)	0	0	100	100
3	Y	30/163~(18%)	27~(90%)	3~(10%)	0	100	100
3	Ζ	26/163~(16%)	26 (100%)	0	0	100	100
All	All	1193/1807~(66%)	1187 (100%)	6 (0%)	0	100	100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	174/190~(92%)	173~(99%)	1 (1%)	84 95
1	В	177/190~(93%)	177~(100%)	0	100 100
1	\mathbf{C}	174/190~(92%)	174 (100%)	0	100 100
1	D	175/190~(92%)	172~(98%)	3~(2%)	56 84
1	Ε	176/190~(93%)	175~(99%)	1 (1%)	84 95
2	F	19/225~(8%)	19 (100%)	0	100 100
3	Υ	28/141~(20%)	28 (100%)	0	100 100
3	Ζ	25/141~(18%)	23~(92%)	2(8%)	10 30
All	All	948/1457~(65%)	941~(99%)	7(1%)	80 94

All (7) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	57	ARG
1	D	114	ARG
1	D	117	ARG
1	D	233	HIS
1	Е	37	THR
3	Ζ	20	VAL
3	Ζ	30	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	18	HIS
1	А	59	GLN
1	А	89	ASN
1	В	7	GLN
1	В	59	GLN
1	В	196	ASN
1	C C C C C C	18	HIS
1	С	59	GLN
1	С	60	GLN
1	С	69	ASN
1	С	89	ASN
1	С	104	ASN
1	С	220	GLN
1	D	89	ASN
1	D	104	ASN
1	Е	18	HIS
1	Е	59	GLN
1	Е	89	ASN
1	Е	201	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain Res		n Res Link		Link Bond lengths			Bond angles		
	Type	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	PEV	В	301	-	48,48,48	0.77	0	$51,\!53,\!53$	0.61	1 (1%)
4	PEV	D	301	-	48,48,48	0.77	1 (2%)	$51,\!53,\!53$	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEV	В	301	-	-	13/52/52/52	-
4	PEV	D	301	-	-	14/52/52/52	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
4	D	301	PEV	C1-C2	2.02	1.57	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	В	301	PEV	C2-O2-C31	2.36	123.44	117.80

There are no chirality outliers.

All (27) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
4	В	301	PEV	C32-C31-O2-C2
4	В	301	PEV	O31-C31-O2-C2
4	D	301	PEV	C32-C31-O2-C2
4	D	301	PEV	O31-C31-O2-C2
4	D	301	PEV	C2-C1-O3P-P
4	D	301	PEV	C22-C23-C24-C25
4	D	301	PEV	C11-C12-C13-C14
4	В	301	PEV	C19-C20-C21-C22
4	D	301	PEV	C5-C4-O4P-P
4	В	301	PEV	C2-C1-O3P-P
4	D	301	PEV	C18-C19-C20-C21
4	В	301	PEV	C1-C2-C3-O3
4	В	301	PEV	C34-C35-C36-C37
4	В	301	PEV	C1-O3P-P-O1P
4	В	301	PEV	C1-O3P-P-O2P
4	В	301	PEV	C1-O3P-P-O4P
4	D	301	PEV	C1-O3P-P-O2P
4	D	301	PEV	C39-C40-C41-C42
4	D	301	PEV	O3P-C1-C2-C3
4	В	301	PEV	C20-C21-C22-C23
4	D	301	PEV	C13-C14-C15-C16
4	В	301	PEV	C39-C40-C41-C42
4	D	301	PEV	O3P-C1-C2-O2
4	В	301	PEV	C44-C45-C46-C47
4	D	301	PEV	O31-C31-C32-C33
4	D	301	PEV	C20-C21-C22-C23
4	В	301	PEV	C45-C46-C47-C48

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-46778. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)


6.2.1 Primary map



X Index: 128




Y Index: 128



Z Index: 128

6.2.2 Raw map



X Index: 128

Y Index: 128




The images above show central slices of the map in three orthogonal directions.





6.3 Largest variance slices (i)

6.3.1 Primary map










Z Index: 89

6.3.2 Raw map



X Index: 145

Y Index: 110




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.08. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 67 nm^3 ; this corresponds to an approximate mass of 60 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.357 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.357 \AA^{-1}



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
Resolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	2.80	-	-	
Author-provided FSC curve	2.80	3.16	2.85	
Unmasked-calculated*	3.60	4.11	3.65	


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.60 differs from the reported value 2.8 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-46778 and PDB model 9DDO. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.08 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.08).



9.4 Atom inclusion (i)



At the recommended contour level, 98% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.



1.0

Map-model fit summary (i) 9.5

The table lists the average atom inclusion at the recommended contour level (0.08) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score	
All	0.8330	0.4890	
А	0.8440	0.4880	
В	0.8250	0.4910	
С	0.8640	0.4980	
D	0.8350	0.4890	
${ m E}$	0.8540	0.4950	
F	0.5840	0.3780	
Y	0.6840	0.4730	
Z	0.7500	0.4790	_ <

