



# Full wwPDB X-ray Structure Validation Report i

Feb 2, 2025 – 01:32 am GMT

PDB ID : 9FY8  
Title : L2A5 Fab in complex with STn-Thr  
Authors : Laugieri, M.E.; Ereno Orbea, J.; Jimenez Barbero, J.; Oyenarte, I.  
Deposited on : 2024-07-03  
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4623 (2.64-2.60)
Clashscore	180529	5071 (2.64-2.60)
Ramachandran outliers	177936	5006 (2.64-2.60)
Sidechain outliers	177891	5006 (2.64-2.60)
RSRZ outliers	164620	4622 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	V	126	 3% 87% 6% 6%
4	A	2	 50% 50%
4	E	2	 50% 50%

## 2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 8062 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L2A5 Fab Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	211	Total	C	N	O	S	0	0	0
			1603	1013	265	320	5			

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	I	209	Total	C	N	O	S	0	0	0
			1585	1002	262	316	5			

- Molecule 2 is a protein called L2A5 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	212	Total	C	N	O	S	0	0	0
			1618	1013	267	330	8			

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1628	1018	269	333	8			


- Molecule 3 is a protein called Human anti kappa variable heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	116	Total	C	N	O	S	0	0	0
			697	426	131	138	2			

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	V	118	Total	C	N	O	S	0	0	0
			708	432	133	141	2			

- Molecule 4 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-2-acetamido-2-deoxy-alpha-D-galactopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	2	Total	C	N	O		0	0	0
			34	19	2	13				

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	2	Total	C	N	O	0	0	0
			34	19	2	13			

- Molecule 5 is THREONINE (three-letter code: THR) (formula: C<sub>4</sub>H<sub>9</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			8	4	1	3		

5	I	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	24	Total	O	0	0
			24	24		
6	C	28	Total	O	0	0
			28	28		
6	D	20	Total	O	0	0
			20	20		
6	I	23	Total	O	0	0
			23	23		
6	L	32	Total	O	0	0
			32	32		
6	V	12	Total	O	0	0
			12	12		


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L2A5 Fab Heavy chain




- Molecule 1: L2A5 Fab Heavy chain




- Molecule 2: L2A5 Fab Light Chain





- Molecule 2: L2A5 Fab Light Chain



- Molecule 3: Human anti kappa variable heavy chain



- Molecule 3: Human anti kappa variable heavy chain



- Molecule 4: N-acetyl-alpha-neuraminic acid-(2-6)-2-acetamido-2-deoxy-alpha-D-galactopyranose



- Molecule 4: N-acetyl-alpha-neuraminic acid-(2-6)-2-acetamido-2-deoxy-alpha-D-galactopyranose



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.33Å 128.14Å 98.90Å 90.00° 98.17° 90.00°	Depositor
Resolution (Å)	53.61 – 2.62 53.61 – 2.62	Depositor EDS
% Data completeness (in resolution range)	97.4 (53.61-2.62) 97.4 (53.61-2.62)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.61 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.21rc1_5109: ????)	Depositor
$R$ , $R_{free}$	0.248 , 0.306 0.248 , 0.304	Depositor DCC
$R_{free}$ test set	1739 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.3	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.47$ , $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	8062	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A2G, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	B	0.25	0/1643	0.49	0/2242
1	I	0.25	0/1625	0.48	0/2218
2	C	0.25	0/1654	0.48	0/2245
2	L	0.25	0/1665	0.47	0/2261
3	D	0.20	0/336	0.47	0/417
3	V	0.22	0/342	0.47	0/425
All	All	0.24	0/7265	0.48	0/9808

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1603	0	1561	12	0
1	I	1585	0	1541	12	0
2	C	1618	0	1567	16	0
2	L	1628	0	1578	12	0
3	D	697	0	406	2	0
3	V	708	0	408	4	0
4	A	34	0	28	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	34	0	28	0	0
5	B	8	0	5	0	0
5	I	8	0	5	0	0
6	B	24	0	0	0	0
6	C	28	0	0	0	0
6	D	20	0	0	0	0
6	I	23	0	0	0	0
6	L	32	0	0	2	0
6	V	12	0	0	0	0
All	All	8062	0	7127	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:76:LYS:O	1:I:76:LYS:HD3	1.91	0.70
3:V:38:GLN:HB2	3:V:44:ARG:HG2	1.81	0.63
2:C:93:ASP:HB3	2:C:94:PRO:HD3	1.80	0.62
1:I:159:LEU:HD21	1:I:182:VAL:HG21	1.82	0.61
1:B:199:HIS:HB3	1:B:204:THR:HG23	1.83	0.61
2:L:34:TRP:HB2	2:L:47:ILE:HB	1.82	0.61
1:I:152:VAL:HG22	1:I:198:VAL:HG22	1.83	0.59
1:I:119:PRO:HB3	1:I:145:TYR:HB3	1.87	0.56
1:I:91:THR:HG23	1:I:110:THR:HA	1.88	0.56
2:L:93:ASP:HB3	2:L:94:PRO:HD3	1.88	0.55
3:V:12:VAL:HG11	3:V:85:LEU:HD12	1.89	0.55
2:C:65:GLY:HA3	2:C:70:TYR:HA	1.90	0.53
3:D:38:GLN:HB2	3:D:44:ARG:HG2	1.91	0.52
1:I:73:ASP:OD1	1:I:75:SER:OG	2.21	0.52
1:B:142:LYS:HE2	2:C:130:THR:HG21	1.91	0.52
1:B:6:GLU:OE2	1:B:94:TYR:HA	2.10	0.51
1:B:98:GLY:O	4:A:2:SIA:O7	2.29	0.50
2:C:143:ARG:HG3	2:C:174:TYR:CG	2.46	0.50
1:B:151:VAL:HG22	1:B:197:VAL:HG22	1.93	0.49
1:I:27:TYR:O	1:I:77:ASN:ND2	2.45	0.48
2:L:38:LYS:HG2	2:L:83:ALA:HB2	1.95	0.48
2:C:34:TRP:HB2	2:C:47:ILE:HB	1.95	0.48
2:L:187:TYR:O	2:L:193:TYR:OH	2.31	0.48
1:I:210:ARG:HG2	1:I:212:GLU:HG2	1.96	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:2:ILE:H	2:C:2:ILE:HD12	1.80	0.47
1:B:118:PRO:HB3	1:B:144:TYR:HB3	1.96	0.47
2:L:1:ASP:OD2	6:L:301:HOH:O	2.21	0.47
1:I:101:ASP:HA	2:L:45:ARG:HD3	1.95	0.47
2:C:29:VAL:O	2:C:70:TYR:OH	2.31	0.46
2:L:184:LYS:HE2	2:L:188:GLU:OE2	2.15	0.46
1:B:147:GLU:OE2	1:B:167:ALA:HB3	2.15	0.46
1:I:72:ARG:HA	1:I:79:PHE:HA	1.98	0.46
2:C:21:LEU:HD22	2:C:103:THR:HG21	1.98	0.46
2:L:186:ASP:HA	2:L:189:LYS:HD2	1.97	0.46
2:C:143:ARG:HG3	2:C:174:TYR:CD1	2.51	0.45
1:B:71:ARG:HA	1:B:78:PHE:HA	1.99	0.45
2:L:123:ASP:O	2:L:127:LYS:HG3	2.16	0.45
2:C:121:PRO:HD3	2:C:133:VAL:HG22	1.99	0.45
3:D:12:VAL:O	3:D:118:VAL:HA	2.17	0.45
1:I:198:VAL:HB	1:I:207:VAL:HG13	1.98	0.44
2:C:146:LYS:HB3	2:C:146:LYS:HE3	1.66	0.44
2:L:21:LEU:HD22	2:L:103:THR:HG21	2.00	0.44
1:B:90:THR:HG23	1:B:109:THR:HA	2.00	0.44
2:C:116:VAL:HA	2:C:136:LEU:O	2.18	0.44
1:B:125:PRO:HG2	1:B:212:PRO:HA	2.00	0.43
2:C:1:ASP:OD2	2:C:94:PRO:HB2	2.17	0.43
1:B:29:ILE:HA	1:B:35:TRP:CZ2	2.54	0.43
3:V:9:GLY:HA2	3:V:18:LEU:HD21	2.01	0.43
2:C:211:ASN:HB2	2:C:214:GLU:HG2	2.01	0.42
2:L:65:GLY:HA3	2:L:70:TYR:HA	2.01	0.42
3:V:12:VAL:O	3:V:118:VAL:HA	2.19	0.42
2:C:143:ARG:HH11	2:C:164:VAL:HG21	1.85	0.42
1:B:153:TRP:CH2	1:B:195:CYS:HB3	2.54	0.41
2:L:150:LYS:NZ	6:L:303:HOH:O	2.42	0.41
2:C:38:LYS:HG2	2:C:83:ALA:HB2	2.03	0.40
1:I:18:LEU:HD22	1:I:109:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	207/216 (96%)	201 (97%)	6 (3%)	0	100 100
1	I	205/216 (95%)	196 (96%)	9 (4%)	0	100 100
2	C	208/215 (97%)	202 (97%)	5 (2%)	1 (0%)	25 45
2	L	212/215 (99%)	206 (97%)	5 (2%)	1 (0%)	25 45
3	D	45/126 (36%)	44 (98%)	1 (2%)	0	100 100
3	V	46/126 (36%)	45 (98%)	1 (2%)	0	100 100
All	All	923/1114 (83%)	894 (97%)	27 (3%)	2 (0%)	44 65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	94	PRO
2	L	94	PRO

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	185/189 (98%)	182 (98%)	3 (2%)	58 78
1	I	182/189 (96%)	179 (98%)	3 (2%)	58 78
2	C	184/186 (99%)	181 (98%)	3 (2%)	58 78
2	L	185/186 (100%)	183 (99%)	2 (1%)	70 86
3	D	37/45 (82%)	37 (100%)	0	100 100
3	V	38/45 (84%)	37 (97%)	1 (3%)	41 66
All	All	811/840 (96%)	799 (98%)	12 (2%)	58 80

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	65	ASP
1	B	204	THR
1	B	205	LYS
2	C	11	MET
2	C	89	GLN
2	C	146	LYS
1	I	36	ASN
1	I	66	ASP
1	I	207	VAL
2	L	23	CYS
2	L	153	ASN
3	V	45	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	I	105	GLN
1	I	192	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	A2G	A	1	5,4	14,14,15	0.36	0	17,19,21	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SIA	A	2	4	20,20,21	0.81	1 (5%)	24,28,31	1.13	2 (8%)
4	A2G	E	1	5,4	14,14,15	0.37	0	17,19,21	0.68	0
4	SIA	E	2	4	20,20,21	0.85	1 (5%)	24,28,31	1.13	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A2G	A	1	5,4	-	2/6/23/26	0/1/1/1
4	SIA	A	2	4	-	1/18/34/38	0/1/1/1
4	A2G	E	1	5,4	-	0/6/23/26	0/1/1/1
4	SIA	E	2	4	-	0/18/34/38	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	2	SIA	C2-C1	2.91	1.55	1.52
4	A	2	SIA	C2-C1	2.74	1.54	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2	SIA	C6-O6-C2	3.31	118.41	111.34
4	E	2	SIA	C6-O6-C2	3.28	118.37	111.34
4	A	2	SIA	O1A-C1-C2	-3.03	115.42	122.57
4	E	2	SIA	O1A-C1-C2	-2.90	115.72	122.57

There are no chirality outliers.

All (3) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	A	1	A2G	C4-C5-C6-O6
4	A	1	A2G	O5-C5-C6-O6
4	A	2	SIA	O1A-C1-C2-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2	SIA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	THR	B	301	4	6,7,7	0.81	0	7,9,9	0.96	0
5	THR	I	301	4	6,7,7	0.80	0	7,9,9	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	THR	B	301	4	-	4/8/8/8	-
5	THR	I	301	4	-	1/8/8/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	301	THR	N-CA-CB-OG1
5	B	301	THR	N-CA-CB-CG2
5	B	301	THR	C-CA-CB-OG1
5	B	301	THR	C-CA-CB-CG2
5	I	301	THR	OXT-C-CA-N

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	211/216 (97%)	0.42	6 (2%) 55 50	19, 31, 48, 61	0
1	I	209/216 (96%)	0.51	8 (3%) 44 39	21, 32, 45, 58	0
2	C	212/215 (98%)	0.42	5 (2%) 59 55	17, 29, 46, 66	0
2	L	214/215 (99%)	0.31	6 (2%) 55 50	21, 31, 43, 59	0
3	D	47/126 (37%)	0.57	1 (2%) 63 59	24, 34, 43, 50	1 (2%)
3	V	48/126 (38%)	0.57	4 (8%) 19 16	20, 32, 46, 55	0
All	All	941/1114 (84%)	0.43	30 (3%) 50 45	17, 31, 45, 66	1 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	41	GLY	3.6
3	V	9	GLY	3.3
1	I	99	GLY	3.3
3	V	2	VAL	3.3
1	I	101	ASP	3.1
1	I	196	CYS	3.0
2	C	19	VAL	2.9
1	I	133	GLY	2.8
1	I	127	SER	2.8
2	L	93	ASP	2.8
2	C	1	ASP	2.7
2	C	94	PRO	2.6
1	I	76	LYS	2.6
2	C	194	ALA	2.5
1	I	195	ILE	2.5
3	V	45	GLU	2.4
3	V	3	GLN	2.4
2	L	195	CYS	2.4
1	B	198	ASN	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	65	ASP	2.3
2	L	169	SER	2.3
1	B	127	SER	2.2
1	B	42	GLY	2.2
2	L	165	THR	2.1
1	I	1	GLU	2.1
2	L	214	GLU	2.1
2	L	31	TYR	2.1
2	C	137	LEU	2.1
1	B	104	GLN	2.1
1	B	141	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

SUGAR-RSR INFOmissingINFO

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q<0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	THR	I	301	8/8	0.65	0.24	54,62,89,101	0
5	THR	B	301	8/8	0.83	0.15	39,42,63,69	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.