



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2025 – 12:10 AM JST

PDB ID : 9IHX / pdb_00009ihx
Title : Co-crystal structure of GH57 family amylopullulanase wild type from Aquifex
aeolicus with maltopentaose
Authors : Zhu, Z.M.; Wang, W.W.; Yu, F.
Deposited on : 2024-06-18
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	477	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 91%, yellow 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 91% 9% </div> </div>
1	B	477	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 88%, yellow 12%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 88% 12% </div> </div>
2	C	3	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 33%, yellow 33%, orange 33%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 33% 33% 33% </div> </div>
2	D	3	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 33%, yellow 33%, orange 33%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 33% 33% 33% </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase family 57 N-terminal domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	476	Total	C	N	O	S	0	0	0
			4044	2646	647	742	9			
1	B	476	Total	C	N	O	S	0	0	0
			4044	2646	647	742	9			

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	3	Total	C	O	0	0	0
			34	18	16			
2	D	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	298	Total	O	0	0
			298	298		
4	B	242	Total	O	0	0
			242	242		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycoside hydrolase family 57 N-terminal domain-containing protein




- Molecule 1: Glycoside hydrolase family 57 N-terminal domain-containing protein



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	61.53Å 41.19Å 195.08Å 90.00° 96.12° 90.00°	Depositor
Resolution (Å)	61.18 – 1.86 61.18 – 1.86	Depositor EDS
% Data completeness (in resolution range)	97.7 (61.18-1.86) 97.8 (61.18-1.86)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.86Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.178 , 0.225 0.180 , 0.223	Depositor DCC
R_{free} test set	4049 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8708	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/4157	0.51	1/5623 (0.0%)
1	B	0.30	0/4157	0.46	0/5623
All	All	0.32	0/8314	0.48	1/11246 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	109	ARG	NE-CZ-NH2	-5.35	114.38	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4044	0	3975	32	0
1	B	4044	0	3975	53	0
2	C	34	0	30	1	0
2	D	34	0	30	1	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	298	0	0	10	1
4	B	242	0	0	7	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8708	0	8026	84	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:324:GLU:CD	1:B:370:LYS:NZ	2.04	1.15
1:B:324:GLU:CD	1:B:370:LYS:HZ3	1.63	1.04
1:B:324:GLU:OE2	1:B:370:LYS:NZ	1.96	0.96
1:A:457:LYS:HG2	1:A:458:ALA:H	1.33	0.91
1:B:324:GLU:OE1	1:B:370:LYS:NZ	2.05	0.85
1:B:327:VAL:HG21	1:B:370:LYS:HG2	1.59	0.85
1:B:474:ARG:O	1:B:474:ARG:NH1	2.11	0.84
1:B:324:GLU:CD	1:B:370:LYS:HZ1	1.78	0.83
1:A:455:GLU:O	4:A:601:HOH:O	2.00	0.78
1:B:458:ALA:HB1	1:B:461:VAL:HB	1.67	0.75
1:B:251:TYR:HE1	1:B:390:MET:SD	2.11	0.74
1:B:120:GLU:OE2	4:B:601:HOH:O	2.06	0.73
1:B:109:ARG:HG3	1:B:114:ARG:HH21	1.55	0.72
1:A:457:LYS:O	4:A:602:HOH:O	2.07	0.72
1:A:394:ASP:OD2	4:A:603:HOH:O	2.08	0.71
1:B:465:ASP:OD2	4:B:602:HOH:O	2.08	0.71
1:B:457:LYS:HG3	1:B:458:ALA:H	1.55	0.70
1:A:458:ALA:HB1	1:A:461:VAL:HB	1.76	0.67
1:A:121:LYS:HD2	1:A:122:LEU:H	1.59	0.67
1:B:282:LEU:HG	1:B:289:ALA:HB1	1.79	0.64
1:B:391:ARG:NH2	4:B:604:HOH:O	2.29	0.64
1:B:457:LYS:CG	1:B:458:ALA:H	2.12	0.61
1:A:111:GLU:HA	1:A:114:ARG:HD3	1.84	0.60
1:B:325:ASP:OD1	1:B:328:ARG:NH1	2.30	0.59
1:A:29:TRP:CD2	2:C:3:GLC:H2	2.39	0.58
1:B:274:MET:HE3	1:B:390:MET:HB2	1.86	0.57
1:A:457:LYS:HD3	1:A:457:LYS:H	1.70	0.56
1:A:135:LEU:HD11	1:A:166:LEU:HG	1.89	0.55
1:A:81:ARG:HD3	4:A:627:HOH:O	2.09	0.53
1:B:219:ILE:HG12	4:B:616:HOH:O	2.08	0.53
1:A:152:ASN:OD1	4:A:604:HOH:O	2.18	0.53
1:B:462:GLU:H	1:B:462:GLU:CD	2.16	0.52
1:A:176:THR:O	1:A:179:ILE:HG22	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:387:GLU:OE2	4:B:603:HOH:O	2.18	0.52
1:B:100:LYS:HG2	1:B:117:MET:CE	2.39	0.52
1:A:457:LYS:HA	4:A:652:HOH:O	2.11	0.51
1:B:29:TRP:CD2	2:D:3:GLC:H2	2.46	0.50
1:A:153:LYS:HE2	1:A:155:ARG:O	2.11	0.50
1:B:90:GLU:H	1:B:90:GLU:CD	2.20	0.50
1:A:328:ARG:HE	1:B:49:GLU:HB2	1.75	0.50
1:A:324:GLU:HG3	1:A:370:LYS:HD3	1.94	0.50
1:A:228:GLU:HG3	1:A:232:LYS:NZ	2.27	0.49
1:B:224:VAL:HG21	1:B:434:LYS:HG2	1.94	0.49
1:B:282:LEU:HB2	1:B:305:PHE:CD2	2.48	0.49
1:A:80:ILE:HG21	1:A:135:LEU:HG	1.95	0.48
1:A:394:ASP:OD1	1:A:394:ASP:N	2.42	0.48
1:B:115:GLU:OE2	1:B:155:ARG:NH2	2.46	0.48
1:B:253:TRP:CZ3	1:B:350:VAL:HG11	2.49	0.48
1:B:319:HIS:HE1	4:B:656:HOH:O	1.97	0.47
1:B:251:TYR:CE1	1:B:390:MET:SD	2.99	0.47
1:B:438:ASN:ND2	1:B:474:ARG:HH12	2.11	0.47
1:A:253:TRP:CZ3	1:A:350:VAL:HG11	2.49	0.47
1:A:195:PRO:HD2	1:A:255:PRO:HD3	1.96	0.46
1:B:457:LYS:CG	1:B:458:ALA:N	2.79	0.46
1:A:97:GLU:HG3	4:A:621:HOH:O	2.15	0.46
1:B:224:VAL:CG2	1:B:434:LYS:HG2	2.46	0.45
1:B:48:PHE:HE2	1:B:373:SER:HB3	1.80	0.45
1:B:373:SER:O	1:B:377:LYS:HG3	2.16	0.45
1:A:167:ASN:O	1:A:171:GLU:HG3	2.17	0.44
1:B:195:PRO:HD2	1:B:255:PRO:HD3	2.00	0.44
1:B:270:LYS:HD3	1:B:270:LYS:HA	1.70	0.44
1:A:2:LYS:HE3	4:A:658:HOH:O	2.17	0.44
1:B:438:ASN:HD21	1:B:474:ARG:HH12	1.66	0.44
1:A:457:LYS:HG2	1:A:458:ALA:N	2.15	0.44
1:A:252:MET:HG3	1:A:272:ILE:HD12	2.00	0.43
1:A:379:GLU:OE1	1:A:379:GLU:N	2.49	0.43
1:B:27:LEU:HG	1:B:463:VAL:HG11	2.01	0.43
1:B:438:ASN:OD1	1:B:440:TYR:HB2	2.18	0.43
1:B:437:LYS:NZ	1:B:476:GLN:O	2.42	0.43
1:B:394:ASP:OD1	1:B:394:ASP:N	2.39	0.43
1:B:251:TYR:OH	1:B:387:GLU:HA	2.20	0.42
1:A:395:VAL:HG13	4:A:676:HOH:O	2.19	0.42
1:B:77:LEU:O	1:B:77:LEU:HD23	2.20	0.42
1:B:49:GLU:N	1:B:49:GLU:CD	2.78	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:ARG:NH2	4:A:611:HOH:O	2.43	0.42
1:B:176:THR:O	1:B:179:ILE:HG22	2.20	0.42
1:A:258:SER:HA	1:A:406:GLY:O	2.19	0.41
1:B:195:PRO:HG3	1:B:234:VAL:HG22	2.02	0.41
1:B:197:TYR:CZ	1:B:443:VAL:HG22	2.55	0.41
1:A:380:TRP:H	1:A:380:TRP:CD1	2.37	0.41
1:B:125:GLU:CD	1:B:125:GLU:H	2.29	0.41
1:B:195:PRO:HB3	1:B:233:HIS:HB3	2.02	0.41
1:B:114:ARG:HG2	4:B:771:HOH:O	2.21	0.41
1:B:380:TRP:CD1	1:B:380:TRP:H	2.39	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:651:HOH:O	4:B:803:HOH:O[2_455]	2.01	0.19
4:A:713:HOH:O	4:A:861:HOH:O[1_545]	2.04	0.16

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/477 (99%)	467 (98%)	6 (1%)	1 (0%)	44	32
1	B	474/477 (99%)	464 (98%)	9 (2%)	1 (0%)	44	32
All	All	948/954 (99%)	931 (98%)	15 (2%)	2 (0%)	44	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	457	LYS
1	A	457	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	436/437 (100%)	434 (100%)	2 (0%)	86	84
1	B	436/437 (100%)	436 (100%)	0	100	100
All	All	872/874 (100%)	870 (100%)	2 (0%)	92	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	286	VAL
1	A	435	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	453	GLN
1	A	476	GLN
1	B	134	ASN
1	B	273	ASN
1	B	319	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	C	1	2	12,12,12	1.43	2 (16%)	17,17,17	1.67	4 (23%)
2	GLC	C	2	2	11,11,12	0.57	0	15,15,17	0.93	0
2	GLC	C	3	2	11,11,12	0.64	0	15,15,17	1.21	1 (6%)
2	GLC	D	1	2	12,12,12	0.91	0	17,17,17	1.24	1 (5%)
2	GLC	D	2	2	11,11,12	0.57	0	15,15,17	0.89	0
2	GLC	D	3	2	11,11,12	0.67	0	15,15,17	1.20	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	C	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	3	2	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	GLC	O5-C5	-2.62	1.38	1.44
2	C	1	GLC	O5-C1	-2.31	1.37	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GLC	C1-O5-C5	3.58	120.42	113.66
2	C	3	GLC	C1-O5-C5	3.45	116.87	112.19
2	C	1	GLC	O5-C5-C4	2.64	114.50	109.69
2	D	3	GLC	O5-C1-C2	-2.44	107.01	110.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GLC	O5-C1-C2	2.43	114.61	110.28
2	D	3	GLC	C1-O5-C5	2.43	115.48	112.19
2	D	1	GLC	C1-O5-C5	2.30	118.01	113.66
2	C	1	GLC	O2-C2-C3	2.13	115.26	110.35

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	GLC	1	0
2	D	3	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	501	-	5,5,5	1.09	1 (20%)	5,5,5	0.95	0
3	GOL	B	501	-	5,5,5	0.84	0	5,5,5	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	501	-	-	2/4/4/4	-
3	GOL	B	501	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	GOL	O2-C2	-2.00	1.37	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	GOL	C1-C2-C3-O3
3	A	501	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/477 (99%)	-0.40	4 (0%) 82 85	11, 20, 41, 70	0
1	B	476/477 (99%)	-0.19	5 (1%) 77 81	14, 25, 47, 69	0
All	All	952/954 (99%)	-0.30	9 (0%) 81 83	11, 23, 45, 70	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	109	ARG	3.1
1	B	251	TYR	3.0
1	B	223	ALA	2.6
1	B	90	GLU	2.5
1	A	457	LYS	2.5
1	B	456	GLU	2.2
1	A	121	LYS	2.1
1	B	457	LYS	2.0
1	A	120	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	1	12/12	0.95	0.07	11,16,29,39	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	C	1	12/12	0.96	0.06	10,14,22,37	0
2	GLC	D	3	11/12	0.96	0.06	17,19,23,28	0
2	GLC	D	2	11/12	0.97	0.04	7,11,14,17	0
2	GLC	C	3	11/12	0.98	0.04	12,14,16,18	0
2	GLC	C	2	11/12	0.98	0.04	6,10,12,13	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	501	6/6	0.96	0.08	15,30,34,38	0
3	GOL	B	501	6/6	0.97	0.09	16,26,31,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.